



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 05:05 AM GMT

PDB ID : 2PBI
Title : The multifunctional nature of Gbeta5/RGS9 revealed from its crystal structure
Authors : Cheever, M.L.; Snyder, J.T.; Gershburg, S.; Siderovski, D.P.; Harden, T.K.; Sondek, J.
Deposited on : 2007-03-28
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	904	-	-	-	X
3	GOL	A	905	-	-	-	X
3	GOL	A	906	-	-	-	X
3	GOL	B	902	-	-	-	X
3	GOL	B	909	-	-	-	X
3	GOL	C	907	-	-	-	X
3	GOL	D	901	-	-	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 13309 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulator of G-protein signaling 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	415	Total	C	N	O	S	0	7	0
			3484	2238	598	631	17			

Mol	Chain	Residues	Total	C	N	O	S	ZeroOcc	AltConf	Trace
1	C	415	Total	C	N	O	S	0	8	0
			3488	2242	595	632	19			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	EXPRESSION TAG	UNP A1L352
A	0	ALA	-	EXPRESSION TAG	UNP A1L352
C	-1	GLY	-	EXPRESSION TAG	UNP A1L352
C	0	ALA	-	EXPRESSION TAG	UNP A1L352

- Molecule 2 is a protein called Guanine nucleotide-binding protein subunit beta 5.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	345	Total	C	N	O	S	0	3	0
			2664	1657	464	519	24			

Mol	Chain	Residues	Total	C	N	O	S	ZeroOcc	AltConf	Trace
2	D	354	Total	C	N	O	S	0	4	0
			2729	1695	474	535	25			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	GLY	-	EXPRESSION TAG	UNP P62881
D	0	GLY	-	EXPRESSION TAG	UNP P62881

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	B	1	Total C O 6 3 3	0	0
3	C	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	276	Total O 276 276	0	0
4	B	214	Total O 214 214	0	0
4	C	199	Total O 199 199	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	201	Total O 201 201	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Regulator of G-protein signaling 9



- Molecule 1: Regulator of G-protein signaling 9




- Molecule 2: Guanine nucleotide-binding protein subunit beta 5



- Molecule 2: Guanine nucleotide-binding protein subunit beta 5





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	110.17 Å 119.03 Å 131.95 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.95 44.19 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (20.00-1.95) 99.7 (44.19-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.65 (at 1.95 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.183 , 0.225 0.207 , 0.232	Depositor DCC
R_{free} test set	6304 reflections (5.02%)	DCC
Wilson B-factor (Å ²)	30.2	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 56.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	1 of 125589 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13309	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/3593	0.57	2/4852 (0.0%)
1	C	0.44	0/3599	0.53	0/4859
2	B	0.46	0/2727	0.63	0/3685
2	D	0.52	0/2795	0.62	0/3776
All	All	0.48	0/12714	0.58	2/17172 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	207	ARG	NE-CZ-NH1	7.71	124.15	120.30
1	A	207	ARG	NE-CZ-NH2	-6.68	116.96	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3484	0	3478	50	0
1	C	3488	0	3482	15	0
2	B	2664	0	2573	17	0
2	D	2729	0	2637	12	0
3	A	18	0	24	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	12	0	16	1	0
3	C	6	0	8	0	0
3	D	18	0	24	0	0
4	A	276	0	0	1	0
4	B	214	0	0	3	0
4	C	199	0	0	0	0
4	D	201	0	0	2	0
All	All	13309	0	12242	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:LYS:HB2	1:A:135:ILE:CG2	1.82	1.10
1:A:132:LYS:CB	1:A:135:ILE:HG22	1.82	1.09
1:A:314:GLN:HE21	1:A:318:LYS:HE3	1.17	1.06
2:B:64:ASN:HD22	2:B:65:LYS:H	1.10	0.98
1:A:132:LYS:HB3	1:A:135:ILE:HG22	1.49	0.93
2:D:64:ASN:HD22	2:D:65:LYS:H	1.09	0.93
1:A:132:LYS:HB2	1:A:135:ILE:HG22	1.45	0.89
1:A:132:LYS:HD3	1:A:135:ILE:HB	1.55	0.89
1:C:214:VAL:O	1:C:214:VAL:HG13	1.73	0.87
1:A:132:LYS:CB	1:A:135:ILE:CG2	2.51	0.82
1:A:314:GLN:NE2	1:A:318:LYS:HE3	1.93	0.82
1:C:214:VAL:O	1:C:214:VAL:CG1	2.30	0.80
1:A:135:ILE:HG12	1:A:135:ILE:O	1.85	0.76
1:C:283:LYS:HG2	1:C:307:PRO:HG2	1.71	0.73
1:A:340:GLN:OE1	1:A:343:VAL:HG11	1.89	0.72
2:D:64:ASN:HD22	2:D:65:LYS:N	1.87	0.71
1:A:215:LYS:H	1:A:215:LYS:HD3	1.57	0.70
2:B:64:ASN:HD22	2:B:65:LYS:N	1.89	0.69
1:A:129:ASN:ND2	1:A:133:LYS:O	2.25	0.68
1:A:132:LYS:CD	1:A:135:ILE:HB	2.24	0.67
1:A:132:LYS:HB2	1:A:135:ILE:HG21	1.76	0.66
1:A:132:LYS:HG3	1:A:133:LYS:H	1.61	0.65
2:B:86:LYS:HE2	2:B:100:ALA:HB1	1.78	0.65
1:A:207:ARG:HD2	4:B:973:HOH:O	1.97	0.64
1:C:210:ASN:HB3	1:C:213:GLU:HB3	1.80	0.62
2:D:64:ASN:ND2	2:D:65:LYS:H	1.90	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:51:PHE:HE2	2:B:321:LEU:HD21	1.65	0.61
1:C:90:LYS:HD3	1:C:189:LEU:HD21	1.86	0.58
1:A:132:LYS:CG	1:A:135:ILE:HB	2.34	0.57
1:A:134:GLY:C	1:A:136:LEU:H	2.08	0.57
1:A:132:LYS:HB2	1:A:135:ILE:CB	2.36	0.56
2:D:109:MET:HB2	4:D:1048:HOH:O	2.06	0.56
2:B:64:ASN:ND2	2:B:65:LYS:H	1.92	0.56
2:B:77:ARG:HD3	2:B:91:ASP:HA	1.88	0.55
1:C:210:ASN:O	1:C:213:GLU:HG2	2.07	0.54
1:A:231:GLN:OE1	1:A:235:ARG:NE	2.41	0.54
1:A:39:GLN:HE22	1:A:43:VAL:HB	1.72	0.54
2:D:152:HIS:HE1	2:D:170:THR:OG1	1.90	0.54
1:A:21:GLU:HG2	1:A:109:TRP:CZ3	2.43	0.54
1:A:132:LYS:HG2	1:A:135:ILE:N	2.22	0.54
2:B:86:LYS:CE	2:B:100:ALA:HB1	2.38	0.53
1:A:135:ILE:CG1	1:A:135:ILE:O	2.56	0.53
1:C:315:TYR:HA	1:C:318:LYS:HE3	1.91	0.52
2:D:318:VAL:HG13	4:D:980:HOH:O	2.10	0.52
1:A:355:LEU:HD23	1:A:363:ILE:HD11	1.92	0.52
1:C:407:SER:O	1:C:411:LYS:HG2	2.10	0.52
1:A:132:LYS:HB3	1:A:135:ILE:H	1.75	0.51
2:B:168:ILE:HG22	2:B:180:TRP:HB2	1.92	0.51
1:A:210:ASN:ND2	1:A:213:GLU:HG3	2.26	0.51
1:A:239:LYS:HA	2:B:33:LEU:HB3	1.91	0.50
1:A:230:GLN:O	1:A:234:MET:HG2	2.12	0.49
2:D:34:HIS:HE1	2:D:270:ALA:O	1.96	0.49
1:A:355:LEU:HD23	1:A:363:ILE:CD1	2.44	0.47
1:A:132:LYS:CG	1:A:133:LYS:H	2.28	0.47
1:A:340:GLN:OE1	1:A:343:VAL:CG1	2.61	0.47
1:C:301:SER:O	1:C:305:ARG:HG3	2.15	0.47
1:A:355:LEU:CD2	1:A:363:ILE:HD11	2.44	0.47
1:A:127:LYS:HG2	1:A:131:LYS:HE3	1.97	0.47
1:C:339:ASP:OD2	2:D:166:MET:HG2	2.15	0.47
1:C:132:LYS:HB3	1:C:135:ILE:HB	1.97	0.47
1:A:132:LYS:HG2	1:A:135:ILE:H	1.80	0.46
1:A:134:GLY:C	1:A:136:LEU:N	2.69	0.46
1:A:132:LYS:HG3	1:A:133:LYS:N	2.29	0.46
1:C:23:LEU:O	1:C:27:MET:HG3	2.16	0.46
2:B:51:PHE:CE2	2:B:321:LEU:HD21	2.48	0.45
1:A:132:LYS:CG	1:A:133:LYS:N	2.80	0.45
1:A:286:GLU:HG2	1:A:287:ILE:HG12	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:132:LYS:CB	1:A:135:ILE:CB	2.94	0.44
2:B:279:LYS:HE3	2:B:279:LYS:HB3	1.83	0.44
2:B:40:GLN:HA	2:B:43:GLU:HG3	1.99	0.44
2:B:277:TYR:HB3	2:B:310:TRP:CD2	2.53	0.43
2:D:264:ARG:HD3	2:D:273:GLU:OE2	2.18	0.43
2:D:171:ALA:HB2	2:D:201:LEU:HD23	2.00	0.43
1:A:168:THR:O	1:A:171:GLU:HG2	2.17	0.43
1:A:340:GLN:O	1:A:343:VAL:HG22	2.19	0.43
1:C:27:MET:HG2	1:C:58:TRP:CZ3	2.53	0.43
1:A:86:LEU:HD21	1:A:102:ARG:HG3	2.01	0.43
2:B:86:LYS:HE3	4:B:998:HOH:O	2.19	0.42
1:C:60:THR:HG21	1:C:67:ASN:ND2	2.34	0.42
1:A:45:SER:HB3	1:A:172:ARG:NH2	2.35	0.42
1:A:318:LYS:HD2	4:A:1099:HOH:O	2.19	0.42
2:D:277:TYR:HB3	2:D:310:TRP:CD2	2.54	0.42
2:D:284:PHE:CD2	2:D:304:ASP:HB3	2.56	0.41
2:B:228:SER:OG	2:B:230:GLN:HG2	2.21	0.41
2:B:302:TYR:CE1	2:B:308:ASN:HB2	2.55	0.41
1:A:132:LYS:CB	1:A:135:ILE:H	2.33	0.41
2:B:246:ARG:NH2	4:B:993:HOH:O	2.47	0.41
1:A:301:SER:OG	3:B:909:GOL:H32	2.21	0.40
1:A:43:VAL:HG22	1:A:175:ALA:HB1	2.03	0.40
1:A:132:LYS:HB2	1:A:135:ILE:HB	2.04	0.40
1:A:362:TRP:CH2	1:A:364:ASN:HB2	2.56	0.40
1:C:314:GLN:HG3	1:C:318:LYS:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	420/424 (99%)	406 (97%)	12 (3%)	2 (0%)	34 21

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	C	420/424 (99%)	410 (98%)	10 (2%)	0	100 100
2	B	346/354 (98%)	331 (96%)	14 (4%)	1 (0%)	46 35
2	D	356/354 (101%)	340 (96%)	16 (4%)	0	100 100
All	All	1542/1556 (99%)	1487 (96%)	52 (3%)	3 (0%)	52 43

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	420	PRO
1	A	135	ILE
2	B	162	THR

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	377/377 (100%)	360 (96%)	17 (4%)	34 18
1	C	378/377 (100%)	364 (96%)	14 (4%)	41 27
2	B	292/295 (99%)	281 (96%)	11 (4%)	40 25
2	D	299/295 (101%)	289 (97%)	10 (3%)	45 32
All	All	1346/1344 (100%)	1294 (96%)	52 (4%)	39 24

All (52) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	LEU
1	A	25	LYS
1	A	133	LYS
1	A	148	ASN
1	A	149	LYS
1	A	185	LYS
1	A	205	LEU
1	A	207	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	215	LYS
1	A	224	LYS
1	A	231	GLN
1	A	236	SER
1	A	284	LEU
1	A	286	GLU
1	A	299	ASN
1	A	363	ILE
1	A	395	LEU
2	B	12	LEU
2	B	64	ASN
2	B	77	ARG
2	B	113	TYR
2	B	151	MET
2	B	168	ILE
2	B	183	GLU
2	B	187	LEU
2	B	197	ASP
2	B	208	THR
2	B	305	TYR
1	C	44	THR
1	C	61	GLN
1	C	63	LEU
1	C	73	LEU
1	C	128	ARG
1	C	189	LEU
1	C	284	LEU
1	C	299	ASN
1	C	319	LYS
1	C	349	GLU
1	C	353	LEU
1	C	360	ARG
1	C	386	ASP
1	C	412	GLU
2	D	1	MET
2	D	64	ASN
2	D	109	MET
2	D	113	TYR
2	D	145	LYS
2	D	147	LYS
2	D	154	ASN
2	D	197	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	305	TYR
2	D	317	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	A	39	GLN
1	A	129	ASN
1	A	299	ASN
1	A	314	GLN
2	B	64	ASN
1	C	61	GLN
1	C	67	ASN
1	C	161	GLN
1	C	299	ASN
2	D	34	HIS
2	D	39	HIS
2	D	64	ASN
2	D	152	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	904	-	5,5,5	0.35	0	5,5,5	0.45	0
3	GOL	A	905	-	5,5,5	0.30	0	5,5,5	0.28	0
3	GOL	A	906	-	5,5,5	0.42	0	5,5,5	0.38	0
3	GOL	B	902	-	5,5,5	0.37	0	5,5,5	0.34	0
3	GOL	B	909	-	5,5,5	0.35	0	5,5,5	0.17	0
3	GOL	C	907	-	5,5,5	0.34	0	5,5,5	0.23	0
3	GOL	D	901	-	5,5,5	0.34	0	5,5,5	0.30	0
3	GOL	D	903	-	5,5,5	0.31	0	5,5,5	0.50	0
3	GOL	D	908	-	5,5,5	0.30	0	5,5,5	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	904	-	-	0/4/4/4	0/0/0/0
3	GOL	A	905	-	-	0/4/4/4	0/0/0/0
3	GOL	A	906	-	-	0/4/4/4	0/0/0/0
3	GOL	B	902	-	-	0/4/4/4	0/0/0/0
3	GOL	B	909	-	-	0/4/4/4	0/0/0/0
3	GOL	C	907	-	-	0/4/4/4	0/0/0/0
3	GOL	D	901	-	-	0/4/4/4	0/0/0/0
3	GOL	D	903	-	-	0/4/4/4	0/0/0/0
3	GOL	D	908	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	909	GOL	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	415/424 (97%)	0.36	20 (4%) 34 45	3, 30, 71, 128	0
1	C	415/424 (97%)	0.77	49 (11%) 6 10	11, 40, 132, 191	0
2	B	345/354 (97%)	0.06	9 (2%) 59 69	-1, 17, 69, 104	0
2	D	354/354 (100%)	0.06	12 (3%) 49 60	1, 19, 58, 155	0
All	All	1529/1556 (98%)	0.33	90 (5%) 26 36	-1, 27, 91, 191	0

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	135	ILE	8.7
1	C	214	VAL	8.5
1	C	42	LEU	7.3
2	D	0	GLY	7.0
1	C	135	ILE	6.4
1	A	214	VAL	6.0
1	A	216	LYS	5.9
1	C	155	TRP	5.5
1	C	132	LYS	5.4
1	C	133	LYS	5.2
1	A	132	LYS	4.8
2	B	209	GLY	4.7
2	D	2	ALA	4.6
1	A	133	LYS	4.4
1	A	178	TYR	4.3
1	C	144	TYR	4.1
2	D	1	MET	3.9
1	A	421	GLN	3.8
1	C	421[A]	GLN	3.8
1	C	156	ASP	3.8
2	B	138	LYS	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	149	LYS	3.7
1	C	40	ARG	3.6
1	C	157	PHE	3.6
1	C	166	TYR	3.5
1	C	158	VAL	3.4
1	C	159	ILE	3.4
2	D	3	THR	3.3
1	C	41	VAL	3.3
1	A	107	TYR	3.3
1	C	146	PHE	3.2
1	C	131	LYS	3.2
1	C	130	ILE	3.2
2	D	136	PHE	3.1
1	A	134	GLY	3.1
2	D	39	HIS	3.1
1	A	131	LYS	3.1
1	A	345	GLU	3.1
2	B	14	SER	3.0
1	C	213	GLU	3.0
1	C	136	LEU	2.9
1	C	178	TYR	2.9
2	D	138	LYS	2.8
1	A	152[A]	ASN	2.8
2	D	44	ARG	2.8
1	C	147	LEU	2.7
1	C	153	TYR	2.7
1	A	215	LYS	2.7
1	C	64	TRP	2.7
1	C	154	LYS	2.6
1	C	39	GLN	2.6
1	C	43	VAL	2.6
1	C	216	LYS	2.6
1	C	141	LYS	2.6
1	A	342	LYS	2.5
2	B	51	PHE	2.5
2	B	13	ALA	2.5
1	C	148	ASN	2.5
2	B	10	GLU	2.5
1	C	31	GLU	2.4
1	C	164	GLU	2.4
1	A	223	ARG	2.4
1	A	227	MET	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	126	ALA	2.4
1	C	345	GLU	2.4
2	D	139	ASN	2.4
1	A	234	MET	2.4
2	D	33	LEU	2.3
1	C	321	PHE	2.3
1	C	38	ASN	2.3
1	C	61	GLN	2.3
2	B	136	PHE	2.3
1	C	160	MET	2.3
1	A	31	GLU	2.3
2	B	12	LEU	2.2
1	C	46	VAL	2.2
2	D	4	ASP	2.2
1	C	167	ARG	2.2
1	C	30	PRO	2.1
1	C	49	ALA	2.1
1	C	36	MET	2.1
2	B	9	ASN	2.0
1	C	151	ILE	2.0
1	C	56	LEU	2.0
2	D	52	VAL	2.0
1	A	171	GLU	2.0
1	A	38	ASN	2.0
1	C	349	GLU	2.0
1	C	145	ASP	2.0
1	C	219	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	GOL	A	906	6/6	0.86	0.44	21.38	42,45,46,46	0
3	GOL	B	902	6/6	0.96	0.31	13.23	22,25,26,27	0
3	GOL	D	901	6/6	0.92	0.22	8.46	27,28,28,29	0
3	GOL	B	909	6/6	0.76	0.29	7.69	76,77,77,77	0
3	GOL	C	907	6/6	0.63	0.31	4.97	69,69,70,71	0
3	GOL	A	905	6/6	0.70	0.28	3.48	49,51,51,51	0
3	GOL	A	904	6/6	0.76	0.28	2.55	51,52,53,53	0
3	GOL	D	903	6/6	0.88	0.16	1.72	47,50,50,51	0
3	GOL	D	908	6/6	0.86	0.14	0.57	52,53,53,53	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.