



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:31 PM GMT

PDB ID : 1PG6  
Title : X-Ray Crystal Structure of Protein SPYM3\_0169 from Streptococcus pyogenes. Northeast Structural Genomics Consortium Target DR2.  
Authors : Kuzin, A.; Lee, I.; Edstrom, W.; Xiao, R.; Acton, T.; Rost, B.; Montelione, G.; Hunt, J.; Tong, L.; Northeast Structural Genomics Consortium (NESG)  
Deposited on : 2003-05-27  
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	243	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CA	A	237	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 1834 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hypothetical protein SpyM3\_0169.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	216	Total	C	N	O	Se	0	0	0
			1650	1037	284	319	10			

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	LEU	-	EXPRESSION TAG	UNP Q8P2Q3
A	-5	GLU	-	EXPRESSION TAG	UNP Q8P2Q3
A	-4	HIS	-	EXPRESSION TAG	UNP Q8P2Q3
A	-3	HIS	-	EXPRESSION TAG	UNP Q8P2Q3
A	-2	HIS	-	EXPRESSION TAG	UNP Q8P2Q3
A	-1	HIS	-	EXPRESSION TAG	UNP Q8P2Q3
A	0	HIS	-	EXPRESSION TAG	UNP Q8P2Q3
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	6	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	14	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	36	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	67	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	73	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	79	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	124	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	143	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	169	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3
A	194	MSE	MET	MODIFIED RESIDUE	UNP Q8P2Q3

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Ca	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	182	Total 182	O 182	0	0



● Molecule 1: Hypothetical protein SpyM3 0169



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.54Å 71.54Å 216.93Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.99 – 1.70 19.99 – 1.70	Depositor EDS
% Data completeness (in resolution range)	90.0 (19.99-1.70) 98.5 (19.99-1.70)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.86 (at 1.70Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.213 , 0.244 0.222 , 0.251	Depositor DCC
$R_{free}$ test set	1198 reflections (5.45%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtriage
Anisotropy	0.263	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 55.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 45195 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	1834	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.66% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1663	0.62	1/2220 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	59	LEU	N-CA-C	-6.56	93.28	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1650	0	1630	33	1
2	A	2	0	0	0	0
3	A	182	0	0	2	0
All	All	1834	0	1630	33	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:LEU:HD12	1:A:59:LEU:HD21	1.64	0.79
1:A:180:SER:HB3	1:A:182:GLU:HG2	1.74	0.70
1:A:156:PHE:H	1:A:216:GLN:NE2	1.91	0.69
1:A:66:SER:HB3	1:A:73:MSE:HE3	1.77	0.65
1:A:58:LYS:HA	1:A:58:LYS:HE2	1.77	0.64
1:A:91:ASN:OD1	1:A:92:THR:HG23	2.01	0.60
1:A:95:GLN:NE2	1:A:159:ILE:H	2.00	0.58
1:A:59:LEU:HD23	1:A:60:VAL:HG23	1.86	0.58
1:A:208:ARG:HD3	3:A:466:HOH:O	2.03	0.58
1:A:6:MSE:HG2	1:A:117:ASP:OD2	2.04	0.57
1:A:66:SER:CB	1:A:73:MSE:HE3	2.34	0.57
1:A:73:MSE:HE2	1:A:73:MSE:HA	1.88	0.56
1:A:107:ARG:HA	1:A:143:MSE:O	2.08	0.54
1:A:56:LEU:HA	1:A:59:LEU:HD22	1.91	0.53
1:A:128:ASN:O	1:A:129:ILE:HG22	2.10	0.52
1:A:193:PHE:O	1:A:197:ILE:HG12	2.11	0.51
1:A:48:LYS:HG2	1:A:73:MSE:HE1	1.96	0.48
1:A:77:GLN:HG2	1:A:79:MSE:SE	2.64	0.48
1:A:180:SER:HB3	1:A:182:GLU:CG	2.42	0.48
1:A:42:ASN:HD22	1:A:42:ASN:N	2.10	0.48
1:A:14:MSE:HE1	1:A:18:LEU:HB2	1.97	0.46
1:A:63:ILE:O	1:A:67:MSE:HG3	2.16	0.46
1:A:50:ASN:HB3	3:A:369:HOH:O	2.15	0.46
1:A:59:LEU:CD2	1:A:60:VAL:HG23	2.45	0.45
1:A:59:LEU:HD23	1:A:60:VAL:N	2.31	0.45
1:A:58:LYS:C	1:A:61:GLY:H	2.21	0.43
1:A:57:GLY:H	1:A:59:LEU:CD2	2.32	0.43
1:A:48:LYS:CG	1:A:73:MSE:HE1	2.49	0.42
1:A:42:ASN:ND2	1:A:42:ASN:N	2.67	0.42
1:A:108:LEU:HD12	1:A:108:LEU:N	2.34	0.42
1:A:57:GLY:H	1:A:59:LEU:HD22	1.84	0.41
1:A:57:GLY:N	1:A:59:LEU:HD22	2.36	0.41
1:A:59:LEU:HD23	1:A:59:LEU:H	1.86	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:15:GLN:OE1	1:A:15:GLN:OE1[3_665]	1.99	0.21



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/243 (86%)	206 (98%)	3 (1%)	1 (0%)	34	15

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	175/186 (94%)	170 (97%)	5 (3%)	50	27

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	31	LEU
1	A	42	ASN
1	A	81	ASN
1	A	108	LEU
1	A	117	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	ASN
1	A	15	GLN
1	A	32	GLN
1	A	33	GLN
1	A	42	ASN
1	A	81	ASN
1	A	95	GLN
1	A	109	ASN
1	A	121	GLN
1	A	128	ASN
1	A	173	ASN
1	A	195	GLN
1	A	210	HIS
1	A	216	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	206/243 (84%)	0.99	30 (14%) 3 4	8, 22, 54, 63	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	58	LYS	10.0
1	A	128	ASN	9.0
1	A	138	GLY	8.6
1	A	129	ILE	8.3
1	A	50	ASN	6.9
1	A	68	VAL	6.6
1	A	56	LEU	5.5
1	A	59	LEU	5.3
1	A	130	GLY	5.0
1	A	60	VAL	4.7
1	A	191	ASN	4.4
1	A	5	ARG	4.2
1	A	193	PHE	4.1
1	A	41	GLU	3.9
1	A	57	GLY	3.7
1	A	231	LEU	3.5
1	A	46	ASN	3.4
1	A	189	LEU	3.4
1	A	139	GLY	3.2
1	A	110	ASP	2.9
1	A	65	ARG	2.8
1	A	192	GLY	2.7
1	A	165	ASP	2.6
1	A	63	ILE	2.6
1	A	55	GLY	2.5
1	A	103	GLU	2.2
1	A	7	ARG	2.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	123	LYS	2.1
1	A	121	GLN	2.1
1	A	208	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	237	1/1	0.92	0.25	9.61	31,31,31,31	0
2	CA	A	238	1/1	0.99	0.04	-2.66	10,10,10,10	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.