



Full wwPDB X-ray Structure Validation Report i

Jan 31, 2016 – 09:32 PM GMT

PDB ID : 1PH6

Title : Crystal Structure of THE OXYTRICHA NOVA TELOMERE END-BINDING PROTEIN COMPLEXED WITH NONCOGNATE SSDNA GGGGTTTTGTGG

Authors : Theobald, D.L.; Schultz, S.C.

Deposited on : 2003-05-29

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.7 (RC4), CSD as536be (2015)

Xtriage (Phenix) : 1.9-1692

EDS : rb-20026688

Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)

Refmac : 5.8.0135

CCP4 : 6.5.0

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition [\(i\)](#)

There are 6 unique types of molecules in this entry. The entry contains 6448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*GP*GP*GP*GP*TP*TP*TP*TP*GP*GP*GP*G P*T)-3'.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	G	12	Total	C	N	O	P	0	0
			253	120	48	74	11		

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
1	H	12	Total	C	N	O	P	0	0
			253	120	48	74	11		

- Molecule 2 is a DNA chain called 5'-D(*GP*GP*GP*GP*TP*TP*TP*TP*GP*TP*GP*G -3'.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	D	11	Total	C	N	O	P	0	0
			229	110	40	69	10		

- Molecule 3 is a protein called Telomere-binding protein alpha subunit.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	A	452	Total	C	N	O	S	0	0
			3665	2333	628	702	2		

- Molecule 4 is a protein called Telomere-binding protein beta subunit.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	B	217	Total	C	N	O	S	0	0
			1740	1117	295	327	1		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Na	0	0
			5	5		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	230	Total O 230 230	0	0
6	B	45	Total O 45 45	0	0
6	D	18	Total O 18 18	0	0
6	G	5	Total O 5 5	0	0
6	H	5	Total O 5 5	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(*GP*GP*GP*GP*TP*TP*TP*TP*GP*GP*GP*GP*T)-3'




- Molecule 1: 5'-D(*GP*GP*GP*GP*TP*TP*TP*TP*GP*GP*GP*GP*T)-3'




- Molecule 2: 5'-D(*GP*GP*GP*GP*TP*TP*TP*TP*GP*TP*GP*G)-3'




- Molecule 3: Telomere-binding protein alpha subunit



- Molecule 4: Telomere-binding protein beta subunit





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	93.47Å 93.47Å 423.94Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.90 – 2.10 29.90 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.5 (29.90-2.10) 90.8 (29.90-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	1.56 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R , R_{free}	0.242 , 0.276 0.242 , 0.277	Depositor DCC
R_{free} test set	5996 reflections (9.91%)	DCC
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.302	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 40.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	1 of 68597 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	6448	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	G	0.54	0/284	0.73	0/440
1	H	0.45	0/284	0.65	0/440
2	D	0.73	0/256	0.83	0/396
3	A	0.55	0/3736	0.73	3/5050 (0.1%)
4	B	0.40	0/1779	0.54	0/2402
All	All	0.52	0/6339	0.68	3/8728 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	A	86	LYS	N-CA-C	-6.82	92.58	111.00
3	A	406	LYS	N-CA-C	5.60	126.11	111.00
3	A	266	ARG	NE-CZ-NH2	-5.23	117.69	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	253	0	138	7	0
1	H	253	0	138	1	0
2	D	229	0	128	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3665	0	3664	64	0
4	B	1740	0	1726	47	0
5	A	5	0	0	0	0
6	A	230	0	0	1	0
6	B	45	0	0	1	0
6	D	18	0	0	0	0
6	G	5	0	0	0	0
6	H	5	0	0	0	0
All	All	6448	0	5794	123	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (123) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:80:ILE:HD12	4:B:218:ILE:HD11	1.52	0.91
4:B:10:GLN:HE21	4:B:12:SER:H	1.19	0.91
1:G:11:DG:H2'	1:G:12:DG:H5'	1.53	0.89
2:D:6:DT:H2'	2:D:7:DT:H5'	1.57	0.85
2:D:6:DT:H2'	2:D:7:DT:C5'	2.17	0.75
2:D:8:DT:H2'	2:D:9:DG:H5'	1.68	0.74
3:A:226:VAL:HG13	3:A:273:ILE:HB	1.71	0.73
4:B:125:LEU:HD12	4:B:125:LEU:H	1.52	0.73
4:B:40:LYS:HD3	4:B:128:ARG:NH2	2.03	0.71
4:B:40:LYS:HD3	4:B:128:ARG:HH21	1.56	0.70
3:A:400:LEU:HD13	3:A:450:ASN:HA	1.74	0.69
4:B:62:TYR:HB3	4:B:212:THR:HG21	1.77	0.66
4:B:10:GLN:HE21	4:B:12:SER:N	1.94	0.66
4:B:132:THR:HG22	4:B:133:ARG:O	1.95	0.66
4:B:48:PRO:O	4:B:215:ILE:HG21	1.97	0.65
3:A:389:GLY:HA3	3:A:407:GLY:HA3	1.76	0.65
3:A:345:LEU:HD12	3:A:346:PRO:HD2	1.78	0.65
3:A:285:LYS:HB3	3:A:331:ASN:HD21	1.61	0.65
3:A:71:ARG:HH21	3:A:103:TYR:HE2	1.45	0.64
3:A:399:SER:O	3:A:401:LYS:N	2.29	0.64
2:D:12:DG:H2'	2:D:12:DG:N3	2.12	0.64
3:A:46:LEU:HD23	3:A:127:LEU:HG	1.81	0.63
3:A:399:SER:C	3:A:401:LYS:H	2.00	0.62
3:A:155:SER:H	3:A:158:GLN:NE2	1.97	0.61
4:B:28:PHE:O	4:B:31:VAL:HG12	2.00	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:6:DT:H2”	1:G:8:DT:O4	2.03	0.59
3:A:423:GLN:HG3	6:A:707:HOH:O	2.03	0.59
3:A:76:LEU:HD23	3:A:76:LEU:N	2.18	0.58
3:A:401:LYS:O	3:A:406:LYS:HB2	2.03	0.58
4:B:215:ILE:O	4:B:218:ILE:HG22	2.04	0.57
4:B:126:GLN:HE21	4:B:128:ARG:HD3	1.68	0.57
4:B:195:SER:HB3	6:B:256:HOH:O	2.03	0.57
3:A:71:ARG:NE	3:A:73:ILE:HD11	2.21	0.56
3:A:67:THR:HG21	3:A:73:ILE:HD12	1.88	0.56
4:B:10:GLN:NE2	4:B:12:SER:H	1.98	0.55
3:A:339:ASP:OD1	3:A:341:LYS:HG2	2.07	0.55
3:A:281:THR:HB	4:B:10:GLN:HE22	1.72	0.54
3:A:387:VAL:HG12	3:A:400:LEU:HD12	1.89	0.54
2:D:6:DT:C2’	2:D:7:DT:H5”	2.32	0.54
3:A:86:LYS:N	3:A:96:ASP:HB2	2.22	0.54
4:B:81:VAL:HG23	4:B:222:GLU:OE1	2.08	0.54
4:B:80:ILE:CD1	4:B:218:ILE:HD11	2.33	0.53
3:A:285:LYS:CB	3:A:331:ASN:HD21	2.22	0.53
3:A:336:THR:HG22	3:A:493:LEU:HD12	1.91	0.53
1:H:4:DG:H1’	1:H:5:DT:H5’	1.90	0.52
3:A:400:LEU:HD11	3:A:409:ASN:OD1	2.08	0.52
1:G:6:DT:H2”	1:G:8:DT:C4	2.45	0.51
3:A:479:VAL:HG12	3:A:488:ILE:HD13	1.93	0.51
4:B:57:PHE:CD2	4:B:136:VAL:HG23	2.46	0.51
4:B:44:LYS:HG2	4:B:45:GLU:HG3	1.93	0.50
3:A:155:SER:H	3:A:158:GLN:HE21	1.59	0.50
3:A:306:ALA:O	3:A:310:ARG:HB2	2.12	0.50
4:B:129:LEU:O	4:B:130:ASN:CB	2.61	0.49
4:B:125:LEU:N	4:B:125:LEU:HD12	2.26	0.49
3:A:400:LEU:C	3:A:406:LYS:H	2.14	0.49
3:A:61:ALA:HB2	3:A:76:LEU:HB3	1.95	0.48
3:A:391:ASP:CG	3:A:394:THR:HG22	2.35	0.47
4:B:47:TYR:CD1	4:B:48:PRO:HA	2.49	0.47
3:A:85:LEU:C	3:A:86:LYS:O	2.49	0.47
4:B:92:ASN:HD22	4:B:92:ASN:C	2.18	0.47
3:A:399:SER:C	3:A:401:LYS:N	2.67	0.47
4:B:81:VAL:O	4:B:81:VAL:HG12	2.15	0.47
3:A:414:GLN:OE1	4:B:170:VAL:HG23	2.15	0.47
4:B:44:LYS:HE3	4:B:45:GLU:OE2	2.14	0.47
3:A:458:ARG:O	3:A:462:GLU:HG3	2.15	0.46
4:B:87:VAL:CG1	4:B:128:ARG:HG3	2.45	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:149:PHE:CZ	3:A:171:PRO:HG3	2.51	0.46
4:B:87:VAL:CG1	4:B:124:ASN:HB2	2.45	0.46
3:A:85:LEU:HG	3:A:86:LYS:O	2.15	0.46
3:A:391:ASP:HB3	3:A:394:THR:HG22	1.98	0.46
3:A:432:LEU:HB2	3:A:487:LEU:HD23	1.98	0.45
1:G:9:DG:N3	1:G:9:DG:H2'	2.30	0.45
3:A:200:PHE:CZ	3:A:205:VAL:HG11	2.51	0.45
3:A:450:ASN:OD1	3:A:452:HIS:HB2	2.16	0.45
3:A:79:VAL:HG13	3:A:84:TYR:HB3	1.98	0.45
4:B:90:ILE:HG23	4:B:118:VAL:HG13	1.98	0.45
3:A:215:ASN:HA	3:A:251:GLN:HG3	1.99	0.45
4:B:62:TYR:HB3	4:B:212:THR:CG2	2.45	0.45
3:A:400:LEU:HA	3:A:400:LEU:HD23	1.64	0.44
3:A:351:GLN:O	3:A:355:HIS:HB2	2.16	0.44
3:A:61:ALA:CB	3:A:76:LEU:HB3	2.48	0.44
3:A:400:LEU:CD1	3:A:450:ASN:HA	2.44	0.44
2:D:8:DT:H2"	2:D:9:DG:C5'	2.44	0.44
2:D:6:DT:O2	2:D:8:DT:H73	2.18	0.44
4:B:42:TYR:CE1	4:B:85:ASP:HA	2.53	0.44
3:A:71:ARG:HE	3:A:73:ILE:HD11	1.81	0.44
4:B:206:LYS:O	4:B:206:LYS:HG3	2.18	0.43
3:A:422:THR:HB	3:A:425:ASN:HD21	1.83	0.43
3:A:79:VAL:CG1	3:A:80:ASP:N	2.81	0.43
4:B:129:LEU:O	4:B:130:ASN:HB2	2.18	0.43
1:G:4:DG:OP1	3:A:105:LYS:NZ	2.50	0.43
4:B:122:LYS:HA	4:B:123:PRO:HD3	1.82	0.43
4:B:41:CYS:HB2	4:B:53:THR:O	2.18	0.43
3:A:85:LEU:HA	3:A:96:ASP:H	1.84	0.43
1:G:4:DG:H1'	1:G:5:DT:H5'	2.00	0.43
3:A:379:GLU:HA	3:A:380:PRO:C	2.39	0.43
3:A:230:ILE:HD13	3:A:245:LEU:CD2	2.49	0.43
4:B:55:GLY:O	4:B:132:THR:HG23	2.18	0.43
4:B:98:ARG:HG3	4:B:115:ARG:HD2	2.01	0.42
3:A:418:LYS:HD3	4:B:156:LEU:HD21	2.00	0.42
1:G:11:DG:C2'	1:G:12:DG:H5'	2.34	0.42
3:A:387:VAL:HG12	3:A:400:LEU:CD1	2.49	0.42
4:B:132:THR:HG22	4:B:133:ARG:N	2.35	0.42
3:A:434:TYR:HB3	3:A:436:GLN:OE1	2.20	0.42
3:A:121:ARG:O	3:A:146:TRP:HA	2.20	0.42
4:B:17:LEU:HD11	4:B:35:LEU:HB3	2.01	0.42
4:B:216:ALA:O	4:B:220:VAL:HG23	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:87:VAL:HG11	4:B:128:ARG:HG3	2.02	0.41
4:B:212:THR:HG22	4:B:213:LEU:N	2.34	0.41
4:B:55:GLY:HA2	4:B:132:THR:HG21	2.03	0.41
3:A:48:SER:C	3:A:50:GLN:H	2.23	0.41
4:B:132:THR:CG2	4:B:133:ARG:N	2.84	0.41
3:A:348:THR:CG2	3:A:353:LEU:HD13	2.51	0.41
3:A:285:LYS:HD3	3:A:331:ASN:ND2	2.35	0.41
2:D:6:DT:H2"	2:D:7:DT:H5'	1.99	0.41
3:A:400:LEU:HD13	3:A:450:ASN:CA	2.45	0.41
3:A:366:GLN:HB3	3:A:369:PHE:CZ	2.56	0.41
4:B:214:LYS:HB2	4:B:217:ASP:OD2	2.20	0.41
4:B:86:LYS:NZ	4:B:125:LEU:HD23	2.36	0.40
3:A:320:GLU:O	3:A:324:LEU:HG	2.21	0.40
3:A:444:PHE:O	3:A:445:ASN:HB2	2.21	0.40
3:A:73:ILE:HG22	3:A:74:CYS:N	2.36	0.40
3:A:149:PHE:CE2	3:A:171:PRO:HG3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	450/461 (98%)	429 (95%)	16 (4%)	5 (1%)	17 11
4	B	215/217 (99%)	194 (90%)	18 (8%)	3 (1%)	14 7
All	All	665/678 (98%)	623 (94%)	34 (5%)	8 (1%)	16 10

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	93	ASP
3	A	406	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
4	B	130	ASN
4	B	126	GLN
4	B	210	THR
3	A	400	LEU
3	A	49	ALA
3	A	68	ASN

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	406/409 (99%)	400 (98%)	6 (2%)	72 78
4	B	190/190 (100%)	186 (98%)	4 (2%)	61 66
All	All	596/599 (100%)	586 (98%)	10 (2%)	68 74

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	76	LEU
3	A	226	VAL
3	A	239	TYR
3	A	309	LEU
3	A	353	LEU
3	A	493	LEU
4	B	66	GLU
4	B	92	ASN
4	B	125	LEU
4	B	215	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	158	GLN
3	A	161	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	198	GLN
3	A	327	ASN
3	A	331	ASN
3	A	372	GLN
4	B	10	GLN
4	B	92	ASN
4	B	126	GLN
4	B	149	GLN
4	B	221	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	G	12/13 (92%)	0.25	0 100 100	32, 42, 50, 53	0
1	H	12/13 (92%)	0.10	0 100 100	40, 45, 52, 56	0
2	D	11/11 (100%)	-0.29	1 (9%) 11 16	23, 31, 47, 51	0
3	A	452/461 (98%)	0.13	22 (4%) 33 42	16, 27, 54, 69	0
4	B	217/217 (100%)	0.91	31 (14%) 4 5	20, 46, 74, 89	0
All	All	704/715 (98%)	0.37	54 (7%) 16 22	16, 32, 67, 89	0

All (54) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	B	127	GLU	10.4
3	A	406	LYS	10.0
3	A	407	GLY	8.9
4	B	8	PRO	7.8
4	B	126	GLN	6.9
4	B	196	LYS	5.8
4	B	129	LEU	5.7
4	B	78	VAL	5.7
3	A	400	LEU	5.3
4	B	9	GLN	5.1
4	B	73	ALA	5.0
4	B	130	ASN	4.9
3	A	87	GLN	4.7
4	B	194	ALA	4.4
4	B	223	LYS	4.2
4	B	215	ILE	4.2
4	B	195	SER	4.1
3	A	93	ASP	4.1
3	A	70	GLU	4.1
3	A	401	LYS	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	B	219	PHE	3.8
4	B	77	ASN	3.7
3	A	94	ALA	3.5
4	B	224	GLY	3.4
4	B	218	ILE	3.4
3	A	76	LEU	3.1
3	A	69	GLN	3.1
3	A	71	ARG	3.1
4	B	179	ALA	2.9
4	B	25	GLU	2.8
3	A	183	LYS	2.8
3	A	362	GLU	2.7
4	B	125	LEU	2.6
4	B	122	LYS	2.6
4	B	197	GLY	2.5
4	B	65	LYS	2.4
4	B	213	LEU	2.4
4	B	131	PRO	2.3
4	B	128	ARG	2.3
3	A	344	ALA	2.2
3	A	78	ILE	2.2
3	A	153	LYS	2.2
3	A	120	ILE	2.2
4	B	214	LYS	2.1
2	D	8	DT	2.1
3	A	168	ASP	2.1
3	A	68	ASN	2.1
4	B	59	VAL	2.1
3	A	340	LYS	2.0
4	B	74	LYS	2.0
3	A	366	GLN	2.0
4	B	87	VAL	2.0
4	B	220	VAL	2.0
3	A	86	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	NA	A	500	1/1	0.80	0.67	-	85,85,85,85	0
5	NA	A	499	1/1	0.18	0.37	-	85,85,85,85	0
5	NA	A	496	1/1	0.36	0.98	-	99,99,99,99	0
5	NA	A	498	1/1	-0.06	1.91	-	94,94,94,94	0
5	NA	A	497	1/1	0.57	0.70	-	101,101,101,101	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.