



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:11 AM GMT

PDB ID : 2PRX
Title : Crystal structure of Thioesterase superfamily protein (ZP_00837258.1) from *Shewanella loihica* PV-4 at 1.65 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2007-05-04
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	<div> <div>19%</div> <div>58%</div> <div>11%</div> <div>•</div> <div>29%</div> </div>
1	B	160	<div> <div>21%</div> <div>58%</div> <div>11%</div> <div>•</div> <div>29%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1724 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Thioesterase superfamily protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	114	Total	C	N	O	S	Se	0	2	0
			852	549	143	153	3	4			
1	B	114	Total	C	N	O	S	Se	0	1	0
			863	555	149	152	3	4			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP A3QJH3
A	1	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
A	44	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
A	109	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
A	145	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
A	150	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
A	154	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
B	0	GLY	-	LEADER SEQUENCE	UNP A3QJH3
B	1	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
B	44	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
B	109	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
B	145	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
B	150	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3
B	154	MSE	MET	MODIFIED RESIDUE	UNP A3QJH3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	O	0	0
			3	3		
2	B	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Thioesterase superfamily protein



• Molecule 1: Thioesterase superfamily protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, α , β , γ	38.50Å 38.50Å 172.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.74 – 1.50 33.34 – 1.50	Depositor EDS
% Data completeness (in resolution range)	72.7 (28.74-1.50) 76.2 (33.34-1.50)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.48 (at 1.50Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.199 , 0.277 0.195 , 0.257	Depositor DCC
R_{free} test set	1800 reflections (5.44%)	DCC
Wilson B-factor (Å ²)	12.0	Xtriage
Anisotropy	0.410	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.4	EDS
Estimated twinning fraction	0.386 for -h,-k,l 0.317 for h,-h-k,-l 0.319 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.35$, $\langle L^2 \rangle = 0.18$	Xtriage
Outliers	0 of 35102 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	1724	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/870	1.23	5/1168 (0.4%)
1	B	0.44	0/878	1.31	4/1173 (0.3%)
All	All	0.45	0/1748	1.27	9/2341 (0.4%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	18	TYR	CB-CG-CD1	-9.68	115.19	121.00
1	A	51	ALA	C-N-CA	8.28	142.41	121.70
1	A	134	SER	C-N-CA	8.06	141.86	121.70
1	B	18	TYR	CB-CG-CD2	7.71	125.63	121.00
1	B	17	CYS	C-N-CA	6.26	137.36	121.70
1	A	35	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	A	101	ILE	C-N-CA	5.31	134.98	121.70
1	B	125	ARG	NE-CZ-NH1	5.07	122.84	120.30
1	A	17	CYS	C-N-CA	5.01	134.22	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	852	0	846	12	0
1	B	863	0	861	15	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	3	0	0	0	0
2	B	6	0	0	3	0
All	All	1724	0	1707	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:19:GLY:HA3	2:B:162:HOH:O	1.75	0.85
1:B:120:LYS:HE2	1:B:130:GLU:OE1	2.01	0.60
1:B:125:ARG:NH1	1:B:149:LYS:HD3	2.18	0.58
1:A:49:HIS:CD2	1:A:60:LEU:HD22	2.39	0.58
1:A:17:CYS:HA	2:B:164:HOH:O	2.05	0.57
1:A:118:GLU:HG3	1:A:120:LYS:CD	2.36	0.55
1:A:51:ALA:O	1:B:17:CYS:HB2	2.08	0.54
1:A:118:GLU:OE1	1:A:118:GLU:HA	2.08	0.54
1:A:100:ASN:OD1	1:B:100:ASN:HB3	2.08	0.53
1:A:94:PHE:CE2	1:A:127:VAL:HG23	2.46	0.50
1:B:40:ILE:HA	1:B:115:LEU:O	2.11	0.49
1:A:100:ASN:HB3	1:B:100:ASN:OD1	2.11	0.49
1:B:46:LYS:O	1:B:49:HIS:HB2	2.13	0.48
1:A:42:HIS:HA	1:A:113:LEU:O	2.13	0.48
1:B:125:ARG:N	2:B:163:HOH:O	2.50	0.45
1:B:130:GLU:OE2	1:B:142:ARG:NH2	2.50	0.45
1:B:104:LEU:HD23	1:B:104:LEU:N	2.32	0.44
1:A:49:HIS:CG	1:A:60:LEU:HD22	2.54	0.43
1:B:60:LEU:O	1:B:63:SER:HB2	2.19	0.42
1:A:35:ARG:HB2	1:A:40:ILE:HG22	2.02	0.42
1:B:125:ARG:O	1:B:149:LYS:N	2.50	0.41
1:A:96:THR:OG1	1:A:145:MSE:HE2	2.20	0.41
1:B:126:LYS:NZ	1:B:146:VAL:HG22	2.36	0.41
1:B:139:LEU:HD21	1:B:142:ARG:HG2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/160 (65%)	98 (94%)	6 (6%)	0	100	100
1	B	101/160 (63%)	98 (97%)	3 (3%)	0	100	100
All	All	205/320 (64%)	196 (96%)	9 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	88/117 (75%)	83 (94%)	5 (6%)	25	3
1	B	89/117 (76%)	84 (94%)	5 (6%)	26	3
All	All	177/234 (76%)	167 (94%)	10 (6%)	25	3

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	15	SER
1	A	52	ILE
1	A	93	ARG
1	A	118	GLU
1	A	121	GLU
1	B	15	SER
1	B	79	ARG
1	B	130	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	142	ARG
1	B	145	MSE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	144	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.







5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	110/160 (68%)	1.71	31 (28%)  	7, 15, 22, 29	0
1	B	110/160 (68%)	1.80	34 (30%)  	8, 15, 22, 29	0
All	All	220/320 (68%)	1.75	65 (29%)  	7, 15, 22, 29	0

All (65) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	18	TYR	8.7
1	A	18	TYR	8.6
1	B	6	PHE	6.1
1	B	99	LEU	5.6
1	B	111	VAL	5.4
1	A	6	PHE	5.1
1	B	103	TYR	4.5
1	B	75	ALA	4.2
1	B	76	ALA	4.2
1	B	115	LEU	4.2
1	B	95	VAL	3.9
1	B	19	GLY	3.7
1	B	10	TYR	3.6
1	A	35	ARG	3.4
1	A	55	PHE	3.3
1	A	51	ALA	3.3
1	A	77	ALA	3.2
1	B	101	ILE	3.1
1	B	17	CYS	3.1
1	B	122	VAL	3.0
1	A	33	TYR	2.9
1	A	135	ALA	2.9
1	A	111	VAL	2.9
1	B	70	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	17	CYS	2.9
1	B	121	GLU	2.8
1	B	33	TYR	2.8
1	A	30	LEU	2.8
1	A	75	ALA	2.8
1	B	35	ARG	2.7
1	A	151	PRO	2.7
1	A	10	TYR	2.6
1	B	144	HIS	2.6
1	B	30	LEU	2.6
1	B	78	GLN	2.6
1	A	34	TRP	2.6
1	B	79	ARG	2.6
1	A	43	PHE	2.6
1	B	146	VAL	2.5
1	B	43	PHE	2.5
1	A	60	LEU	2.4
1	B	55	PHE	2.4
1	B	94	PHE	2.4
1	B	51	ALA	2.4
1	A	57	TYR	2.4
1	A	52	ILE	2.4
1	B	142	ARG	2.4
1	A	65	ILE	2.3
1	A	67	CYS	2.3
1	B	15	SER	2.3
1	A	94	PHE	2.3
1	A	97	ALA	2.2
1	A	92	PRO	2.2
1	B	126	LYS	2.2
1	A	116	VAL	2.1
1	A	9	ALA	2.1
1	B	49	HIS	2.1
1	A	113	LEU	2.1
1	B	148	VAL	2.1
1	B	130	GLU	2.1
1	A	104	LEU	2.1
1	A	131	ILE	2.0
1	A	95	VAL	2.0
1	A	121	GLU	2.0
1	B	105	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.