



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:38 PM GMT

PDB ID : 1PTO  
Title : THE STRUCTURE OF A PERTUSSIS TOXIN-SUGAR COMPLEX AS A  
MODEL FOR RECEPTOR BINDING  
Authors : Stein, P.E.; Read, R.J.  
Deposited on : 1994-03-22  
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1157 (3.60-3.40)
Ramachandran outliers	100387	1120 (3.60-3.40)
Sidechain outliers	100360	1121 (3.60-3.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	244	
1	G	244	
2	B	196	
3	C	196	
3	I	196	
4	D	110	
4	E	110	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	J	110	 42% 55%
4	K	110	 51% 38% 11%
5	F	98	 34% 46% 19%
5	L	98	 46% 39% 15%
6	H	198	 42% 45% 12%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 14614 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PERTUSSIS TOXIN (SUBUNIT S1).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	Total	C	N	O	S	0	0	0
			1769	1095	318	350	6			
1	G	224	Total	C	N	O	S	0	0	0
			1769	1095	318	350	6			

- Molecule 2 is a protein called PERTUSSIS TOXIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	196	Total	C	N	O	S	0	0	0
			1522	961	260	292	9			

- Molecule 3 is a protein called PERTUSSIS TOXIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	196	Total	C	N	O	S	0	0	0
			1521	969	258	285	9			
3	I	196	Total	C	N	O	S	0	0	0
			1521	969	258	285	9			

- Molecule 4 is a protein called PERTUSSIS TOXIN (SUBUNIT S4).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	110	Total	C	N	O	S	0	0	0
			838	536	143	147	12			
4	E	110	Total	C	N	O	S	0	0	0
			838	536	143	147	12			
4	J	110	Total	C	N	O	S	0	0	0
			838	536	143	147	12			
4	K	110	Total	C	N	O	S	0	0	0
			838	536	143	147	12			

- Molecule 5 is a protein called PERTUSSIS TOXIN (SUBUNIT S5).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	98	Total	C	N	O	S	0	0	0
			764	489	125	144	6			
5	L	98	Total	C	N	O	S	0	0	0
			764	489	125	144	6			

- Molecule 6 is a protein called PERTUSSIS TOXIN (SUBUNIT S2).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	H	198	Total	C	N	O	S	7	0	0
			1536	970	262	295	9			

- Molecule 7 is a polymer of unknown type called SUGAR (2-MER).


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	2	Total	C	N	O	0	0
			32	17	1	14		
7	C	2	Total	C	N	O	0	0
			32	17	1	14		
7	I	2	Total	C	N	O	0	0
			32	17	1	14		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: PERTUSSIS TOXIN (SUBUNIT S1)



- Molecule 1: PERTUSSIS TOXIN (SUBUNIT S1)




- Molecule 2: PERTUSSIS TOXIN







• Molecule 3: PERTUSSIS TOXIN




• Molecule 3: PERTUSSIS TOXIN




• Molecule 4: PERTUSSIS TOXIN (SUBUNIT S4)




• Molecule 4: PERTUSSIS TOXIN (SUBUNIT S4)




- Molecule 4: PERTUSSIS TOXIN (SUBUNIT S4)



- Molecule 4: PERTUSSIS TOXIN (SUBUNIT S4)




- Molecule 5: PERTUSSIS TOXIN (SUBUNIT S5)



- Molecule 5: PERTUSSIS TOXIN (SUBUNIT S5)



- Molecule 6: PERTUSSIS TOXIN (SUBUNIT S2)



K151	M152	L153	Y154	L155	I156	V157	V158	I161	S162	V163	R164	V165	Q172	Y173	Y174	D175	Y176	E177	D178	A179	T180	F181	Y184	A185	L186	T187	G188	I189	S190	I191	C192	M193	S196	C199
------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	163.80Å 98.20Å 194.50Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 3.50	Depositor
% Data completeness (in resolution range)	(Not available) (10.00-3.50)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, $R_{free}$	0.183 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	14614	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SIA, GAL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.73	0/1809	0.85	1/2457 (0.0%)
1	G	0.53	0/1809	0.75	0/2457
2	B	0.75	0/1558	0.94	1/2115 (0.0%)
3	C	0.78	0/1557	0.90	1/2115 (0.0%)
3	I	0.69	0/1557	0.89	3/2115 (0.1%)
4	D	0.74	0/856	0.97	1/1155 (0.1%)
4	E	0.79	0/856	0.90	0/1155
4	J	0.67	0/856	0.92	1/1155 (0.1%)
4	K	0.62	0/856	0.87	1/1155 (0.1%)
5	F	0.74	0/782	0.94	1/1059 (0.1%)
5	L	0.62	0/782	0.86	0/1059
6	H	0.67	0/1573	0.92	2/2137 (0.1%)
All	All	0.70	0/14851	0.89	12/20134 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
3	I	0	1
4	D	0	1
5	F	0	1
5	L	0	1
6	H	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	119	LEU	N-CA-C	-6.30	93.98	111.00
6	H	191	ILE	N-CA-C	-5.84	95.22	111.00
3	I	119	LEU	N-CA-C	-5.83	95.27	111.00
3	I	112	LEU	CA-CB-CG	5.63	128.25	115.30
4	D	31	CYS	CA-CB-SG	-5.47	104.16	114.00

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	126	TYR	Sidechain
4	D	52	PHE	Sidechain
5	F	64	TYR	Sidechain
6	H	146	TYR	Sidechain
3	I	103	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1769	0	1655	127	0
1	G	1769	0	1655	110	0
2	B	1522	0	1473	125	0
3	C	1521	0	1484	145	0
3	I	1521	0	1484	133	0
4	D	838	0	874	54	0
4	E	838	0	874	58	0
4	J	838	0	874	62	0
4	K	838	0	874	55	0
5	F	764	0	747	60	0
5	L	764	0	747	47	0
6	H	1536	0	1487	114	0
7	B	32	0	28	1	0
7	C	32	0	28	5	0
7	I	32	0	28	7	0
All	All	14614	0	14312	998	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 998 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:H:28:ARG:HH21	6:H:177:GLU:HA	1.22	1.01
2:B:28:ARG:HH21	2:B:177:GLU:HA	1.26	0.99
3:I:60:GLY:HA2	3:I:74:ALA:HB3	1.46	0.97
1:G:51:VAL:HB	1:G:132:ALA:HB3	1.51	0.93
2:B:163:VAL:HG21	2:B:189:ILE:HG22	1.52	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/244 (90%)	198 (90%)	17 (8%)	5 (2%)	8	48
1	G	220/244 (90%)	193 (88%)	21 (10%)	6 (3%)	6	44
2	B	194/196 (99%)	154 (79%)	30 (16%)	10 (5%)	2	25
3	C	194/196 (99%)	171 (88%)	17 (9%)	6 (3%)	5	41
3	I	194/196 (99%)	170 (88%)	19 (10%)	5 (3%)	7	45
4	D	108/110 (98%)	96 (89%)	10 (9%)	2 (2%)	10	51
4	E	108/110 (98%)	91 (84%)	15 (14%)	2 (2%)	10	51
4	J	108/110 (98%)	100 (93%)	7 (6%)	1 (1%)	21	68
4	K	108/110 (98%)	92 (85%)	14 (13%)	2 (2%)	10	51
5	F	96/98 (98%)	84 (88%)	8 (8%)	4 (4%)	3	32
5	L	96/98 (98%)	84 (88%)	8 (8%)	4 (4%)	3	32
6	H	196/198 (99%)	159 (81%)	26 (13%)	11 (6%)	2	24
All	All	1842/1910 (96%)	1592 (86%)	192 (10%)	58 (3%)	5	41

5 of 58 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	5	ILE
2	B	94	THR
3	C	73	ASP
4	E	22	GLU
5	F	50	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	185/197 (94%)	166 (90%)	19 (10%)	9 40
1	G	185/197 (94%)	168 (91%)	17 (9%)	11 45
2	B	163/163 (100%)	136 (83%)	27 (17%)	3 16
3	C	155/155 (100%)	129 (83%)	26 (17%)	2 15
3	I	155/155 (100%)	129 (83%)	26 (17%)	2 15
4	D	94/94 (100%)	83 (88%)	11 (12%)	7 32
4	E	94/94 (100%)	81 (86%)	13 (14%)	4 25
4	J	94/94 (100%)	83 (88%)	11 (12%)	7 32
4	K	94/94 (100%)	81 (86%)	13 (14%)	4 25
5	F	83/83 (100%)	63 (76%)	20 (24%)	1 5
5	L	83/83 (100%)	62 (75%)	21 (25%)	1 4
6	H	165/165 (100%)	132 (80%)	33 (20%)	1 9
All	All	1550/1574 (98%)	1313 (85%)	237 (15%)	3 21

5 of 237 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	F	57	LEU
1	G	231	ILE
5	L	19	LEU
5	F	74	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	G	106	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	182	GLN
1	G	28	ASN
3	I	172	GLN
3	C	193	ASN
4	E	86	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
7	SIA	B	200	7	16,20,21	0.79	0	18,28,31	1.22	2 (11%)
7	GAL	B	201	7	12,12,12	0.68	0	17,17,17	0.54	0
7	SIA	C	200	7	16,20,21	0.62	0	18,28,31	1.08	2 (11%)
7	GAL	C	201	7	12,12,12	0.72	0	17,17,17	0.65	0
7	SIA	I	200	7	16,20,21	0.89	1 (6%)	18,28,31	1.24	1 (5%)
7	GAL	I	201	7	12,12,12	0.63	0	17,17,17	0.80	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SIA	B	200	7	-	0/14/34/38	0/1/1/1
7	GAL	B	201	7	-	0/2/22/22	0/1/1/1
7	SIA	C	200	7	-	0/14/34/38	0/1/1/1
7	GAL	C	201	7	-	0/2/22/22	0/1/1/1
7	SIA	I	200	7	-	0/14/34/38	0/1/1/1
7	GAL	I	201	7	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	I	200	SIA	C7-C6	-2.10	1.50	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	I	200	SIA	C8-C7-C6	-4.10	104.78	113.01
7	B	200	SIA	C8-C7-C6	-3.53	105.91	113.01
7	C	200	SIA	C8-C7-C6	-2.45	108.08	113.01
7	C	200	SIA	C3-C4-C5	2.21	113.94	111.47
7	B	200	SIA	C7-C6-C5	2.90	118.70	114.32

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	200	SIA	1	0
7	C	200	SIA	5	0
7	C	201	GAL	2	0
7	I	200	SIA	7	0

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers

EDS was not executed - this section will therefore be empty.