



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:46 PM GMT

PDB ID : 4PVD
Title : Crystal structure of yeast methylglyoxal/isovaleraldehyde reductase Gre2 complexed with NADPH
Authors : Guo, P.C.; Bao, Z.Z.; Li, W.F.; Zhou, C.Z.
Deposited on : 2014-03-17
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	<div> <div>2%</div> <div> <div></div> <div>94%</div> <div>5%</div> </div> </div>
1	B	342	<div> <div></div> <div> <div>88%</div> <div>12%</div> </div> </div>
1	C	342	<div> <div>8%</div> <div> <div></div> <div>80%</div> <div>9%</div> <div>10%</div> </div> </div>
1	D	342	<div> <div>10%</div> <div> <div></div> <div>79%</div> <div>9%</div> <div>11%</div> </div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 10416 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADPH-dependent methylglyoxal reductase GRE2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	0	0
			2690	1714	452	514	10			
1	B	342	Total	C	N	O	S	0	0	0
			2690	1714	452	514	10			
1	C	308	Total	C	N	O	S	0	0	0
			2422	1551	409	456	6			
1	D	305	Total	C	N	O	S	0	0	0
			2391	1531	405	449	6			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	40	Total 40	O 40	0	0
3	B	39	Total 39	O 39	0	0
3	C	28	Total 28	O 28	0	0
3	D	20	Total 20	O 20	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: NADPH-dependent methylglyoxal reductase GRE2





- Molecule 1: NADPH-dependent methylglyoxal reductase GRE2




- Molecule 1: NADPH-dependent methylglyoxal reductase GRE2



- Molecule 1: NADPH-dependent methylglyoxal reductase GRE2





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	90.31Å 92.89Å 201.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.40 44.21 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-2.40) 99.3 (44.21-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.73 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.242 , 0.288 0.237 , 0.277	Depositor DCC
R_{free} test set	3358 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.091	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.2	EDS
Estimated twinning fraction	0.000 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	1 of 66457 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10416	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.89 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 4.6663e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/2745	0.53	0/3705
1	B	0.37	0/2745	0.51	0/3705
1	C	0.38	0/2467	0.51	0/3322
1	D	0.38	0/2432	0.53	0/3271
All	All	0.38	0/10389	0.52	0/14003

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2690	0	2679	9	0
1	B	2690	0	2679	30	0
1	C	2422	0	2444	23	0
1	D	2391	0	2422	24	0
2	A	48	0	26	0	0
2	B	48	0	26	3	0
3	A	40	0	0	0	0
3	B	39	0	0	2	0
3	C	28	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	20	0	0	0	0
All	All	10416	0	10276	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:136:LYS:HB3	1:C:142:LEU:HD22	1.19	1.16
1:C:136:LYS:HB3	1:C:142:LEU:CD2	1.79	1.12
1:D:139:ASP:OD2	1:D:142:LEU:HD13	1.66	0.95
1:C:136:LYS:HD3	1:C:142:LEU:HD21	1.57	0.84
1:A:235:LEU:HD23	1:A:272:MET:HE3	1.64	0.78
1:B:256:ARG:O	1:B:259:ILE:HG12	1.85	0.76
1:B:289:ASN:HD22	1:B:289:ASN:N	1.84	0.76
1:B:270:PHE:HE1	1:B:272:MET:CE	1.97	0.76
1:B:270:PHE:HE1	1:B:272:MET:HE2	1.49	0.75
1:C:340:GLY:O	1:C:341:ARG:HB2	1.84	0.75
1:D:139:ASP:OD2	1:D:142:LEU:CD1	2.39	0.71
1:D:217:CYS:O	1:D:218:GLU:HG3	1.92	0.70
1:A:235:LEU:HD23	1:A:272:MET:CE	2.23	0.67
1:C:136:LYS:CB	1:C:142:LEU:CD2	2.67	0.67
1:C:218:GLU:OE1	1:C:218:GLU:HA	1.94	0.66
1:D:174:LYS:O	1:D:178:GLU:HG3	1.97	0.64
1:B:299:ALA:HA	1:B:304:LEU:HD13	1.79	0.64
1:A:154:TRP:O	1:A:155:GLU:HB2	1.98	0.63
1:B:79:LEU:CD2	1:B:123:VAL:HG13	2.29	0.62
1:B:87:PHE:HE1	1:B:165:TYR:HB2	1.63	0.62
1:D:127:SER:HB2	1:D:169:LYS:HD2	1.82	0.62
1:B:76:LYS:HG3	1:B:77:ILE:HG12	1.82	0.62
1:B:289:ASN:N	1:B:289:ASN:ND2	2.48	0.61
1:C:240:ILE:HG12	1:C:265:VAL:HG12	1.81	0.61
1:D:129:ALA:HA	1:D:132:PHE:CE1	2.37	0.60
1:D:235:LEU:HD23	1:D:272:MET:HE3	1.85	0.59
1:B:79:LEU:HD23	1:B:123:VAL:HG13	1.83	0.59
1:B:184:ARG:NH2	1:B:190:GLU:OE1	2.36	0.59
1:B:270:PHE:CE1	1:B:272:MET:HE2	2.35	0.59
1:D:53:GLU:HG3	1:D:67:VAL:HG23	1.85	0.58
1:B:337:LYS:HA	1:B:342:ILE:HD13	1.85	0.58
1:B:132:PHE:CE2	1:B:134:MET:HB3	2.39	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:136:LYS:HB3	1:C:142:LEU:HD21	1.78	0.56
1:C:136:LYS:CD	1:C:142:LEU:HD21	2.32	0.54
1:C:206:ASP:O	1:C:209:VAL:HG23	2.08	0.53
1:D:257:GLU:H	1:D:257:GLU:CD	2.12	0.52
1:B:290:ILE:HB	1:B:291:PRO:HD2	1.90	0.52
1:D:340:GLY:O	1:D:341:ARG:HB2	2.09	0.52
1:D:79:LEU:HD23	1:D:123:VAL:HB	1.92	0.52
1:C:340:GLY:O	1:C:341:ARG:CB	2.58	0.52
1:C:299:ALA:HA	1:C:304:LEU:HD13	1.92	0.52
1:B:274:ASP:OD1	1:B:300:THR:HG22	2.11	0.51
1:C:79:LEU:HD23	1:C:123:VAL:HG13	1.93	0.51
1:D:217:CYS:O	1:D:218:GLU:CG	2.58	0.50
1:B:126:SER:O	2:B:401:NDP:H6N	2.10	0.50
1:A:154:TRP:O	1:A:155:GLU:CB	2.59	0.50
1:D:340:GLY:O	1:D:341:ARG:CB	2.59	0.50
1:C:129:ALA:HA	1:C:132:PHE:CE1	2.47	0.49
1:D:270:PHE:HE1	1:D:272:MET:CE	2.26	0.49
1:D:300:THR:HG22	1:D:301:HIS:CD2	2.47	0.49
1:B:58:ILE:HG12	2:B:401:NDP:C6A	2.44	0.48
1:C:264:ILE:HG21	1:C:308:LEU:HD21	1.96	0.47
1:C:136:LYS:CB	1:C:142:LEU:HD22	2.14	0.47
1:D:217:CYS:C	1:D:218:GLU:HG3	2.35	0.47
1:D:217:CYS:O	1:D:218:GLU:OE2	2.33	0.46
1:B:270:PHE:CE1	1:B:272:MET:CE	2.88	0.46
1:B:257:GLU:H	1:B:257:GLU:CD	2.18	0.46
1:D:1:MET:HG3	1:D:1:MET:O	2.16	0.46
1:A:93:GLU:HA	1:A:97:LEU:HD12	1.99	0.45
1:C:290:ILE:HB	1:C:291:PRO:HD2	1.98	0.45
1:D:251:VAL:HG13	1:D:255:LYS:HE2	1.99	0.45
1:B:152:ALA:O	1:B:170:LYS:NZ	2.35	0.45
1:A:270:PHE:HE1	1:A:272:MET:HE1	1.82	0.44
1:C:218:GLU:OE1	1:C:218:GLU:CA	2.64	0.44
1:D:264:ILE:HG21	1:D:308:LEU:HD21	1.99	0.44
1:B:128:TYR:CE2	1:B:132:PHE:HB3	2.53	0.43
2:B:401:NDP:H4B	3:B:509:HOH:O	2.19	0.43
1:A:240:ILE:HG12	1:A:265:VAL:HG12	1.99	0.43
1:C:150:ASN:HA	1:C:151:PRO:HD3	1.93	0.43
1:D:128:TYR:CZ	1:D:132:PHE:HB3	2.54	0.43
1:B:190:GLU:HG3	1:B:259:ILE:HD12	2.01	0.42
1:B:146:GLU:OE2	1:B:310:ASN:HA	2.19	0.42
1:C:341:ARG:HD3	1:C:341:ARG:HA	1.76	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:18:ASP:O	1:B:22:LYS:HG3	2.19	0.42
1:A:342:ILE:HG23	1:A:342:ILE:OXT	2.20	0.42
1:C:5:VAL:O	1:C:29:GLY:HA2	2.20	0.42
1:B:302:ASN:HB2	1:B:304:LEU:CD1	2.50	0.42
1:D:270:PHE:HE1	1:D:272:MET:HE1	1.83	0.42
1:D:153:THR:HA	1:D:171:PHE:HZ	1.85	0.42
1:B:184:ARG:HD2	1:B:184:ARG:HA	1.83	0.41
1:C:310:ASN:HB2	1:C:313:SER:HB2	2.02	0.41
1:B:106:GLY:HA3	3:B:527:HOH:O	2.19	0.41
1:B:264:ILE:HG21	1:B:308:LEU:HD21	2.02	0.41
1:B:223:LEU:HD21	1:B:276:LEU:HG	2.02	0.41
1:C:256:ARG:O	1:C:259:ILE:HG13	2.20	0.41
1:D:198:TYR:OH	1:D:235:LEU:HD11	2.21	0.41
1:A:225:HIS:CE1	1:A:339:GLU:OE2	2.74	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	340/342 (99%)	336 (99%)	2 (1%)	2 (1%)	30	43
1	B	340/342 (99%)	335 (98%)	4 (1%)	1 (0%)	46	63
1	C	300/342 (88%)	292 (97%)	7 (2%)	1 (0%)	46	63
1	D	293/342 (86%)	288 (98%)	4 (1%)	1 (0%)	46	63
All	All	1273/1368 (93%)	1251 (98%)	17 (1%)	5 (0%)	39	56

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	155	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	197	VAL
1	B	197	VAL
1	D	197	VAL
1	A	197	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/297 (100%)	291 (98%)	6 (2%)	63	81
1	B	297/297 (100%)	293 (99%)	4 (1%)	76	89
1	C	265/297 (89%)	253 (96%)	12 (4%)	34	52
1	D	263/297 (89%)	257 (98%)	6 (2%)	58	78
All	All	1122/1188 (94%)	1094 (98%)	28 (2%)	55	76

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	76	LYS
1	A	83	SER
1	A	117	ASP
1	A	132	PHE
1	A	160	ASP
1	A	184	ARG
1	B	118	SER
1	B	132	PHE
1	B	163	ASN
1	B	289	ASN
1	C	2	SER
1	C	24	ASP
1	C	58	ILE
1	C	123	VAL
1	C	132	PHE
1	C	137	GLU
1	C	184	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	230	ASP
1	C	256	ARG
1	C	285	VAL
1	C	333	SER
1	C	341	ARG
1	D	81	THR
1	D	105	LYS
1	D	132	PHE
1	D	218	GLU
1	D	285	VAL
1	D	300	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	HIS
1	A	203	GLN
1	B	289	ASN
1	C	109	HIS
1	D	203	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	A	401	-	42,52,52	1.56	5 (11%)	55,80,80	1.93	5 (9%)
2	NDP	B	401	-	42,52,52	1.53	5 (11%)	55,80,80	1.86	5 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	401	-	-	0/30/77/77	0/5/5/5
2	NDP	B	401	-	-	0/30/77/77	0/5/5/5

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	NDP	C2N-C3N	2.26	1.40	1.34
2	B	401	NDP	C2N-C3N	2.45	1.40	1.34
2	A	401	NDP	C2A-N1A	2.49	1.38	1.33
2	B	401	NDP	C2A-N1A	2.52	1.38	1.33
2	A	401	NDP	C6N-C5N	2.95	1.39	1.33
2	B	401	NDP	C6N-C5N	3.09	1.39	1.33
2	A	401	NDP	C2A-N3A	3.60	1.38	1.32
2	B	401	NDP	C2A-N3A	3.66	1.38	1.32
2	B	401	NDP	O7N-C7N	6.65	1.41	1.24
2	A	401	NDP	O7N-C7N	7.01	1.42	1.24

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	NDP	N3A-C2A-N1A	-12.54	119.29	128.89
2	B	401	NDP	N3A-C2A-N1A	-11.65	119.97	128.89
2	B	401	NDP	PN-O3-PA	-2.76	124.98	132.73
2	B	401	NDP	C1D-N1N-C6N	-2.54	115.12	120.81
2	A	401	NDP	C1D-N1N-C6N	-2.44	115.35	120.81
2	A	401	NDP	PN-O3-PA	-2.19	126.59	132.73
2	B	401	NDP	C4A-C5A-N7A	-2.15	107.50	109.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	A	401	NDP	C1B-N9A-C4A	-2.01	123.90	126.94
2	A	401	NDP	O4D-C1D-N1N	2.55	113.46	108.07
2	B	401	NDP	O4D-C1D-N1N	2.78	113.93	108.07

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	NDP	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/342 (100%)	-0.01	6 (1%) 71 71	26, 44, 69, 90	0
1	B	342/342 (100%)	0.02	1 (0%) 94 94	26, 49, 77, 95	0
1	C	308/342 (90%)	0.18	29 (9%) 11 10	28, 47, 96, 109	0
1	D	305/342 (89%)	0.45	34 (11%) 7 7	28, 50, 86, 131	0
All	All	1297/1368 (94%)	0.15	70 (5%) 29 30	26, 48, 78, 131	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	131	VAL	9.3
1	D	97	LEU	6.9
1	D	138	ASN	6.6
1	D	307	THR	6.1
1	D	128	TYR	5.8
1	D	145	ASN	5.3
1	D	340	GLY	5.2
1	D	143	THR	5.1
1	D	171	PHE	5.1
1	C	97	LEU	5.0
1	D	84	PRO	4.8
1	D	142	LEU	4.7
1	D	168	SER	4.7
1	D	133	ASP	4.6
1	C	171	PHE	4.5
1	D	303	THR	4.4
1	D	98	ILE	4.3
1	D	305	GLY	4.3
1	C	139	ASP	4.0
1	C	1	MET	4.0
1	D	129	ALA	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	96	LEU	3.8
1	D	130	ALA	3.8
1	D	306	ALA	3.6
1	D	144	PHE	3.6
1	D	151	PRO	3.5
1	C	150	ASN	3.5
1	C	147	GLU	3.4
1	C	143	THR	3.4
1	C	148	SER	3.3
1	C	211	LYS	3.2
1	C	133	ASP	3.2
1	D	140	LYS	3.1
1	C	84	PRO	3.1
1	C	146	GLU	3.1
1	D	137	GLU	3.1
1	D	152	ALA	3.0
1	D	211	LYS	3.0
1	C	82	ALA	2.9
1	D	304	LEU	2.8
1	D	308	LEU	2.8
1	C	307	THR	2.8
1	C	142	LEU	2.8
1	C	138	ASN	2.8
1	C	140	LYS	2.7
1	C	141	SER	2.6
1	D	136	LYS	2.6
1	C	308	LEU	2.6
1	D	217	CYS	2.6
1	B	135	ALA	2.6
1	C	303	THR	2.6
1	C	135	ALA	2.6
1	C	144	PHE	2.6
1	C	304	LEU	2.5
1	A	338	PHE	2.5
1	A	88	ASP	2.4
1	D	132	PHE	2.4
1	A	342	ILE	2.4
1	D	139	ASP	2.3
1	C	136	LYS	2.2
1	C	145	ASN	2.2
1	A	182	GLU	2.2
1	C	117	ASP	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	134	MET	2.2
1	A	1	MET	2.1
1	C	129	ALA	2.1
1	C	169	LYS	2.1
1	D	169	LYS	2.1
1	A	207	LYS	2.0
1	D	170	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NDP	B	401	48/48	0.97	0.12	-0.76	34,36,39,41	0
2	NDP	A	401	48/48	0.96	0.13	-0.87	34,38,41,42	0

6.5 Other polymers [i](#)

There are no such residues in this entry.