



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 11:48 AM GMT

PDB ID : 3Q2S  
Title : Crystal Structure of CFIm68 RRM/CFIm25 complex  
Authors : Yang, Q.; Coseno, M.; Gilmartin, G.M.; Doublié, S.  
Deposited on : 2010-12-20  
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1668 (2.90-2.90)
Ramachandran outliers	100387	1630 (2.90-2.90)
Sidechain outliers	100360	1632 (2.90-2.90)
RSRZ outliers	91569	1456 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	207	<div> <div>2%</div> <div>57%</div> <div>40%</div> <div>.</div> </div>
1	B	207	<div> <div>%</div> <div>65%</div> <div>30%</div> <div>..</div> </div>
2	C	229	<div> <div>%</div> <div>18%</div> <div>17%</div> <div>5%</div> <div>59%</div> </div>
2	D	229	<div> <div>21%</div> <div>15%</div> <div>.</div> <div>60%</div> </div>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4863 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cleavage and polyadenylation specificity factor subunit 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	207	Total	C	N	O	S	4	2	0
			1709	1113	288	304	4			
1	B	203	Total	C	N	O	S	0	2	0
			1677	1097	282	294	4			

- Molecule 2 is a protein called Cleavage and polyadenylation specificity factor subunit 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	93	Total	C	N	O	S	0	0	0
			731	465	125	139	2			
2	D	92	Total	C	N	O	S	0	0	0
			726	462	124	138	2			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	159	VAL	CYS	ENGINEERED MUTATION	UNP Q16630
C	236	HIS	-	EXPRESSION TAG	UNP Q16630
C	237	HIS	-	EXPRESSION TAG	UNP Q16630
C	238	HIS	-	EXPRESSION TAG	UNP Q16630
C	239	HIS	-	EXPRESSION TAG	UNP Q16630
C	240	HIS	-	EXPRESSION TAG	UNP Q16630
C	241	HIS	-	EXPRESSION TAG	UNP Q16630
D	159	VAL	CYS	ENGINEERED MUTATION	UNP Q16630
D	236	HIS	-	EXPRESSION TAG	UNP Q16630
D	237	HIS	-	EXPRESSION TAG	UNP Q16630
D	238	HIS	-	EXPRESSION TAG	UNP Q16630
D	239	HIS	-	EXPRESSION TAG	UNP Q16630
D	240	HIS	-	EXPRESSION TAG	UNP Q16630
D	241	HIS	-	EXPRESSION TAG	UNP Q16630

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total 5	O 5	0	0
3	B	12	Total 12	O 12	0	0
3	C	2	Total 2	O 2	0	0
3	D	1	Total 1	O 1	0	0



PRO	GLY	GLY	ASP	ARG	PHE	PRO	GLY	PRO	GLY	ALA	GLY	GLY	PRO	GLY	GLY	PRO	PRO	PRO	PRO	PRO	PHE	PRO	ALA	GLY	GLN	THR	HIS	HIS	HIS	HIS	HIS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

● Molecule 2: Cleavage and polyadenylation specificity factor subunit 6



ASP	VAL	GLY	GLU	THR	PHE	ASN	GLN	GLU	ALA	ALA	THR	GLY	HIS	ASP	GLN	ILE	ASP	LEU	TYR	ASP	ASP	VAL	ILE	SER	PRO	SER	SER	ALA	ASN	ASN	GLY	ASP	ALA	PRO	GLU	ASP	ARG	ASP	TYR	MET	ASP	THR	LEU	PRO	PRO	THR	VAL	GLY	ASP	VAL	GLY	LYS	GLY	ALA	ALA	PRO	ASN	VAL
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

VAL	TYR	THR	THR	THR	GLY	LYS	ARG	I81	A82	L83	Y84	I85	G86	N87	L88	L97	T98	E99	A100	V101	H102	S103	L104	G105	V106	N107	D108	I109	I112	K113	R118	S123	A127	L128	V129	S136	S137	K138	K139	L140	M141	D142	L143	L144	P145	K146	R147	E148	L149	Q152	V156
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

T157	F158	V159	M160	F163	Q166	F167	E168	M169	R172	LYS	THR	THR	THR	GLN	SER	GLY	GLN	GLN	MET	SER	SER	GLY	GLY	GLY	LYS	ALA	GLY	PRO	PRO	GLY	GLY	GLY	ARG	GLY	ARG	GLY	ARG	PHE	PRO	GLN	GLY	GLY	ARG	GLY	VAL	PRO	GLY	GLY	ASP	ARG	PHE	PRO	GLY
------	------	------	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

PRO	ALA	GLY	PRO	GLY	GLY	PRO	PRO	PRO	PRO	PHE	PRO	ALA	ALA	GLY	GLN	THR	THR	HIS	HIS	HIS	HIS	HIS
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.32Å 139.32Å 139.32Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.90 – 2.90 19.90 – 2.90	Depositor EDS
% Data completeness (in resolution range)	82.4 (19.90-2.90) 82.6 (19.90-2.90)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.88Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6 _289)	Depositor
R, $R_{free}$	0.213 , 0.278 (Not available) , (Not available)	Depositor DCC
$R_{free}$ test set	NotAvailable	DCC
Wilson B-factor (Å <sup>2</sup> )	67.6	Xtriage
Anisotropy	0.000	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.26 , 47.8	EDS
Estimated twinning fraction	0.040 for l,-k,h 0.053 for l,-k,h	Xtriage
Reported twinning fraction	0.040 for l,-k,h	Depositor
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 20174 reflections (0.005%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4863	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.32	0/1763	0.50	0/2394
1	B	0.33	0/1730	0.52	0/2347
2	C	0.29	0/745	0.49	0/1007
2	D	0.29	0/740	0.45	0/1000
All	All	0.32	0/4978	0.50	0/6748

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1709	0	1719	79	0
1	B	1677	0	1698	72	0
2	C	731	0	723	50	0
2	D	726	0	721	45	0
3	A	5	0	0	0	0
3	B	12	0	0	1	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
All	All	4863	0	4861	241	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 25.



The worst 5 of 241 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:LEU:HG	1:A:191:TYR:CE2	1.65	1.31
1:B:99:LEU:HD12	1:B:99:LEU:N	1.51	1.18
1:B:76:MET:HE1	1:B:171:HIS:HB2	1.35	1.07
1:A:99:LEU:HG	1:A:191:TYR:CD2	1.92	1.05
1:A:99:LEU:CG	1:A:191:TYR:CE2	2.43	1.01

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	207/207 (100%)	194 (94%)	10 (5%)	3 (1%)	14	44
1	B	201/207 (97%)	185 (92%)	16 (8%)	0	100	100
2	C	91/229 (40%)	78 (86%)	12 (13%)	1 (1%)	17	51
2	D	90/229 (39%)	76 (84%)	12 (13%)	2 (2%)	8	31
All	All	589/872 (68%)	533 (90%)	50 (8%)	6 (1%)	19	54

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	59	SER
1	A	60	VAL
2	C	111	GLU
1	A	113	PRO
2	D	109	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/185 (101%)	168 (90%)	19 (10%)	9	27
1	B	183/185 (99%)	171 (93%)	12 (7%)	21	51
2	C	80/180 (44%)	65 (81%)	15 (19%)	2	6
2	D	80/180 (44%)	71 (89%)	9 (11%)	7	22
All	All	530/730 (73%)	475 (90%)	55 (10%)	10	26

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	128[A]	ILE
1	B	217	GLN
2	D	141	MET
1	B	128[B]	ILE
1	B	129[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	217	GLN
1	B	164	HIS
2	C	102	HIS
1	A	180	GLN
2	C	87	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	207/207 (100%)	-0.32	4 (1%) 70 66	36, 56, 100, 117	11 (5%)
1	B	203/207 (98%)	-0.29	3 (1%) 76 74	33, 53, 94, 118	9 (4%)
2	C	93/229 (40%)	-0.11	3 (3%) 51 43	46, 76, 103, 116	5 (5%)
2	D	92/229 (40%)	-0.11	1 (1%) 82 80	55, 79, 101, 105	7 (7%)
All	All	595/872 (68%)	-0.25	11 (1%) 71 68	33, 61, 101, 118	32 (5%)

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	C	134	GLU	3.8
1	B	133	ASP	3.0
2	C	132	GLY	2.9
1	A	132	GLN	2.9
1	A	101	THR	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.