



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:49 AM GMT

PDB ID : 3Q6B
Title : The high-resolution and new form crystal structure of BamA POTRA4-5 from E.coli
Authors : Gao, Z.Q.; Zhang, H.; Dong, Y.H.
Deposited on : 2010-12-31
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1422 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Outer membrane protein assembly complex, YaeT protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	155	1281	800	230	247	4	0	7	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	232	MET	-	EXPRESSION TAG	UNP C9QRL1
A	233	GLY	-	EXPRESSION TAG	UNP C9QRL1
A	234	SER	-	EXPRESSION TAG	UNP C9QRL1
A	235	SER	-	EXPRESSION TAG	UNP C9QRL1
A	236	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	237	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	238	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	239	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	240	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	241	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	242	SER	-	EXPRESSION TAG	UNP C9QRL1
A	243	SER	-	EXPRESSION TAG	UNP C9QRL1
A	244	GLY	-	EXPRESSION TAG	UNP C9QRL1
A	245	GLU	-	EXPRESSION TAG	UNP C9QRL1
A	246	ASN	-	EXPRESSION TAG	UNP C9QRL1
A	247	LEU	-	EXPRESSION TAG	UNP C9QRL1
A	248	TYR	-	EXPRESSION TAG	UNP C9QRL1
A	249	PHE	-	EXPRESSION TAG	UNP C9QRL1
A	250	GLU	-	EXPRESSION TAG	UNP C9QRL1
A	251	GLY	-	EXPRESSION TAG	UNP C9QRL1
A	252	SER	-	EXPRESSION TAG	UNP C9QRL1
A	253	HIS	-	EXPRESSION TAG	UNP C9QRL1
A	254	MET	-	EXPRESSION TAG	UNP C9QRL1
A	255	ALA	-	EXPRESSION TAG	UNP C9QRL1
A	256	SER	-	EXPRESSION TAG	UNP C9QRL1
A	257	MET	-	EXPRESSION TAG	UNP C9QRL1
A	258	THR	-	EXPRESSION TAG	UNP C9QRL1

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	259	GLY	-	EXPRESSION TAG	UNP C9QRL1
A	260	GLY	-	EXPRESSION TAG	UNP C9QRL1
A	261	GLN	-	EXPRESSION TAG	UNP C9QRL1
A	262	GLN	-	EXPRESSION TAG	UNP C9QRL1
A	263	MET	-	EXPRESSION TAG	UNP C9QRL1
A	264	GLY	-	EXPRESSION TAG	UNP C9QRL1
A	265	ARG	-	EXPRESSION TAG	UNP C9QRL1


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	141	Total O 141 141	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Outer membrane protein assembly complex, YaeT protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.53Å 39.01Å 32.56Å 90.00° 101.00° 90.00°	Depositor
Resolution (Å)	25.79 – 1.50 26.97 – 1.50	Depositor EDS
% Data completeness (in resolution range)	92.9 (25.79-1.50) 96.8 (26.97-1.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.50 (at 1.50Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.147 , 0.189 0.148 , 0.188	Depositor DCC
R_{free} test set	1344 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	15.4	Xtriage
Anisotropy	0.411	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 53.7	EDS
Estimated twinning fraction	0.035 for -h-2*k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 26677 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1422	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.32	0/1320	0.49	0/1773

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1281	0	1296	13	0
2	A	141	0	0	5	0
All	All	1422	0	1296	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:370:ARG:NH2	2:A:140:HOH:O	2.33	0.60
1:A:370:ARG:CZ	2:A:140:HOH:O	2.49	0.59
1:A:281:SER:O	1:A:285:GLU:HG2	2.03	0.59
1:A:391[B]:ARG:NH1	2:A:32:HOH:O	2.36	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:THR:HG22	1:A:304:MET:SD	2.49	0.52
1:A:361:LYS:O	1:A:365[B]:LEU:HD12	2.13	0.49
1:A:266:TYR:HE1	1:A:333:LYS:HG3	1.81	0.46
1:A:283[A]:GLU:HG3	2:A:67:HOH:O	2.15	0.45
1:A:365[A]:LEU:HD11	1:A:395:PHE:HZ	1.82	0.45
1:A:277:LEU:HD23	1:A:281:SER:HA	1.99	0.43
1:A:336:LYS:HE2	1:A:338:ARG:HE	1.83	0.43
1:A:365[A]:LEU:HD11	1:A:395:PHE:CZ	2.56	0.41
1:A:381:LEU:HD21	2:A:43:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/189 (85%)	158 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/161 (88%)	135 (96%)	6 (4%)	35	7

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	325	MET
1	A	330	ASP
1	A	333	LYS
1	A	345	ASN
1	A	365[A]	LEU
1	A	365[B]	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	345	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	155/189 (82%)	0.23	10 (6%)	22 23	10, 24, 49, 59	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	293	GLY	6.4
1	A	292	PRO	6.0
1	A	331	ALA	5.1
1	A	332	ASP	3.8
1	A	333	LYS	3.5
1	A	406	PRO	3.3
1	A	267	LYS	3.1
1	A	338	ARG	2.3
1	A	266	TYR	2.2
1	A	407	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.