



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 04:22 PM BST

PDB ID : 1QBH
Title : SOLUTION STRUCTURE OF A BACULOVIRAL INHIBITOR OF APOPTOSIS (IAP) REPEAT
Authors : Hinds, M.G.; Norton, R.S.; Vaux, D.L.; Day, C.L.
Deposited on : 1999-04-20

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	101	

2 Ensemble composition and analysis

This entry contains 20 models. Model 6 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:271-A:299, A:304-A:309, A:321-A:352 (67)	0.23	6

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	2, 3, 5, 6, 7, 9, 10, 11, 12, 13, 14, 16
2	1, 4, 18, 19
3	8, 15, 20
Single-model clusters	17

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1595 atoms, of which 768 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1).

Mol	Chain	Residues	Atoms						Trace
1	A	101	Total	C	H	N	O	S	0
			1594	524	768	144	149	9	

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	263	GLY	-	SEE REMARK 999	UNP Q13490
A	264	SER	-	SEE REMARK 999	UNP Q13490
A	265	HIS	-	SEE REMARK 999	UNP Q13490

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1


- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)



4.2.2 Score per residue for model 2


- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)






4.2.3 Score per residue for model 3

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.4 Score per residue for model 4

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.5 Score per residue for model 5

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.6 Score per residue for model 6 (medoid)

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.7 Score per residue for model 7

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.8 Score per residue for model 8

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.9 Score per residue for model 9

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.10 Score per residue for model 10

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.11 Score per residue for model 11

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.12 Score per residue for model 12

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)



4.2.13 Score per residue for model 13


- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)






4.2.14 Score per residue for model 14

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.15 Score per residue for model 15

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.16 Score per residue for model 16


- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)



4.2.17 Score per residue for model 17


- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)






4.2.18 Score per residue for model 18

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)




4.2.19 Score per residue for model 19

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)



4.2.20 Score per residue for model 20

- Molecule 1: INHIBITOR OF APOPTOSIS PROTEIN (2MIHB/C-IAP-1)



5 Refinement protocol and experimental data overview

The models were refined using the following method: *TORSION ANGLE DYNAMICS DISTANCE GEOMETRY SIMULATED ANNEALING MOLECULAR DYNAMICS*.

Of the 1000 calculated structures, 20 were deposited, based on the following criterion: *STRUCTURES WITH ACCEPTABLE COVALENT GEOMETRY, STRUCTURES WITH FAVORABLE NON- BOND ENERGY, STRUCTURES WITH THE LEAST RESTRAINT VIOLATIONS, STRUCTURES WITH THE LOWEST ENERGY, TARGET FUNCTION*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	1.5
X-PLOR	refinement	3.851

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.09±0.01	0±0/585 (0.0±0.0%)	1.98±0.04	20±2/794 (2.5±0.3%)
All	All	1.09	0/11700 (0.0%)	1.98	403/15880 (2.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.7±1.1
All	All	0	34

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	296	TYR	CB-CG-CD1	15.77	130.46	121.00	9	20
1	A	330	PHE	CB-CG-CD1	15.09	131.36	120.80	17	20
1	A	296	TYR	CB-CG-CD2	-13.45	112.93	121.00	14	20
1	A	297	TYR	CB-CG-CD1	12.93	128.76	121.00	13	20
1	A	335	PHE	CB-CG-CD1	12.12	129.28	120.80	12	20
1	A	297	TYR	CB-CG-CD2	-11.55	114.07	121.00	3	20
1	A	330	PHE	CB-CG-CD2	-11.25	112.92	120.80	5	20
1	A	276	PHE	CB-CG-CD1	10.69	128.28	120.80	10	20
1	A	335	PHE	CB-CG-CD2	-10.62	113.37	120.80	18	20
1	A	344	PHE	CB-CG-CD2	-9.49	114.16	120.80	19	20
1	A	344	PHE	CB-CG-CD1	9.21	127.25	120.80	19	20
1	A	352	TYR	CB-CG-CD2	-9.02	115.59	121.00	14	16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	352	TYR	CB-CG-CD1	8.46	126.08	121.00	14	16
1	A	286	GLN	N-CA-C	-8.44	88.22	111.00	20	17
1	A	276	PHE	CB-CG-CD2	-7.78	115.35	120.80	17	20
1	A	282	SER	N-CA-CB	7.15	121.23	110.50	4	1
1	A	283	VAL	CA-CB-CG1	6.56	120.74	110.90	15	12
1	A	327	ALA	CB-CA-C	6.46	119.79	110.10	2	2
1	A	333	CYS	N-CA-C	-6.43	93.63	111.00	4	8
1	A	297	TYR	CA-CB-CG	6.30	125.37	113.40	10	9
1	A	349	GLN	N-CA-CB	6.20	121.75	110.60	13	8
1	A	296	TYR	CB-CA-C	6.15	122.70	110.40	13	12
1	A	290	LEU	CB-CG-CD1	6.14	121.44	111.00	18	5
1	A	279	TRP	CA-CB-CG	6.10	125.29	113.70	20	8
1	A	274	ARG	NE-CZ-NH1	6.02	123.31	120.30	4	1
1	A	326	HIS	CA-CB-CG	5.96	123.74	113.60	19	9
1	A	277	MET	CA-CB-CG	5.88	123.30	113.30	15	3
1	A	309	CYS	CA-CB-SG	5.84	124.51	114.00	10	2
1	A	278	TYR	CB-CG-CD1	5.76	124.46	121.00	17	1
1	A	276	PHE	N-CA-CB	-5.65	100.43	110.60	16	8
1	A	332	ARG	N-CA-C	-5.64	95.77	111.00	16	4
1	A	295	PHE	CB-CA-C	-5.63	99.14	110.40	1	7
1	A	327	ALA	N-CA-CB	5.57	117.90	110.10	7	3
1	A	338	ARG	NE-CZ-NH2	-5.54	117.53	120.30	9	3
1	A	338	ARG	NE-CZ-NH1	5.33	122.96	120.30	1	1
1	A	272	ARG	NE-CZ-NH1	5.30	122.95	120.30	2	1
1	A	296	TYR	CA-CB-CG	5.17	123.22	113.40	9	1
1	A	274	ARG	NE-CZ-NH2	-5.13	117.74	120.30	4	1
1	A	306	CYS	CA-CB-SG	5.10	123.19	114.00	2	1
1	A	306	CYS	CB-CA-C	5.10	120.61	110.40	1	1
1	A	333	CYS	N-CA-CB	5.08	119.73	110.60	11	1
1	A	273	MET	CG-SD-CE	-5.05	92.11	100.20	3	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	279	TRP	Mainchain	12
1	A	272	ARG	Sidechain	8
1	A	274	ARG	Sidechain	5
1	A	332	ARG	Sidechain	5
1	A	351	ARG	Sidechain	3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	338	ARG	Sidechain	1

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	563	532	532	13±2
All	All	11280	10640	10640	264

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:299:GLY:HA3	1:A:304:VAL:HG13	0.79	1.53	9	8
1:A:298:VAL:HG13	1:A:307:PHE:CD2	0.73	2.18	7	1
1:A:323:TRP:CZ3	1:A:336:LEU:HD21	0.66	2.25	14	20
1:A:323:TRP:HZ3	1:A:336:LEU:HD21	0.65	1.52	19	20
1:A:326:HIS:CE1	1:A:336:LEU:HD23	0.63	2.29	20	19
1:A:279:TRP:HE1	1:A:283:VAL:HG23	0.60	1.56	13	13
1:A:344:PHE:CE1	1:A:348:ILE:HG13	0.57	2.34	9	16
1:A:348:ILE:HD12	1:A:352:TYR:CE2	0.56	2.36	13	15
1:A:336:LEU:HD11	1:A:340:LYS:HE3	0.56	1.77	2	2
1:A:299:GLY:HA3	1:A:304:VAL:HG12	0.55	1.78	14	3
1:A:293:ALA:O	1:A:326:HIS:CD2	0.54	2.61	18	18
1:A:279:TRP:HB2	1:A:280:PRO:HD3	0.51	1.83	6	10
1:A:323:TRP:HB3	1:A:344:PHE:CZ	0.50	2.42	9	20
1:A:323:TRP:HB3	1:A:344:PHE:CE2	0.49	2.42	10	20
1:A:287:PRO:HA	1:A:290:LEU:HD13	0.49	1.84	8	2
1:A:271:ALA:O	1:A:275:THR:HG22	0.49	2.07	12	18
1:A:297:TYR:CE1	1:A:298:VAL:HG22	0.48	2.43	5	4
1:A:299:GLY:CA	1:A:304:VAL:HG13	0.48	2.37	2	1
1:A:326:HIS:CE1	1:A:330:PHE:CD2	0.48	3.01	17	8
1:A:277:MET:HG2	1:A:278:TYR:CD1	0.47	2.44	2	2
1:A:340:LYS:HG3	1:A:344:PHE:CD2	0.45	2.46	5	12
1:A:275:THR:HG23	1:A:297:TYR:HA	0.45	1.88	17	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:298:VAL:HA	1:A:307:PHE:CZ	0.44	2.47	7	1
1:A:348:ILE:HD12	1:A:352:TYR:CZ	0.44	2.47	7	1
1:A:283:VAL:HG21	1:A:290:LEU:CD2	0.43	2.43	6	4
1:A:279:TRP:HZ2	1:A:290:LEU:HD23	0.43	1.74	7	1
1:A:306:CYS:SG	1:A:307:PHE:N	0.43	2.90	3	2
1:A:286:GLN:O	1:A:290:LEU:HD13	0.43	2.14	18	1
1:A:326:HIS:CE1	1:A:330:PHE:CE2	0.42	3.07	14	3
1:A:283:VAL:HG11	1:A:296:TYR:HD2	0.42	1.75	1	1
1:A:325:GLU:HG3	1:A:329:TRP:CH2	0.41	2.50	11	1
1:A:277:MET:HG3	1:A:278:TYR:CD1	0.41	2.50	15	1
1:A:297:TYR:CD1	1:A:298:VAL:HG22	0.41	2.50	9	1
1:A:326:HIS:ND1	1:A:326:HIS:O	0.41	2.48	4	2
1:A:298:VAL:HA	1:A:307:PHE:CE2	0.41	2.51	9	2
1:A:297:TYR:C	1:A:297:TYR:CD1	0.41	2.93	5	4
1:A:326:HIS:O	1:A:326:HIS:ND1	0.41	2.47	3	2
1:A:306:CYS:HB3	1:A:309:CYS:HB2	0.41	1.93	3	1
1:A:325:GLU:HB2	1:A:329:TRP:CH2	0.41	2.51	10	1
1:A:287:PRO:HB3	1:A:296:TYR:CD2	0.40	2.51	1	1
1:A:279:TRP:CB	1:A:280:PRO:HD3	0.40	2.47	3	1

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	67/101 (66%)	45±2 (67±3%)	14±2 (21±3%)	8±2 (12±2%)	1	8
All	All	1340/2020 (66%)	904 (67%)	279 (21%)	157 (12%)	1	8

All 15 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	277	MET	20
1	A	280	PRO	20
1	A	296	TYR	20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	285	VAL	20
1	A	304	VAL	15
1	A	331	PRO	12
1	A	306	CYS	12
1	A	327	ALA	11
1	A	279	TRP	10
1	A	344	PHE	7
1	A	282	SER	5
1	A	299	GLY	2
1	A	309	CYS	1
1	A	278	TYR	1
1	A	337	ILE	1

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	59/88 (67%)	40±2 (68±3%)	19±2 (32±3%)	1	14
All	All	1180/1760 (67%)	806 (68%)	374 (32%)	1	14

All 36 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	330	PHE	20
1	A	296	TYR	20
1	A	278	TYR	20
1	A	275	THR	20
1	A	297	TYR	20
1	A	335	PHE	20
1	A	338	ARG	20
1	A	344	PHE	20
1	A	348	ILE	20
1	A	276	PHE	20
1	A	340	LYS	20
1	A	279	TRP	18
1	A	332	ARG	14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	286	GLN	13
1	A	329	TRP	13
1	A	337	ILE	12
1	A	304	VAL	12
1	A	321	ASP	11
1	A	309	CYS	8
1	A	283	VAL	8
1	A	272	ARG	5
1	A	336	LEU	5
1	A	328	LYS	5
1	A	308	CYS	4
1	A	349	GLN	4
1	A	305	LYS	4
1	A	282	SER	3
1	A	346	ASP	3
1	A	351	ARG	3
1	A	274	ARG	3
1	A	347	GLU	1
1	A	325	GLU	1
1	A	298	VAL	1
1	A	277	MET	1
1	A	289	GLN	1
1	A	339	MET	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided