



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 11:55 AM GMT

PDB ID : 3QFS
Title : Crystal Structure of NADPH-Cytochrome P450 Reductase (FAD/NADPH domain)
Authors : Xia, C.; Marohnic, C.; Panda, S.P.; Masters, B.S.; Kim, J.-J.P.
Deposited on : 2011-01-22
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1199 (1.40-1.40)
Clashscore	102246	1295 (1.40-1.40)
Ramachandran outliers	100387	1259 (1.40-1.40)
Sidechain outliers	100360	1258 (1.40-1.40)
RSRZ outliers	91569	1198 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	458	 6% 79% 15% • 5%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3984 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NADPH–cytochrome P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	435	3476	2199	615	646	16	0	0	0

There are 22 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	223	MET	-	EXPRESSION TAG	UNP P16435
A	224	GLY	-	EXPRESSION TAG	UNP P16435
A	225	SER	-	EXPRESSION TAG	UNP P16435
A	226	SER	-	EXPRESSION TAG	UNP P16435
A	227	HIS	-	EXPRESSION TAG	UNP P16435
A	228	HIS	-	EXPRESSION TAG	UNP P16435
A	229	HIS	-	EXPRESSION TAG	UNP P16435
A	230	HIS	-	EXPRESSION TAG	UNP P16435
A	231	HIS	-	EXPRESSION TAG	UNP P16435
A	232	HIS	-	EXPRESSION TAG	UNP P16435
A	233	SER	-	EXPRESSION TAG	UNP P16435
A	234	SER	-	EXPRESSION TAG	UNP P16435
A	235	GLY	-	EXPRESSION TAG	UNP P16435
A	236	LEU	-	EXPRESSION TAG	UNP P16435
A	237	VAL	-	EXPRESSION TAG	UNP P16435
A	238	PRO	-	EXPRESSION TAG	UNP P16435
A	239	ARG	-	EXPRESSION TAG	UNP P16435
A	240	GLY	-	EXPRESSION TAG	UNP P16435
A	241	SER	-	EXPRESSION TAG	UNP P16435
A	242	HIS	-	EXPRESSION TAG	UNP P16435
A	243	MET	-	EXPRESSION TAG	UNP P16435
A	503	VAL	ALA	SEE REMARK 999	UNP P16435

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	31	10	5	13	3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	424	Total 424	O 424	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NADPH-cytochrome P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.64Å 58.54Å 130.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.71 – 1.40 19.71 – 1.40	Depositor EDS
% Data completeness (in resolution range)	84.4 (19.71-1.40) 84.6 (19.71-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.24 (at 1.40Å)	Xtrriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.211 , 0.231 0.210 , 0.231	Depositor DCC
R_{free} test set	3544 reflections (5.08%)	DCC
Wilson B-factor (Å ²)	14.7	Xtrriage
Anisotropy	0.362	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 48.6	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	1 of 69748 reflections (0.001%)	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3984	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.40% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, FAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.34	1/3555 (0.0%)	0.58	1/4816 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	475	CYS	C-O	-5.67	1.12	1.23

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	505	GLU	OE1-CD-OE2	-5.34	116.89	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3476	0	3438	70	0
2	A	53	0	31	0	0
3	A	31	0	11	1	0
4	A	424	0	0	8	0
All	All	3984	0	3480	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:503:VAL:HG12	1:A:504:GLY:N	1.57	1.15
1:A:503:VAL:HG12	1:A:504:GLY:H	1.11	1.08
1:A:609:GLN:H	1:A:609:GLN:HE21	1.19	0.91
1:A:503:VAL:CG1	1:A:504:GLY:N	2.31	0.88
1:A:247:GLN:HE21	1:A:353:ASN:HD21	1.27	0.83
1:A:609:GLN:HE22	3:A:753:NAP:H2A	1.44	0.80
1:A:297:GLN:H	1:A:297:GLN:NE2	1.81	0.79
1:A:504:GLY:O	1:A:505:GLU:O	2.03	0.76
1:A:656:MET:HB2	1:A:660:GLN:HE21	1.49	0.75
1:A:609:GLN:NE2	1:A:609:GLN:H	1.87	0.73
1:A:504:GLY:O	1:A:507:GLY:O	2.07	0.72
1:A:609:GLN:HE21	1:A:609:GLN:N	1.91	0.68
1:A:505:GLU:O	1:A:507:GLY:N	2.27	0.68
1:A:247:GLN:HE21	1:A:353:ASN:ND2	1.93	0.66
1:A:635:ALA:HB2	1:A:678:VAL:HB	1.77	0.66
1:A:249:GLU:HB3	1:A:354:LEU:HD21	1.79	0.64
1:A:353:ASN:HD22	1:A:355:ASP:H	1.46	0.63
1:A:353:ASN:ND2	1:A:355:ASP:H	1.97	0.63
1:A:297:GLN:H	1:A:297:GLN:HE21	1.47	0.61
1:A:631:VAL:HB	1:A:676:LEU:CD2	2.31	0.61
1:A:303:LEU:HD22	1:A:577:LEU:HD21	1.81	0.61
1:A:503:VAL:O	1:A:508:GLY:O	2.19	0.60
1:A:253:HIS:HE1	1:A:350:SER:OG	1.84	0.59
1:A:656:MET:HB2	1:A:660:GLN:NE2	2.18	0.59
1:A:305:HIS:HD2	1:A:578:TYR:OH	1.86	0.59
1:A:674:TYR:HE2	1:A:676:LEU:HD21	1.69	0.58
1:A:322:HIS:HD2	4:A:25:HOH:O	1.86	0.57
1:A:353:ASN:HD22	1:A:354:LEU:N	2.00	0.57
1:A:613:LYS:HG3	1:A:616:ARG:NH2	2.19	0.57
1:A:504:GLY:O	1:A:505:GLU:C	2.43	0.57
1:A:593:GLN:HE21	1:A:595:ASN:HD21	1.53	0.57
1:A:505:GLU:O	1:A:506:ASN:C	2.38	0.56
1:A:505:GLU:C	1:A:507:GLY:N	2.59	0.56
1:A:503:VAL:CG1	1:A:504:GLY:H	1.93	0.55
1:A:631:VAL:HB	1:A:676:LEU:HD23	1.88	0.54
1:A:247:GLN:NE2	1:A:353:ASN:HD21	2.00	0.54
1:A:276:LYS:HB3	1:A:276:LYS:NZ	2.22	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:399:PRO:O	1:A:403:GLU:HG3	2.08	0.54
1:A:667:LYS:O	1:A:671:LYS:HG3	2.08	0.52
1:A:429:HIS:CD2	1:A:432:ALA:H	2.28	0.52
1:A:613:LYS:HE2	1:A:616:ARG:CZ	2.40	0.52
1:A:305:HIS:HE1	1:A:307:GLU:OE1	1.95	0.50
1:A:335:ASN:ND2	1:A:371:ARG:HH12	2.10	0.49
1:A:429:HIS:HD2	1:A:432:ALA:H	1.59	0.48
1:A:505:GLU:C	1:A:507:GLY:H	2.17	0.48
1:A:362:HIS:HE1	4:A:704:HOH:O	1.96	0.48
1:A:587:ARG:HG3	4:A:690:HOH:O	2.13	0.47
1:A:666:LYS:O	1:A:670:THR:HG23	2.13	0.47
1:A:393:ALA:O	1:A:402:GLN:HG3	2.17	0.45
1:A:297:GLN:N	1:A:297:GLN:NE2	2.57	0.45
1:A:637:ASN:HB3	4:A:839:HOH:O	2.17	0.45
1:A:324:ALA:HB2	1:A:458:TYR:CE1	2.52	0.45
1:A:485:ALA:HB3	4:A:142:HOH:O	2.17	0.45
1:A:636:ARG:HG2	1:A:637:ASN:ND2	2.32	0.44
1:A:564:LEU:HD12	1:A:564:LEU:N	2.33	0.44
1:A:628:HIS:HE1	4:A:200:HOH:O	2.01	0.44
1:A:330:ASP:OD2	1:A:332:ALA:HB3	2.18	0.43
1:A:335:ASN:HD21	1:A:371:ARG:NH1	2.17	0.43
1:A:620:TRP:CZ3	1:A:650:VAL:HG13	2.53	0.43
1:A:313:SER:O	1:A:314:LYS:HB2	2.19	0.43
1:A:353:ASN:HD22	1:A:353:ASN:C	2.22	0.43
1:A:571:ARG:HB2	1:A:574:GLU:HB3	2.01	0.42
1:A:541:ALA:HB3	1:A:542:PRO:CD	2.50	0.42
1:A:626:GLY:HA2	1:A:673:ARG:CZ	2.49	0.42
1:A:351:LEU:O	1:A:362:HIS:HD2	2.02	0.41
1:A:501:GLU:HG3	1:A:503:VAL:H	1.86	0.41
1:A:635:ALA:HB2	1:A:678:VAL:CB	2.48	0.41
1:A:468:HIS:HE1	4:A:136:HOH:O	2.03	0.41
1:A:639:ALA:HB2	1:A:678:VAL:HG11	2.03	0.40
1:A:628:HIS:HD2	4:A:152:HOH:O	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	433/458 (94%)	418 (96%)	12 (3%)	3 (1%)	26 6

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	505	GLU
1	A	508	GLY
1	A	503	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	373/394 (95%)	367 (98%)	6 (2%)	70 38

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	297	GLN
1	A	349	MET
1	A	353	ASN
1	A	559	GLU
1	A	576	TYR
1	A	609	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	253	HIS
1	A	274	ASN
1	A	283	ASN
1	A	297	GLN
1	A	305	HIS
1	A	322	HIS
1	A	335	ASN
1	A	353	ASN
1	A	362	HIS
1	A	394	GLN
1	A	429	HIS
1	A	468	HIS
1	A	470	ASN
1	A	489	ASN
1	A	506	ASN
1	A	584	GLN
1	A	593	GLN
1	A	609	GLN
1	A	628	HIS
1	A	637	ASN
1	A	643	GLN
1	A	660	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FAD	A	752	-	48,58,58	2.69	21 (43%)	54,89,89	3.12	12 (22%)
3	NAP	A	753	-	27,33,52	2.31	5 (18%)	34,52,80	1.75	4 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	752	-	-	0/30/50/50	0/6/6/6
3	NAP	A	753	-	-	0/17/37/67	0/3/3/5

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	752	FAD	C5'-C4'	-5.03	1.43	1.51
3	A	753	NAP	PA-O5B	-2.36	1.48	1.59
2	A	752	FAD	P-O5'	-2.01	1.49	1.59
2	A	752	FAD	C5A-C4A	2.17	1.45	1.40
2	A	752	FAD	C6-C7	2.20	1.43	1.37
3	A	753	NAP	C2A-N1A	2.21	1.38	1.33
2	A	752	FAD	C1'-N10	2.26	1.50	1.48
2	A	752	FAD	C4-C4X	2.42	1.46	1.41
2	A	752	FAD	C8A-N7A	2.50	1.39	1.34
2	A	752	FAD	C5X-N5	2.78	1.39	1.35
2	A	752	FAD	O2'-C2'	2.87	1.49	1.43
3	A	753	NAP	C5A-C4A	2.93	1.47	1.40
2	A	752	FAD	C10-N1	3.03	1.40	1.35
2	A	752	FAD	C6-C5X	3.05	1.46	1.41
2	A	752	FAD	C9A-C5X	3.09	1.48	1.42
3	A	753	NAP	P2B-O1X	3.13	1.61	1.51
2	A	752	FAD	O4B-C1B	3.15	1.45	1.41
2	A	752	FAD	C2A-N3A	3.21	1.37	1.32
2	A	752	FAD	C8-C7	3.22	1.49	1.41
2	A	752	FAD	C4X-C10	3.27	1.47	1.41
2	A	752	FAD	C2'-C3'	3.42	1.60	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	752	FAD	C4X-N5	4.24	1.40	1.33
2	A	752	FAD	C4-N3	5.65	1.43	1.33
2	A	752	FAD	C4A-N3A	6.97	1.46	1.35
2	A	752	FAD	C9A-N10	8.19	1.50	1.38
3	A	753	NAP	C4A-N3A	9.74	1.50	1.35

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	752	FAD	N3A-C2A-N1A	-9.09	121.93	128.89
2	A	752	FAD	C4X-C4-N3	-6.96	114.08	123.59
3	A	753	NAP	N3A-C2A-N1A	-6.67	123.78	128.89
2	A	752	FAD	C4X-C10-N10	-4.83	117.67	120.52
2	A	752	FAD	C1B-N9A-C4A	-4.08	120.79	126.94
2	A	752	FAD	C4-C4X-C10	-3.17	117.91	119.94
2	A	752	FAD	C5X-C9A-N10	-2.78	115.51	117.62
2	A	752	FAD	P-O3P-PA	-2.74	125.04	132.73
3	A	753	NAP	C4B-O4B-C1B	-2.46	107.02	109.72
2	A	752	FAD	O2P-P-O3P	2.13	114.76	105.09
3	A	753	NAP	C2A-N1A-C6A	2.38	123.03	118.77
2	A	752	FAD	O2'-C2'-C1'	2.59	116.31	109.94
3	A	753	NAP	C4A-C5A-N7A	3.60	112.79	109.48
2	A	752	FAD	C1'-N10-C9A	4.55	123.97	118.86
2	A	752	FAD	C2A-N1A-C6A	5.13	127.94	118.77
2	A	752	FAD	C4-N3-C2	14.96	128.18	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	753	NAP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	435/458 (94%)	0.34	29 (6%) 21 18	11, 18, 28, 35	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	503	VAL	8.2
1	A	636	ARG	8.0
1	A	507	GLY	8.0
1	A	504	GLY	7.4
1	A	680	SER	5.5
1	A	390	TYR	4.2
1	A	505	GLU	4.2
1	A	506	ASN	4.1
1	A	635	ALA	3.7
1	A	679	TRP	3.7
1	A	658	HIS	3.5
1	A	300	GLU	3.4
1	A	255	ASP	3.3
1	A	357	GLU	2.8
1	A	587	ARG	2.8
1	A	484	LYS	2.8
1	A	299	THR	2.7
1	A	670	THR	2.7
1	A	558	LYS	2.5
1	A	356	GLU	2.5
1	A	387	ASN	2.3
1	A	637	ASN	2.2
1	A	502	PRO	2.2
1	A	634	ASP	2.1
1	A	655	ALA	2.1
1	A	525	PHE	2.1
1	A	380	ILE	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	640	ARG	2.0
1	A	659	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	NAP	A	753	31/48	0.94	0.08	-0.26	14,16,31,34	0
2	FAD	A	752	53/53	0.96	0.08	-0.36	10,13,23,27	0

6.5 Other polymers [i](#)

There are no such residues in this entry.