



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 09:46 PM GMT

PDB ID : 1QGH
Title : THE X-RAY STRUCTURE OF THE UNUSUAL DODECAMERIC FER-
RITIN FROM LISTERIA INNOCUA, REVEALS A NOVEL INTERSUB-
UNIT IRON BINDING SITE.
Authors : Ilari, A.; Stefanini, S.; Chiancone, E.; Tsernoglou, D.
Deposited on : 1999-04-27
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1456 (2.38-2.34)
Ramachandran outliers	100387	1435 (2.38-2.34)
Sidechain outliers	100360	1436 (2.38-2.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	156	69% 24% . .
1	B	156	69% 24% . .
1	C	156	70% 22% . .
1	D	156	68% 24% . .
1	E	156	67% 26% . .
1	F	156	65% 25% 6% .
1	G	156	69% 19% 8% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	H	156	 67% 25% . .
1	I	156	 63% 27% 5% . . .
1	J	156	 68% 25% . . .
1	K	156	 65% 24% 8% .
1	L	156	 67% 25% . . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 15155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NON-HEME IRON-CONTAINING FERRITIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	150	1222	784	196	235	7	32	0	0
1	B	150	1222	784	196	235	7	57	0	0
1	C	150	1222	784	196	235	7	44	0	0
1	D	150	1222	784	196	235	7	47	0	0
1	E	150	1222	784	196	235	7	46	0	0
1	F	150	1222	784	196	235	7	42	0	0
1	G	150	1222	784	196	235	7	41	0	0
1	H	150	1222	784	196	235	7	34	0	0
1	I	150	1222	784	196	235	7	45	0	0
1	J	150	1222	784	196	235	7	44	0	0
1	K	150	1222	784	196	235	7	43	0	0
1	L	150	1222	784	196	235	7	50	0	0

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total	Fe	0	0
			1	1		
2	J	1	Total	Fe	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Fe 1 1	0	0
2	K	1	Total Fe 1 1	0	0
2	E	1	Total Fe 1 1	0	0
2	H	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0
2	I	1	Total Fe 1 1	0	0
2	C	1	Total Fe 1 1	0	0
2	A	1	Total Fe 1 1	0	0
2	L	1	Total Fe 1 1	0	0
2	F	1	Total Fe 1 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	38	Total O 38 38	0	0
3	B	22	Total O 22 22	0	0
3	C	36	Total O 36 36	0	0
3	D	28	Total O 28 28	0	0
3	E	50	Total O 50 50	0	0
3	F	42	Total O 42 42	0	0
3	G	54	Total O 54 54	0	0
3	H	44	Total O 44 44	0	0
3	I	33	Total O 33 33	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	J	46	Total 46	O 46	0	0
3	K	41	Total 41	O 41	0	0
3	L	45	Total 45	O 45	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN




- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN



- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN




- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN






- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain E: 67% 26%




- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain F: 65% 25% 6%



- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain G: 69% 19% 8%




- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN

Chain H: 67% 25%




- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN


Chain I: 63% 27% 5%




- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN



- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN



- Molecule 1: NON-HEME IRON-CONTAINING FERRITIN



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.54Å 137.50Å 173.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.35	Depositor
% Data completeness (in resolution range)	91.0 (10.00-2.35)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
Refinement program	TNT V. 5-E	Depositor
R, R_{free}	0.219 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	15155	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: FE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/1249	1.15	5/1684 (0.3%)
1	B	0.72	0/1249	1.19	1/1684 (0.1%)
1	C	0.70	0/1249	1.14	6/1684 (0.4%)
1	D	0.72	0/1249	1.24	9/1684 (0.5%)
1	E	0.70	0/1249	1.17	7/1684 (0.4%)
1	F	0.68	0/1249	1.20	7/1684 (0.4%)
1	G	0.70	0/1249	1.13	5/1684 (0.3%)
1	H	0.71	0/1249	1.17	6/1684 (0.4%)
1	I	0.71	0/1249	1.21	8/1684 (0.5%)
1	J	0.72	1/1249 (0.1%)	1.20	5/1684 (0.3%)
1	K	0.72	0/1249	1.19	9/1684 (0.5%)
1	L	0.73	1/1249 (0.1%)	1.24	8/1684 (0.5%)
All	All	0.71	2/14988 (0.0%)	1.19	76/20208 (0.4%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	L	156	GLU	CD-OE1	6.00	1.32	1.25
1	J	156	GLU	C-O	5.67	1.34	1.23

The worst 5 of 76 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	156	GLU	CA-C-O	9.79	140.65	120.10
1	L	156	GLU	OE1-CD-OE2	-9.23	112.23	123.30
1	L	7	VAL	O-C-N	-8.84	108.56	122.70
1	D	110	ARG	NE-CZ-NH1	7.78	124.19	120.30
1	I	110	ARG	NE-CZ-NH2	-7.40	116.60	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1222	0	1186	32	0
1	B	1222	0	1186	34	0
1	C	1222	0	1186	33	0
1	D	1222	0	1186	42	0
1	E	1222	0	1186	38	0
1	F	1222	0	1186	53	0
1	G	1222	0	1186	46	0
1	H	1222	0	1186	33	0
1	I	1222	0	1186	46	0
1	J	1222	0	1186	33	0
1	K	1222	0	1186	63	0
1	L	1222	0	1186	38	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
2	I	1	0	0	0	0
2	J	1	0	0	0	0
2	K	1	0	0	0	0
2	L	1	0	0	0	0
3	A	38	0	0	3	0
3	B	22	0	0	0	0
3	C	36	0	0	3	0
3	D	28	0	0	6	0
3	E	50	0	0	7	0
3	F	42	0	0	6	0
3	G	54	0	0	8	0
3	H	44	0	0	3	0
3	I	33	0	0	3	0
3	J	46	0	0	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	K	41	0	0	10	0
3	L	45	0	0	4	0
All	All	15155	0	14232	477	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

The worst 5 of 477 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:94:THR:HG22	1:G:97:GLN:H	1.11	1.12
1:F:94:THR:HB	1:F:97:GLN:HG3	1.20	1.12
1:F:94:THR:HG22	1:F:96:ASP:N	1.69	1.08
1:F:7:VAL:HG12	1:F:8:ASP:H	1.22	1.04
1:I:94:THR:HB	1:I:97:GLN:HG3	1.40	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	B	148/156 (95%)	146 (99%)	2 (1%)	0	100	100
1	C	148/156 (95%)	146 (99%)	2 (1%)	0	100	100
1	D	148/156 (95%)	142 (96%)	6 (4%)	0	100	100
1	E	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	F	148/156 (95%)	143 (97%)	5 (3%)	0	100	100
1	G	148/156 (95%)	146 (99%)	2 (1%)	0	100	100
1	H	148/156 (95%)	146 (99%)	2 (1%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	I	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	J	148/156 (95%)	145 (98%)	3 (2%)	0	100	100
1	K	148/156 (95%)	144 (97%)	4 (3%)	0	100	100
1	L	148/156 (95%)	143 (97%)	5 (3%)	0	100	100
All	All	1776/1872 (95%)	1736 (98%)	40 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	133/139 (96%)	121 (91%)	12 (9%)	12	12
1	B	133/139 (96%)	123 (92%)	10 (8%)	17	18
1	C	133/139 (96%)	123 (92%)	10 (8%)	17	18
1	D	133/139 (96%)	121 (91%)	12 (9%)	12	12
1	E	133/139 (96%)	122 (92%)	11 (8%)	14	15
1	F	133/139 (96%)	121 (91%)	12 (9%)	12	12
1	G	133/139 (96%)	120 (90%)	13 (10%)	10	9
1	H	133/139 (96%)	119 (90%)	14 (10%)	8	8
1	I	133/139 (96%)	121 (91%)	12 (9%)	12	12
1	J	133/139 (96%)	124 (93%)	9 (7%)	20	22
1	K	133/139 (96%)	120 (90%)	13 (10%)	10	9
1	L	133/139 (96%)	123 (92%)	10 (8%)	17	18
All	All	1596/1668 (96%)	1458 (91%)	138 (9%)	13	13

5 of 138 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	49	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	119	LEU
1	K	156	GLU
1	F	119	LEU
1	G	20	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 55 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	15	HIS
1	H	129	ASN
1	L	15	HIS
1	G	28	HIS
1	H	15	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 12 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

6.4 Ligands

EDS was not executed - this section will therefore be empty.

6.5 Other polymers

EDS was not executed - this section will therefore be empty.