



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 11:55 AM GMT

PDB ID : 3QGF

Title : Crystal structure of the hepatitis C virus NS5B RNA-dependent RNA polymerase complex with (2E)-3-(4-{|[(1-{|[(13-cyclohexyl-6-oxo-6,7-dihydro-5H-indolo[1,2-d][1,4]benzodiazepin-10-yl)carbonyl]amino}cyclopentyl)carbonyl]amino}phenyl)prop-2-enoic acid and (2R)-4-(6-chloropyridazin-3-yl)-N-(4-methoxybenzyl)-1-{|[4-(trifluoromethoxy)phenyl}sulfonyl)piperazine-2-carboxamide

Authors : Sheriff, S.

Deposited on : 2011-01-24

Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.7 (RC4), CSD as536be (2015)

Xtriage (Phenix) : 1.9-1692

EDS : rb-20026688

Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)

Refmac : 5.8.0135

CCP4 : 6.5.0

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	574	5%	83%	9%	• 7%
1	B	574	6%	83%	9%	• 7%

2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 8753 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RNA-directed RNA polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	534	4145	2613	733	767	32	0	0	0
1	B	533	4134	2605	731	766	32	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	INITIATING METHIONINE	UNP Q9WMX2
B	0	MET	-	INITIATING METHIONINE	UNP Q9WMX2


- Molecule 2 is (2E)-3-(4-{{[(1-{{{[(13-CYCLOHEXYL-6-OXO-6,7-DIHYDRO-5H-INDOLO[1,2-D][1,4]BENZODIAZEPIN-10-YL)CARBONYL]AMINO}CYCLOPENTYL)CARBONYL]AMINO}PHENYL)PROP-2-ENOIC ACID (three-letter code: 23E) (formula: C₃₈H₃₈N₄O₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			47	38	4	5		


Mol	Chain	Residues	Total	C	N	O	ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			47	38	4	5		

- Molecule 3 is (2R)-4-(6-CHLOROPYRIDAZIN-3-YL)-N-(4-METHOXYBENZYL)-1-{|4-(TRIFLUOROMETHOXY)PHENYL|SULFONYL}PIPERAZINE-2-CARBOXAMIDE (three-letter code: 46F) (formula: C₂₄H₂₃ClF₃N₅O₅S).



Mol	Chain	Residues	Atoms							ZeroOcc	AltConf
			Total	C	Cl	F	N	O	S		
3	A	1	Total	39	24	1	3	5	5	1	0
3	B	1	Total	39	24	1	3	5	5	1	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	A	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0
4	B	1	Total O S 5 4 1	0	0



- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	129	Total O 129 129	0	0
5	B	123	Total O 123 123	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA-directed RNA polymerase



- Molecule 1: RNA-directed RNA polymerase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.90 Å 90.10 Å 231.30 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	35.44 – 2.45 35.44 – 2.44	Depositor EDS
% Data completeness (in resolution range)	85.0 (35.44-2.45) 84.7 (35.44-2.44)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.78 (at 2.45 Å)	Xtriage
Refinement program	BUSTER 2.9.7	Depositor
R , R_{free}	0.225 , 0.269 0.218 , 0.269	Depositor DCC
R_{free} test set	1065 reflections (2.19%)	DCC
Wilson B-factor (Å ²)	26.3	Xtriage
Anisotropy	0.456	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.34$	Xtriage
Outliers	2 of 50113 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	8753	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 23E, 46F, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/4233	0.70	1/5740 (0.0%)
1	B	0.49	0/4222	0.69	1/5726 (0.0%)
All	All	0.50	0/8455	0.69	2/11466 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	554	GLY	C-N-CA	5.35	135.07	121.70
1	B	535	LYS	C-N-CA	5.26	134.85	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4145	0	4162	18	0
1	B	4134	0	4142	21	0
2	A	47	0	37	1	0
2	B	47	0	37	0	0
3	A	39	0	23	1	0
3	B	39	0	23	1	0
4	A	30	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	20	0	0	0	0
5	A	129	0	0	0	0
5	B	123	0	0	0	0
All	All	8753	0	8424	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:465:ARG:HG3	1:A:545:LEU:HB3	1.69	0.74
1:B:535:LYS:HA	1:B:536:LEU:HB2	1.69	0.73
1:B:308:LEU:HB3	1:B:311:CYS:SG	2.30	0.72
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.31	0.69
2:A:601:23E:O6	2:A:601:23E:H7	2.02	0.59
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.89	0.55
1:A:288:SER:CB	1:A:556:SER:HB3	2.37	0.55
1:B:83:LEU:HB2	1:B:173:MET:HA	1.89	0.55
1:A:83:LEU:HB2	1:A:173:MET:HA	1.91	0.53
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.91	0.52
1:A:285:LEU:HD12	1:A:557:GLY:HA3	1.92	0.51
1:B:535:LYS:CA	1:B:536:LEU:HB2	2.36	0.51
1:B:434:LEU:HD11	1:B:511:LEU:HG	1.92	0.51
1:B:554:GLY:O	1:B:555:TYR:HB2	2.11	0.51
1:B:197:PRO:HA	3:B:602:46F:H3	1.93	0.51
1:A:288:SER:HB3	1:A:556:SER:HB3	1.95	0.48
1:B:314:LEU:HB3	1:B:321:VAL:HG23	1.95	0.48
1:A:139:MET:HB3	1:A:160:ILE:HG23	1.95	0.48
1:A:197:PRO:HA	3:A:602:46F:H3	1.95	0.48
1:B:556:SER:N	1:B:557:GLY:HA2	2.28	0.47
1:A:314:LEU:HB3	1:A:321:VAL:HG23	1.96	0.47
1:B:503:ARG:O	1:B:507:VAL:HG23	2.15	0.47
1:B:139:MET:HB3	1:B:160:ILE:HG23	1.96	0.47
1:A:309:GLN:O	1:A:324:CYS:HB2	2.15	0.46
1:B:309:GLN:O	1:B:324:CYS:HB2	2.14	0.46
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.98	0.45
1:B:191:TYR:O	1:B:194:GLN:HG2	2.18	0.44
1:A:191:TYR:O	1:A:194:GLN:HG2	2.18	0.43
1:A:46:SER:O	1:A:50:LYS:HG2	2.19	0.43
1:A:82:LEU:HD13	1:A:249:ALA:HB2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:SER:HB2	1:A:165:LEU:HB3	2.01	0.43
1:B:548:SER:HB2	1:B:551:PHE:HB2	2.00	0.42
1:A:451:CYS:HB2	1:A:552:VAL:O	2.19	0.42
1:B:99:SER:HB2	1:B:165:LEU:HB3	2.02	0.42
1:B:46:SER:O	1:B:50:LYS:HG2	2.19	0.42
1:A:501:ARG:O	1:A:505:ARG:HG3	2.21	0.41
1:B:536:LEU:HA	1:B:536:LEU:HD22	1.88	0.41
1:A:203:PHE:CE2	1:A:314:LEU:HD13	2.56	0.41
1:B:548:SER:HA	1:B:550:TRP:H	1.86	0.41
1:B:388:PRO:O	1:B:392:LEU:HG	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	528/574 (92%)	511 (97%)	13 (2%)	4 (1%)	24 28
1	B	527/574 (92%)	507 (96%)	13 (2%)	7 (1%)	15 15
All	All	1055/1148 (92%)	1018 (96%)	26 (2%)	11 (1%)	19 21

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	147	VAL
1	A	555	TYR
1	B	147	VAL
1	B	536	LEU
1	B	555	TYR
1	B	556	SER
1	A	553	ALA
1	A	557	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	549	SER
1	B	552	VAL
1	B	553	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	449/486 (92%)	423 (94%)	26 (6%)	25 34
1	B	447/486 (92%)	425 (95%)	22 (5%)	31 43
All	All	896/972 (92%)	848 (95%)	48 (5%)	27 38

All (48) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ARG
1	A	96	SER
1	A	150	GLU
1	A	159	LEU
1	A	182	LEU
1	A	201	VAL
1	A	222	ARG
1	A	308	LEU
1	A	309	GLN
1	A	321	VAL
1	A	336	LEU
1	A	341	GLU
1	A	355	LYS
1	A	380	ARG
1	A	381	VAL
1	A	384	LEU
1	A	434	LEU
1	A	440	GLU
1	A	461	GLN
1	A	465	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	487	SER
1	A	545	LEU
1	A	547	LEU
1	A	548	SER
1	A	556	SER
1	A	559	ASP
1	B	56	ARG
1	B	150	GLU
1	B	159	LEU
1	B	182	LEU
1	B	201	VAL
1	B	222	ARG
1	B	308	LEU
1	B	321	VAL
1	B	336	LEU
1	B	341	GLU
1	B	380	ARG
1	B	381	VAL
1	B	384	LEU
1	B	440	GLU
1	B	461	GLN
1	B	487	SER
1	B	506	SER
1	B	531	ARG
1	B	536	LEU
1	B	547	LEU
1	B	555	TYR
1	B	559	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	436	GLN
1	A	514	GLN
1	B	436	GLN
1	B	514	GLN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	23E	A	601	-	46,53,53	1.74	5 (10%)	60,77,77	1.30	6 (10%)
3	46F	A	602	-	38,42,42	1.56	4 (10%)	52,61,61	2.12	15 (28%)
4	SO4	A	651	-	4,4,4	0.14	0	6,6,6	0.19	0
4	SO4	A	652	-	4,4,4	0.29	0	6,6,6	0.13	0
4	SO4	A	653	-	4,4,4	0.69	0	6,6,6	0.46	0
4	SO4	A	654	-	4,4,4	0.04	0	6,6,6	0.07	0
4	SO4	A	659	-	4,4,4	0.35	0	6,6,6	0.31	0
4	SO4	A	660	-	4,4,4	0.15	0	6,6,6	0.28	0
2	23E	B	601	-	46,53,53	1.67	4 (8%)	60,77,77	1.13	3 (5%)
3	46F	B	602	-	38,42,42	1.60	4 (10%)	52,61,61	1.94	13 (25%)
4	SO4	B	651	-	4,4,4	0.20	0	6,6,6	0.11	0
4	SO4	B	652	-	4,4,4	0.72	0	6,6,6	0.31	0
4	SO4	B	653	-	4,4,4	0.30	0	6,6,6	0.15	0
4	SO4	B	657	-	4,4,4	0.11	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	23E	A	601	-	-	0/22/57/57	0/6/7/7
3	46F	A	602	-	-	0/29/45/45	0/4/4/4
4	SO4	A	651	-	-	0/0/0/0	0/0/0/0
4	SO4	A	652	-	-	0/0/0/0	0/0/0/0
4	SO4	A	653	-	-	0/0/0/0	0/0/0/0
4	SO4	A	654	-	-	0/0/0/0	0/0/0/0
4	SO4	A	659	-	-	0/0/0/0	0/0/0/0
4	SO4	A	660	-	-	0/0/0/0	0/0/0/0
2	23E	B	601	-	-	0/22/57/57	0/6/7/7
3	46F	B	602	-	-	0/29/45/45	0/4/4/4
4	SO4	B	651	-	-	0/0/0/0	0/0/0/0
4	SO4	B	652	-	-	0/0/0/0	0/0/0/0
4	SO4	B	653	-	-	0/0/0/0	0/0/0/0
4	SO4	B	657	-	-	0/0/0/0	0/0/0/0

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	23E	C2-C3	-2.09	1.49	1.54
2	A	601	23E	C30-C29	-2.06	1.35	1.41
2	B	601	23E	C24-C23	2.07	1.40	1.36
2	B	601	23E	C32-C30	2.23	1.51	1.49
2	A	601	23E	C16-C2	2.28	1.57	1.54
2	A	601	23E	C32-C30	2.33	1.51	1.49
2	B	601	23E	C25-C22	2.37	1.41	1.37
3	B	602	46F	C21-N28	2.38	1.51	1.48
3	B	602	46F	O31-S38	2.48	1.46	1.43
3	A	602	46F	C19-N28	2.49	1.51	1.48
3	A	602	46F	C21-N28	2.75	1.51	1.48
3	B	602	46F	C17-N29	3.23	1.40	1.33
3	A	602	46F	C17-N29	3.38	1.40	1.33
3	A	602	46F	S38-N28	5.83	1.71	1.63
3	B	602	46F	S38-N28	6.56	1.72	1.63
2	B	601	23E	C29-C28	8.73	1.49	1.40
2	A	601	23E	C29-C28	9.29	1.50	1.40

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	46F	C21-C20-N27	-6.41	97.54	109.69
3	B	602	46F	C21-C20-N27	-5.72	98.84	109.69
3	B	602	46F	CL39-C16-N26	-4.65	111.77	115.16
3	B	602	46F	C19-C18-N27	-4.63	101.45	110.63

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	46F	CL39-C16-N26	-4.57	111.83	115.16
3	A	602	46F	C19-C18-N27	-3.90	102.89	110.63
3	A	602	46F	C21-N28-S38	-3.79	111.76	119.10
3	A	602	46F	C17-C21-N28	-3.36	103.70	111.30
3	A	602	46F	C9-C15-N25	-3.34	118.26	123.64
3	A	602	46F	C8-C14-S38	-3.25	116.20	119.79
2	A	601	23E	C23-C24-C28	-3.25	116.86	121.13
3	A	602	46F	O32-S38-C14	-3.18	103.88	108.00
2	B	601	23E	C29-C28-C27	-3.02	103.72	106.86
2	B	601	23E	C23-C24-C28	-2.93	117.28	121.13
2	A	601	23E	O6-C3-C2	-2.54	117.03	120.97
3	B	602	46F	O32-S38-C14	-2.54	104.71	108.00
2	A	601	23E	C29-C28-C27	-2.51	104.26	106.86
3	B	602	46F	C9-C15-N25	-2.47	119.66	123.64
3	B	602	46F	C21-N28-S38	-2.13	114.98	119.10
3	A	602	46F	C10-C16-N26	-2.12	122.85	124.75
3	B	602	46F	C17-C21-N28	-2.11	106.52	111.30
3	B	602	46F	C11-C23-N29	-2.04	108.21	112.88
3	A	602	46F	C8-C14-C7	2.06	123.17	120.42
3	A	602	46F	C9-C15-N27	2.08	124.99	121.57
3	A	602	46F	C10-C9-C15	2.12	120.58	117.57
3	B	602	46F	C10-C16-CL39	2.23	124.47	119.19
3	B	602	46F	C15-N25-N26	2.24	121.25	119.12
2	A	601	23E	C24-C28-C27	2.57	122.40	119.56
2	A	601	23E	C32-C43-N45	2.58	124.02	120.93
3	A	602	46F	C10-C16-CL39	2.59	125.32	119.19
2	B	601	23E	C24-C28-C27	2.65	122.48	119.56
3	A	602	46F	C15-N25-N26	2.67	121.66	119.12
2	A	601	23E	C39-C32-C43	3.00	120.69	117.92
3	B	602	46F	C18-C19-N28	3.64	112.30	109.07
3	B	602	46F	C23-N29-C17	4.07	127.94	122.34
3	B	602	46F	C20-C21-N28	4.17	113.57	108.96
3	A	602	46F	C20-C21-N28	5.43	114.96	108.96

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	23E	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	46F	1	0
3	B	602	46F	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	534/574 (93%)	-0.02	26 (4%) 33 36	10, 27, 59, 115	0
1	B	533/574 (92%)	0.07	34 (6%) 23 24	12, 31, 76, 114	0
All	All	1067/1148 (92%)	0.03	60 (5%) 28 30	10, 29, 68, 115	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	PRO	8.8
1	B	149	PRO	8.4
1	A	150	GLU	8.3
1	A	152	GLY	7.7
1	B	153	GLY	7.5
1	B	556	SER	7.2
1	A	14	CYS	7.0
1	B	151	LYS	7.0
1	B	152	GLY	6.3
1	A	153	GLY	6.3
1	B	559	ASP	6.0
1	B	150	GLU	5.9
1	A	556	SER	5.9
1	A	151	LYS	5.8
1	B	552	VAL	5.8
1	B	14	CYS	5.6
1	B	146	CYS	5.5
1	B	148	GLN	5.5
1	A	559	ASP	5.5
1	A	148	GLN	5.4
1	A	147	VAL	5.3
1	B	557	GLY	4.6
1	B	154	ARG	4.6
1	B	147	VAL	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	402	HIS	4.3
1	A	10	LEU	3.9
1	A	558	GLY	3.8
1	B	548	SER	3.8
1	B	404	PRO	3.7
1	A	41	THR	3.6
1	A	154	ARG	3.4
1	B	38	TYR	3.3
1	A	555	TYR	3.3
1	B	547	LEU	3.1
1	A	557	GLY	3.1
1	A	7	THR	3.1
1	A	146	CYS	3.1
1	B	555	TYR	3.1
1	A	541	ALA	3.1
1	B	550	TRP	3.0
1	B	541	ALA	2.9
1	B	41	THR	2.9
1	B	558	GLY	2.8
1	B	43	ARG	2.7
1	A	38	TYR	2.6
1	B	549	SER	2.5
1	B	13	PRO	2.4
1	B	42	SER	2.4
1	A	155	LYS	2.3
1	A	45	ALA	2.3
1	B	540	PRO	2.3
1	A	43	ARG	2.3
1	A	13	PRO	2.3
1	B	542	ALA	2.2
1	B	535	LYS	2.2
1	B	110	ASN	2.1
1	A	47	LEU	2.1
1	B	40	THR	2.1
1	B	546	ASP	2.1
1	A	402	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	SO4	B	652	5/5	0.95	0.15	0.02	58,61,64,65	0
4	SO4	A	652	5/5	0.91	0.17	-0.06	97,102,102,103	0
2	23E	A	601	47/47	0.93	0.13	-0.10	12,19,43,55	0
4	SO4	A	653	5/5	0.99	0.11	-0.22	23,26,30,31	0
3	46F	B	602	39/39	0.96	0.12	-0.42	18,31,39,41	0
4	SO4	A	659	5/5	0.97	0.12	-0.48	40,43,45,47	0
3	46F	A	602	39/39	0.97	0.11	-0.49	12,22,33,35	0
2	23E	B	601	47/47	0.94	0.12	-0.78	6,28,48,58	0
4	SO4	B	657	5/5	0.83	0.22	-	117,121,122,123	0
4	SO4	B	651	5/5	0.24	0.60	-	183,188,188,189	0
4	SO4	B	653	5/5	0.98	0.10	-	62,66,67,68	0
4	SO4	A	651	5/5	0.70	0.23	-	121,126,127,127	0
4	SO4	A	660	5/5	0.94	0.21	-	94,98,99,100	0
4	SO4	A	654	5/5	0.58	0.64	-	154,158,159,160	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.