Full wwPDB X-ray Structure Validation Report (i)

Feb 1, 2016 – 07:57 PM GMT

PDB ID	:	4QHR
Title	:	The structure of alanine racemase from Acinetobacter baumannii
Authors	:	Davis, E.; Scaletti-Hutchinson, E.; Nakatani, Y.; Krause, K.L.
Deposited on	:	2014-05-29
Resolution	:	1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4755 (1.90-1.90)
Clashscore	102246	5398 (1.90-1.90)
Ramachandran outliers	100387	5338 (1.90-1.90)
Sidechain outliers	100360	5339 (1.90-1.90)
RSRZ outliers	91569	4766 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	356	3% 94% 6%
1	B	356	% 89% 5% 5%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5902 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alanine racemase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	P	S			
1	A	356	2756	1755	466	514	1	20	0	0	0
1	B	337	2634	1679	445	490	1	19	0	0	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	258	258	258	0	0
2	B	254	254	254	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alanine racemase

Chain A:





- Molecule 1: Alanine racemase

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	46.99Å 82.98Å 93.32Å 90.00° 97.09° 90.00°	Depositor
Resolution (Å)	30.15 – 1.90 30.15 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.2 (30.15-1.90) 99.3 (30.15-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	10.58 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.197 , 0.234 0.210 , 0.244	Depositor DCC
R_{free} test set	2654 reflections (5.01%)	DCC
Wilson B-factor (Å^2)	8.8	Xtriage
Anisotropy	0.327	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.42 , 56.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Outliers	0 of 55666 reflections	Xtriage
F_o,F_c correlation	0.91	EDS
Total number of atoms	5902	wwPDB-VP
Average B, all atoms (Å^2)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.78	0/2787	0.85	4/3780 (0.1%)
1	B	0.79	0/2659	0.88	3/3603 (0.1%)
All	All	0.78	0/5446	0.86	7/7383 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	9	ASP	CB-CG-OD1	7.74	125.27	118.30
1	A	229	ASP	CB-CG-OD1	6.34	124.00	118.30
1	A	9	ASP	CB-CG-OD2	-6.13	112.78	118.30
1	B	19	ARG	NE-CZ-NH1	5.24	122.92	120.30
1	A	303	ASP	CB-CG-OD1	5.24	123.01	118.30
1	B	9	ASP	CB-CG-OD1	5.16	122.94	118.30
1	B	341	TYR	CA-CB-CG	5.09	123.07	113.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2756	0	2703	7	0
1	B	2634	0	2610	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	258	0	0	0	0
2	B	254	0	0	0	0
All	All	5902	0	5313	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:HIS:HD2	1:B:169:LEU:H	1.51	0.57
1:B:276:GLY:HA3	1:B:347:LEU:HA	1.91	0.52
1:A:276:GLY:HA3	1:A:347:LEU:HA	1.93	0.50
1:B:34:LLP:H2'3	1:B:78:ILE:HG12	1.97	0.47
1:A:251:HIS:HA	1:A:258:PHE:O	2.16	0.45
1:A:273:TYR:HA	1:A:277:TYR:O	2.16	0.45
1:A:289:ILE:HD11	1:A:307:ILE:HD12	1.99	0.45
1:A:137:ASP:OD1	1:A:138:PRO:HA	2.17	0.45
1:B:27:SER:OG	1:B:204:ASN:HB3	2.18	0.44
1:B:24:ALA:HA	1:B:204:ASN:ND2	2.34	0.42
1:A:295:PRO:HG2	1:A:308:ASP:HB3	2.02	0.41
1:A:289:ILE:HD11	1:A:307:ILE:CD1	2.51	0.41
1:B:34:LLP:C2'	1:B:78:ILE:HG12	2.51	0.41
1:B:165:VAL:HG23	1:B:167:HIS:HB2	2.01	0.41
1:B:137:ASP:OD1	1:B:138:PRO:HA	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	353/356 (99%)	346 (98%)	6 (2%)	1 (0%)	46	35
1	B	332/356 (93%)	324 (98%)	8 (2%)	0	100	100
All	All	685/712 (96%)	670 (98%)	14 (2%)	1 (0%)	56	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	132	ARG

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/297 (97%)	284 (99%)	4 (1%)	74	71
1	B	279/297 (94%)	275 (99%)	4 (1%)	74	71
All	All	567/594 (96%)	559 (99%)	8 (1%)	74	71

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	HIS
1	A	175	GLN
1	A	185	ASP
1	A	292	LYS
1	B	60	LEU
1	B	101	HIS
1	B	172	GLN
1	B	293	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	244	ASN
1	A	284	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	313	GLN
1	B	167	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
1	LLP	A	34	1	23,24,25	3.24	7 (30%)	28,32,34	1.85	9 (32%)
1	LLP	B	34	1	23,24,25	3.99	7 (30%)	28,32,34	1.68	8 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LLP	A	34	1	-	0/15/17/19	0/1/1/1
1	LLP	B	34	1	-	0/15/17/19	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	34	LLP	C2'-C2	2.02	1.54	1.50
1	A	34	LLP	C6-N1	2.04	1.38	1.34
1	B	34	LLP	C6-N1	2.04	1.38	1.34
1	B	34	LLP	C2'-C2	2.44	1.55	1.50
1	A	34	LLP	C4-C4'	3.49	1.52	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	34	LLP	C4-C4'	4.13	1.53	1.46
1	A	34	LLP	C4-C5	6.32	1.50	1.42
1	A	34	LLP	C4-C3	6.96	1.49	1.40
1	A	34	LLP	C4'-NZ	7.13	1.48	1.27
1	B	34	LLP	C4'-NZ	7.44	1.49	1.27
1	B	34	LLP	C4-C5	8.09	1.52	1.42
1	B	34	LLP	C4-C3	8.49	1.51	1.40
1	A	34	LLP	C3-C2	8.59	1.46	1.40
1	B	34	LLP	C3-C2	11.25	1.48	1.40

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	34	LLP	C5-C6-N1	-2.95	118.74	123.86
1	B	34	LLP	C3-C4-C5	-2.94	115.91	118.11
1	A	34	LLP	C3-C4-C5	-2.65	116.12	118.11
1	A	34	LLP	CG-CD-CE	-2.53	103.63	113.97
1	A	34	LLP	C3-C2-N1	-2.38	117.32	120.61
1	B	34	LLP	CG-CD-CE	-2.33	104.46	113.97
1	A	34	LLP	C4-C4'-NZ	-2.18	112.93	125.06
1	B	34	LLP	C2'-C2-C3	-2.11	118.49	121.04
1	A	34	LLP	C2'-C2-N1	2.14	122.70	117.95
1	B	34	LLP	C2'-C2-N1	2.18	122.78	117.95
1	B	34	LLP	O3-C3-C4	2.31	125.69	119.96
1	A	34	LLP	CE-NZ-C4'	2.33	125.69	118.97
1	B	34	LLP	C5'-C5-C4	2.40	125.51	121.47
1	B	34	LLP	C5-C4-C4'	3.29	126.25	121.52
1	B	34	LLP	C6-N1-C2	3.38	126.17	119.28
1	A	34	LLP	CD-CE-NZ	3.68	117.00	110.98
1	A	34	LLP	C6-N1-C2	4.47	128.40	119.28

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	34	LLP	2	0

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	355/356 (99%)	-0.00	11 (3%) 52 56	6, 12, 30, 46	0
1	B	336/356 (94%)	-0.10	2 (0%) 90 91	5, 12, 25, 34	0
All	All	691/712 (97%)	-0.05	13 (1%) 70 73	5, 12, 28, 46	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	129	GLY	3.6
1	A	257	THR	3.5
1	A	263	ALA	3.2
1	A	311	ASN	3.0
1	A	259	CYS	2.6
1	A	355	GLY	2.6
1	A	258	PHE	2.5
1	A	175	GLN	2.3
1	B	264	MET	2.1
1	A	169	LEU	2.1
1	A	310	THR	2.1
1	B	311	ASN	2.1
1	A	249	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
1	LLP	B	34	24/25	0.94	0.13	-	7,13,17,17	0
1	LLP	A	34	24/25	0.94	0.14	-	5,12,17,18	0

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.