



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:00 PM GMT

PDB ID : 3QOR  
Title : Crystal structure of human nuclear migration protein NudC  
Authors : Derewenda, U.; Derewenda, Z.; Zheng, M.  
Deposited on : 2011-02-10  
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	121	 2% 88% 12%
2	B	121	 4% 94%
3	C	121	 % 87% 10%
4	D	121	 2% 82% 15%
4	E	121	 18% 85% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
5	ACT	A	301	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 5753 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nuclear migration protein nudC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	120	998	636	173	185	4	0	8	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	154	GLY	-	EXPRESSION TAG	UNP Q9Y266
A	155	SER	-	EXPRESSION TAG	UNP Q9Y266
A	156	SER	-	EXPRESSION TAG	UNP Q9Y266
A	157	SER	-	EXPRESSION TAG	UNP Q9Y266
A	236	ALA	GLU	ENGINEERED MUTATION	UNP Q9Y266
A	239	ALA	LYS	ENGINEERED MUTATION	UNP Q9Y266

- Molecule 2 is a protein called Nuclear migration protein nudC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	119	984	626	170	184	4	0	6	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	154	GLY	-	EXPRESSION TAG	UNP Q9Y266
B	155	SER	-	EXPRESSION TAG	UNP Q9Y266
B	156	SER	-	EXPRESSION TAG	UNP Q9Y266
B	157	SER	-	EXPRESSION TAG	UNP Q9Y266
B	236	ALA	GLU	ENGINEERED MUTATION	UNP Q9Y266
B	239	ALA	LYS	ENGINEERED MUTATION	UNP Q9Y266

- Molecule 3 is a protein called Nuclear migration protein nudC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	117	946	603	164	175	4	0	2	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	154	GLY	-	EXPRESSION TAG	UNP Q9Y266
C	155	SER	-	EXPRESSION TAG	UNP Q9Y266
C	156	SER	-	EXPRESSION TAG	UNP Q9Y266
C	157	SER	-	EXPRESSION TAG	UNP Q9Y266
C	236	ALA	GLU	ENGINEERED MUTATION	UNP Q9Y266
C	239	ALA	LYS	ENGINEERED MUTATION	UNP Q9Y266


- Molecule 4 is a protein called Nuclear migration protein nudC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	117	970	615	165	185	5	3	7	0
4	E	118	974	620	168	182	4	0	6	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	154	GLY	-	EXPRESSION TAG	UNP Q9Y266
D	155	SER	-	EXPRESSION TAG	UNP Q9Y266
D	156	SER	-	EXPRESSION TAG	UNP Q9Y266
D	157	SER	-	EXPRESSION TAG	UNP Q9Y266
D	236	ALA	GLU	ENGINEERED MUTATION	UNP Q9Y266
D	239	ALA	LYS	ENGINEERED MUTATION	UNP Q9Y266
E	154	GLY	-	EXPRESSION TAG	UNP Q9Y266
E	155	SER	-	EXPRESSION TAG	UNP Q9Y266
E	156	SER	-	EXPRESSION TAG	UNP Q9Y266
E	157	SER	-	EXPRESSION TAG	UNP Q9Y266
E	236	ALA	GLU	ENGINEERED MUTATION	UNP Q9Y266
E	239	ALA	LYS	ENGINEERED MUTATION	UNP Q9Y266

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	187	Total O 187 187	0	0
6	B	192	Total O 192 192	0	0
6	C	236	Total O 236 236	0	0
6	D	176	Total O 176 176	0	0
6	E	86	Total O 86 86	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Nuclear migration protein nudC




- Molecule 2: Nuclear migration protein nudC




- Molecule 3: Nuclear migration protein nudC



- Molecule 4: Nuclear migration protein nudC



- Molecule 4: Nuclear migration protein nudC



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.73Å 51.82Å 92.85Å 90.00° 90.58° 90.00°	Depositor
Resolution (Å)	28.05 – 1.75 28.05 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.6 (28.05-1.75) 99.7 (28.05-1.75)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.61 (at 1.75Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, $R_{free}$	0.173 , 0.202 0.164 , 0.193	Depositor DCC
$R_{free}$ test set	3225 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	18.8	Xtrriage
Anisotropy	0.704	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 50.0	EDS
Estimated twinning fraction	0.026 for h,-k,-l	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	1 of 63792 reflections (0.002%)	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5753	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 20.86 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 7.9372e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CSD, OCS, CSX, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/1033	0.54	0/1397
2	B	0.33	0/1013	0.51	0/1369
3	C	0.36	0/971	0.54	0/1315
4	D	0.34	0/993	0.54	0/1345
4	E	0.25	0/1001	0.44	0/1353
All	All	0.33	0/5011	0.51	0/6779

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	998	0	1031	15	0
2	B	984	0	1013	2	0
3	C	946	0	961	20	0
4	D	970	0	990	16	0
4	E	974	0	993	13	0
5	A	4	0	3	3	0
6	A	187	0	0	2	0
6	B	192	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	C	236	0	0	8	0
6	D	176	0	0	4	0
6	E	86	0	0	1	0
All	All	5753	0	4991	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:204:ARG:HD3	6:D:402:HOH:O	1.70	0.90
2:B:272:GLU:O	2:B:273:ASN:HB2	1.83	0.76
1:A:227:LYS:NZ	5:A:301:ACT:H1	2.01	0.76
4:D:268:LYS:NZ	4:E:250:LYS:HE3	2.03	0.73
4:D:268:LYS:HZ2	4:E:250:LYS:HE3	1.54	0.73
1:A:272:GLU:O	1:A:273:ASN:HB2	1.87	0.72
4:E:198:MET:HG3	4:E:235[B]:ILE:HD11	1.69	0.72
3:C:228:VAL:HG13	4:E:234[B]:LEU:HD23	1.73	0.71
2:B:199:VAL:HG23	2:B:212[A]:LYS:HG2	1.73	0.71
3:C:160:LYS:HG3	3:C:268:LYS:HE3	1.74	0.70
4:E:188:OCS:OD1	6:E:857:HOH:O	2.10	0.69
3:C:205:ARG:HD2	6:C:855:HOH:O	1.93	0.69
4:D:225:GLU:OE1	6:D:402:HOH:O	2.10	0.68
1:A:227:LYS:HZ3	5:A:301:ACT:H1	1.59	0.67
4:D:205:ARG:HG3	4:D:205:ARG:O	1.95	0.65
3:C:157:SER:N	6:C:373:HOH:O	2.29	0.63
4:D:265:ASN:HD22	4:D:268:LYS:HE2	1.64	0.63
3:C:208:ARG:NH1	6:C:366:HOH:O	2.33	0.62
6:C:869:HOH:O	4:E:234[A]:LEU:HD21	2.01	0.61
4:D:163:LEU:O	4:D:273:ASN:HB3	2.04	0.57
4:E:198:MET:CG	4:E:235[B]:ILE:HD11	2.35	0.57
1:A:184:ALA:O	3:C:251[B]:MET:HE1	2.06	0.56
3:C:268:LYS:NZ	6:C:356:HOH:O	2.40	0.54
1:A:227:LYS:NZ	6:A:121:HOH:O	2.32	0.53
1:A:173:ARG:HG2	3:C:251[B]:MET:HE2	1.91	0.53
1:A:227:LYS:HZ2	5:A:301:ACT:H1	1.74	0.52
1:A:186:PRO:HD3	3:C:251[B]:MET:HE3	1.92	0.52
3:C:265:ASN:HD22	3:C:268:LYS:NZ	2.07	0.52
4:E:194:LYS:N	4:E:194:LYS:HD2	2.25	0.52
1:A:173:ARG:H	3:C:251[B]:MET:HE1	1.75	0.50

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:265:ASN:ND2	4:D:268:LYS:HE2	2.26	0.49
1:A:192[B]:ARG:HG3	1:A:194:LYS:HD3	1.93	0.49
4:E:200:VAL:HG11	4:E:233:TRP:CG	2.49	0.47
1:A:189:VAL:O	3:C:272:GLU:HA	2.15	0.47
4:D:194:LYS:HE3	4:D:196:LYS:HE3	1.96	0.47
4:D:223:TYR:HB3	4:D:254:TRP:CE3	2.51	0.46
1:A:234[A]:LEU:HD11	6:A:82:HOH:O	2.15	0.46
4:D:242[A]:THR:HG21	6:D:466:HOH:O	2.18	0.44
1:A:173:ARG:N	3:C:251[B]:MET:HE1	2.32	0.44
3:C:157:SER:HB3	6:C:349:HOH:O	2.16	0.44
4:D:194:LYS:CE	4:D:196:LYS:HE3	2.48	0.44
1:A:234[B]:LEU:HD13	1:A:242:THR:OG1	2.18	0.44
3:C:265:ASN:HD22	3:C:268:LYS:HZ2	1.66	0.43
3:C:193:LEU:HD11	3:C:211:LEU:HD21	1.99	0.43
4:E:177:THR:OG1	4:E:180[B]:GLU:HG2	2.19	0.43
4:E:265:ASN:OD1	4:E:267:LYS:HB2	2.19	0.42
4:D:212:LYS:NZ	6:D:472:HOH:O	2.51	0.42
1:A:199:VAL:HG11	1:A:208:ARG:CZ	2.51	0.41
4:E:256:ARG:HD3	4:E:261:ASP:O	2.21	0.41
4:D:169[B]:LEU:HD22	4:D:172:TYR:CZ	2.56	0.41
3:C:205:ARG:NH1	6:C:855:HOH:O	2.52	0.40
3:C:221:GLU:CD	4:D:170:PRO:HG3	2.40	0.40
4:D:255[A]:SER:OG	4:D:263:GLU:HB3	2.21	0.40
3:C:268:LYS:HE2	3:C:268:LYS:HB2	1.88	0.40
3:C:157:SER:CA	6:C:373:HOH:O	2.70	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	125/121 (103%)	123 (98%)	2 (2%)	0	<a href="#">100</a> <a href="#">100</a>

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	122/121 (101%)	120 (98%)	2 (2%)	0	100	100
3	C	117/121 (97%)	116 (99%)	1 (1%)	0	100	100
4	D	120/121 (99%)	116 (97%)	4 (3%)	0	100	100
4	E	121/121 (100%)	115 (95%)	6 (5%)	0	100	100
All	All	605/605 (100%)	590 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	114/107 (106%)	112 (98%)	2 (2%)	66	46
2	B	112/107 (105%)	111 (99%)	1 (1%)	84	72
3	C	106/108 (98%)	106 (100%)	0	100	100
4	D	110/107 (103%)	110 (100%)	0	100	100
4	E	109/107 (102%)	104 (95%)	5 (5%)	33	10
All	All	551/536 (103%)	543 (98%)	8 (2%)	82	55

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	180[A]	GLU
1	A	180[B]	GLU
2	B	192	ARG
4	E	180[A]	GLU
4	E	180[B]	GLU
4	E	192[A]	ARG
4	E	192[B]	ARG
4	E	194	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	273	ASN
2	B	273	ASN
3	C	265	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

5 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CSX	A	188	1	3,6,7	0.58	0	3,6,8	1.20	0
2	CSD	B	188	2	3,7,8	1.06	0	3,8,10	4.78	1 (33%)
4	OCS	D	188[A]	-	7,8,9	1.50	1 (14%)	7,11,13	1.80	2 (28%)
4	OCS	D	188[B]	-	7,8,9	1.07	0	7,11,13	1.72	2 (28%)
4	OCS	E	188	4	7,8,9	1.30	1 (14%)	7,11,13	2.19	2 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSX	A	188	1	-	0/1/5/7	0/0/0/0
2	CSD	B	188	2	-	0/2/6/8	0/0/0/0
4	OCS	D	188[A]	-	-	1/4/7/9	0/0/0/0
4	OCS	D	188[B]	-	-	0/4/7/9	0/0/0/0
4	OCS	E	188	4	-	1/4/7/9	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	E	188	OCS	CB-SG	2.67	1.81	1.77
4	D	188[A]	OCS	CB-SG	3.05	1.82	1.77

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	188[A]	OCS	OD3-SG-CB	2.41	108.97	106.94
4	D	188[B]	OCS	OD3-SG-CB	2.62	109.15	106.94
4	D	188[B]	OCS	OD1-SG-CB	2.91	109.39	106.94
4	E	188	OCS	OD3-SG-CB	3.51	109.90	106.94
4	D	188[A]	OCS	OD1-SG-CB	3.51	109.90	106.94
4	E	188	OCS	OD1-SG-CB	3.71	110.07	106.94
2	B	188	CSD	OD1-SG-CB	8.04	118.80	105.40

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	E	188	OCS	SG-CB-CA-N
4	D	188[A]	OCS	SG-CB-CA-N

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	188	OCS	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all  $Z$  scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	ACT	A	301	-	1,3,3	1.48	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	ACT	A	301	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	301	ACT	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	119/121 (98%)	0.14	2 (1%) 73 80	13, 18, 40, 45	0
2	B	118/121 (97%)	0.32	5 (4%) 40 46	15, 21, 40, 52	0
3	C	117/121 (96%)	-0.08	1 (0%) 85 89	13, 20, 36, 52	0
4	D	116/121 (95%)	0.07	3 (2%) 59 65	14, 20, 37, 53	0
4	E	117/121 (96%)	1.16	22 (18%) 2 2	28, 38, 57, 84	0
All	All	587/605 (97%)	0.32	33 (5%) 28 33	13, 22, 47, 84	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	E	190	ASN	5.8
4	E	160	LYS	5.6
4	E	196	LYS	5.4
4	E	159	LEU	5.2
4	D	157	SER	4.8
4	E	191	PHE	4.6
4	E	157	SER	4.5
4	E	189	VAL	4.3
4	D	273	ASN	4.3
1	A	155	SER	4.1
4	E	267	LYS	3.5
2	B	190	ASN	3.3
2	B	273	ASN	3.1
4	E	229	GLU	3.1
4	E	215	PRO	3.0
4	E	216	ALA	2.9
4	E	201	ASP	2.8
4	E	200	VAL	2.8
2	B	189	VAL	2.6
2	B	155	SER	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
4	E	158	LYS	2.5
1	A	273	ASN	2.5
4	E	214	GLN	2.4
3	C	272	GLU	2.4
4	D	192	ARG	2.4
4	E	248	ILE	2.3
4	E	194	LYS	2.3
2	B	251[A]	MET	2.3
4	E	169	LEU	2.2
4	E	213	GLY	2.1
4	E	260	SER	2.1
4	E	195	GLY	2.1
4	E	251[A]	MET	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	OCS	E	188	9/10	0.83	0.34	-	43,46,61,61	3
4	OCS	D	188[B]	9/10	0.94	0.12	-	18,26,30,35	4
4	OCS	D	188[A]	9/10	0.94	0.12	-	18,26,39,39	4
1	CSX	A	188	7/8	0.93	0.07	-	17,25,35,38	0
2	CSD	B	188	8/9	0.89	0.13	-	21,26,40,47	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q < 0.9
5	ACT	A	301	4/4	0.84	0.15	-	26,33,36,45	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.