



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 07:12 PM GMT

PDB ID : 4QOF  
Title : Crystal structure of fmn quinone reductase 2 AT 1.55Å  
Authors : Serriere, J.; Boutin, J.A.; Isabet, T.; Antoine, M.; Ferry, G.  
Deposited on : 2014-06-20  
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1665 (1.58-1.54)
Clashscore	102246	1014 (1.56-1.56)
Ramachandran outliers	100387	1704 (1.58-1.54)
Sidechain outliers	100360	1702 (1.58-1.54)
RSRZ outliers	91569	1668 (1.58-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	B	231	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 4312 atoms, of which 38 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosyldihyronicotinamide dehydrogenase [quinone].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	0	6	0
			1865	1201	309	346	9			
1	B	228	Total	C	N	O	S	0	18	0
			1957	1264	321	362	10			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	46	PHE	LEU	VARIANT	UNP P16083
B	46	PHE	LEU	VARIANT	UNP P16083

- Molecule 2 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	H	N	O	0	0
			50	17	19	4	9		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	B	1	Total	C	H	N	O	P	0	0
			50	17	19	4	9	1		

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	A	1	Total	Zn	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	222	Total	O	0	0
			222	222		
4	B	166	Total	O	0	0
			166	166		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribosyldihydropyridine dehydrogenase [quinone]



- Molecule 1: Ribosyldihydropyridine dehydrogenase [quinone]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.17Å 83.27Å 106.44Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.69 – 1.55 46.57 – 1.55	Depositor EDS
% Data completeness (in resolution range)	99.8 (22.69-1.55) 99.3 (46.57-1.55)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.97 (at 1.55Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, $R_{free}$	0.214 , 0.235 0.218 , 0.233	Depositor DCC
$R_{free}$ test set	3648 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	16.9	Xtriage
Anisotropy	0.593	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 39.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 73159 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4312	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.53	0/1916	0.60	0/2597
1	B	0.49	0/2012	0.59	0/2727
All	All	0.51	0/3928	0.59	0/5324

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1865	0	1811	16	0
1	B	1957	0	1894	9	0
2	A	31	19	19	0	0
2	B	31	19	19	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	222	0	0	10	0
4	B	166	0	0	3	0
All	All	4274	38	3743	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:157:LYS:HE2	4:A:617:HOH:O	1.39	1.20
1:A:193:GLU:CD	4:A:620:HOH:O	1.93	1.07
1:A:85[B]:THR:HG23	4:A:622:HOH:O	1.53	1.06
1:A:193:GLU:CG	4:A:620:HOH:O	2.20	0.89
1:B:193:GLU:CD	4:B:560:HOH:O	2.13	0.86
1:B:193:GLU:CG	4:B:560:HOH:O	2.32	0.76
1:A:85[A]:THR:HG22	4:A:622:HOH:O	1.94	0.68
1:B:193:GLU:HG2	4:B:560:HOH:O	1.98	0.61
1:A:193:GLU:HG2	4:A:620:HOH:O	1.97	0.58
1:B:122:GLN:NE2	1:B:128[A]:ILE:H	2.06	0.54
1:A:82:SER:HA	1:A:85[B]:THR:HG22	1.89	0.54
1:A:193:GLU:OE2	4:A:620:HOH:O	2.16	0.53
1:B:122:GLN:HG3	1:B:127[A]:ASP:HA	1.92	0.51
1:B:4[B]:LYS:HG2	1:B:35:THR:HB	1.92	0.51
1:A:4[B]:LYS:HG2	1:A:35:THR:HB	1.95	0.48
1:A:229:GLY:O	1:A:230:GLN:HB2	2.14	0.47
1:A:85[A]:THR:CG2	4:A:622:HOH:O	2.59	0.46
1:A:157:LYS:CE	4:A:617:HOH:O	2.23	0.46
1:B:125:ALA:HB1	1:B:179:CYS:SG	2.59	0.42
1:A:165:ARG:HA	1:A:168:LEU:HD12	2.01	0.42
1:B:169:TRP:HB3	1:B:170:PRO:HD3	2.01	0.42
1:A:169:TRP:HB3	1:A:170:PRO:HD3	2.01	0.41
1:A:157:LYS:CD	4:A:617:HOH:O	2.65	0.41
1:A:24:VAL:HG11	1:A:204:VAL:CG1	2.50	0.41
1:B:165:ARG:HA	1:B:168:LEU:HD12	2.02	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles ⓘ

### 5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	232/231 (100%)	223 (96%)	9 (4%)	0	100	100
1	B	244/231 (106%)	234 (96%)	8 (3%)	2 (1%)	24	5
All	All	476/462 (103%)	457 (96%)	17 (4%)	2 (0%)	52	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	130[A]	GLY
1	B	130[B]	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	200/195 (103%)	198 (99%)	2 (1%)	82	62
1	B	210/195 (108%)	207 (99%)	3 (1%)	74	48
All	All	410/390 (105%)	405 (99%)	5 (1%)	82	54

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	127	ASP
1	A	198	GLU
1	B	127[A]	ASP
1	B	127[B]	ASP
1	B	198	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	72	HIS
1	B	122	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMN	A	301	-	31,33,33	1.22	5 (16%)	32,50,50	1.45	2 (6%)
2	FMN	B	301	-	31,33,33	1.21	5 (16%)	32,50,50	1.52	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	301	-	-	0/18/18/18	0/3/3/3
2	FMN	B	301	-	-	0/18/18/18	0/3/3/3

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	301	FMN	C4-N3	2.09	1.39	1.36

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	FMN	C4-N3	2.31	1.40	1.36
2	B	301	FMN	C5A-N5	2.49	1.39	1.35
2	B	301	FMN	C4-C4A	2.70	1.46	1.40
2	A	301	FMN	C5A-N5	2.76	1.39	1.35
2	A	301	FMN	C4A-C10	2.78	1.46	1.40
2	A	301	FMN	C4-C4A	2.82	1.46	1.40
2	A	301	FMN	C9A-N10	2.85	1.42	1.38
2	B	301	FMN	C4A-C10	3.05	1.46	1.40
2	B	301	FMN	C9A-N10	3.25	1.43	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	FMN	C4A-C10-N10	-5.41	116.59	120.52
2	A	301	FMN	C4A-C10-N10	-4.94	116.93	120.52
2	B	301	FMN	O5'-P-O1P	2.68	113.83	107.08
2	B	301	FMN	C2-N1-C10	5.44	118.03	113.39
2	A	301	FMN	C2-N1-C10	5.87	118.40	113.39

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	228/231 (98%)	1.05	26 (11%)	7 6	10, 18, 40, 72	0
1	B	228/231 (98%)	1.43	47 (20%)	1 1	10, 25, 54, 88	0
All	All	456/462 (98%)	1.24	73 (16%)	3 2	10, 20, 51, 88	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	230	GLN	7.3
1	B	131[A]	PHE	7.3
1	B	128[A]	ILE	7.1
1	B	57	GLY	5.3
1	B	129[A]	PRO	5.1
1	B	130[A]	GLY	5.0
1	B	58	THR	4.7
1	B	157	LYS	4.6
1	B	230	GLN	4.5
1	B	196	SER	4.5
1	B	198	GLU	4.4
1	B	59	LEU	4.2
1	B	228	PHE	4.1
1	B	197	GLU	3.9
1	A	198	GLU	3.8
1	A	197	GLU	3.5
1	A	158	THR	3.5
1	A	160	VAL	3.4
1	B	132[A]	TYR	3.4
1	B	160	VAL	3.3
1	B	64	VAL	3.3
1	A	196	SER	3.3
1	B	222	CYS	3.3
1	B	134[A]	SER	3.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	153	GLU	3.2
1	A	24	VAL	3.2
1	A	128	ILE	3.2
1	B	136[A]	LEU	3.2
1	B	158	THR	3.1
1	A	159	GLY	3.1
1	B	135[A]	GLY	3.1
1	B	229	GLY	3.1
1	B	56	THR	3.1
1	B	225	HIS	3.1
1	B	126[A]	PHE	3.0
1	A	85[A]	THR	3.0
1	B	205	ALA	3.0
1	A	62	PRO	3.0
1	A	157	LYS	2.9
1	B	194	ILE	2.9
1	A	213	THR	2.9
1	B	201	LYS	2.9
1	B	209	GLN	2.9
1	B	133[A]	ASP	2.8
1	A	222	CYS	2.8
1	A	64	VAL	2.8
1	B	204	VAL	2.7
1	A	201	LYS	2.7
1	B	226	TRP	2.6
1	B	193	GLU	2.5
1	B	123	GLY	2.5
1	B	195	ALA	2.5
1	A	209	GLN	2.5
1	B	82	SER	2.5
1	B	151	THR	2.5
1	A	56	THR	2.4
1	B	61	ASN	2.4
1	B	178	PHE	2.3
1	B	218	GLU	2.3
1	B	153	GLU	2.3
1	B	122	GLN	2.2
1	A	194	ILE	2.2
1	A	207	TRP	2.2
1	A	199	GLU	2.2
1	A	195	ALA	2.1
1	A	192	PRO	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	137	LEU	2.1
1	A	215	TRP	2.1
1	A	176	LEU	2.1
1	B	214	ILE	2.1
1	B	219	PRO	2.1
1	B	169	TRP	2.1
1	B	106[A]	PHE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	FMN	B	301	31/31	0.90	0.12	-0.48	12,14,19,20	0
2	FMN	A	301	31/31	0.90	0.11	-0.74	15,17,21,24	0
3	ZN	B	302	1/1	0.94	0.10	-2.42	24,24,24,24	0
3	ZN	A	302	1/1	0.98	0.12	-	17,17,17,17	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.