



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 12:01 PM GMT

PDB ID : 3QQ4

Title : Crystal structure of swine major histocompatibility complex class I SLA-1 0401 and identification of 2009 pandemic swine-origin influenza A H1N1 virus cytotoxic T lymphocyte epitope peptides

Authors : Zhang, N.; Qi, J.; Gao, F.; Pan, X.; Chen, R.; Li, Q.; Chen, Z.; Li, X.; Xia, C.; Gao, G.F.

Deposited on : 2011-02-15

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.7 (RC4), CSD as536be (2015)

Xtriage (Phenix) : 1.9-1692

EDS : rb-20026688

Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)

Refmac : 5.8.0135

CCP4 : 6.5.0

Ideal geometry (proteins) : Engh & Huber (2001)

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MHC class I antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C 2228	N 1392	O 395	S 432	9	0	0

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	99	Total	C 817	N 524	O 142	S 148	3	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	GLU	-	EXPRESSION TAG	UNP Q07717
B	2	PHE	-	EXPRESSION TAG	UNP Q07717

- Molecule 3 is a protein called VP35.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C 61	N 37	O 9	O 15	0	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	237	Total	O 237	0	0
4	B	90	Total	O 90	0	0
4	C	3	Total	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: MHC class I antigen



- Molecule 2: Beta-2-microglobulin



- Molecule 3: VP35



4 Data and refinement statistics (i)

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	88.68 Å 40.24 Å 103.63 Å 90.00° 89.93° 90.00°	Depositor
Resolution (Å)	23.82 – 2.10 34.55 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.3 (23.82-2.10) 99.3 (34.55-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	6.43 (at 2.10 Å)	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
R , R_{free}	0.192 , 0.233 0.192 , 0.231	Depositor DCC
R_{free} test set	1109 reflections (5.13%)	DCC
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.141	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.7	EDS
Estimated twinning fraction	0.048 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 21639 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3436	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.26	0/2288	0.65	8/3108 (0.3%)
2	B	0.22	0/843	0.41	0/1145
3	C	0.25	0/61	0.43	0/82
All	All	0.25	0/3192	0.59	8/4335 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	20	SER	CB-CA-C	12.69	134.21	110.10
1	A	226	GLN	N-CA-CB	-11.33	90.20	110.60
1	A	16	ASP	N-CA-CB	9.77	128.18	110.60
1	A	225	SER	N-CA-C	7.85	132.20	111.00
1	A	20	SER	C-N-CA	5.77	136.13	121.70
1	A	195	SER	N-CA-C	-5.47	96.23	111.00
1	A	226	GLN	N-CA-C	5.35	125.44	111.00
1	A	16	ASP	CB-CG-OD2	5.20	122.98	118.30

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	15	PRO	Mainchain,Peptide
1	A	17	ARG	Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	20	SER	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2228	0	2087	49	2
2	B	817	0	788	18	0
3	C	61	0	56	3	0
4	A	237	0	0	3	2
4	B	90	0	0	0	0
4	C	3	0	0	1	0
All	All	3436	0	2931	65	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:ASP:HB2	1:A:18:GLY:CA	1.08	1.55
1:A:16:ASP:CB	1:A:18:GLY:HA2	1.37	1.48
1:A:16:ASP:CB	1:A:18:GLY:CA	2.03	1.25
1:A:16:ASP:HB2	1:A:18:GLY:HA3	1.13	1.08
2:B:6:PRO:HA	2:B:87:THR:HG21	1.42	0.98
1:A:16:ASP:HA	1:A:17:ARG:HB2	1.44	0.98
1:A:15:PRO:HA	1:A:16:ASP:HB3	1.52	0.90
1:A:16:ASP:HB3	1:A:18:GLY:HA2	1.54	0.86
1:A:16:ASP:N	1:A:17:ARG:HG3	1.94	0.82
4:A:506:HOH:O	3:C:7:GLU:HB2	1.80	0.79
1:A:16:ASP:HA	1:A:17:ARG:CB	2.12	0.79
1:A:196:ASP:O	1:A:251:PRO:HG2	1.87	0.74
1:A:16:ASP:CB	1:A:18:GLY:HA3	1.94	0.73
1:A:16:ASP:HB2	1:A:18:GLY:HA2	0.73	0.72
1:A:16:ASP:CA	1:A:17:ARG:CB	2.74	0.65
1:A:21:ARG:NE	1:A:39:TYR:HB2	2.12	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:ARG:HH12	2:B:100:HIS:CD2	2.17	0.62
1:A:234:ARG:HH12	2:B:100:HIS:HD2	1.48	0.61
1:A:93:HIS:HD2	1:A:119:ASP:OD2	1.84	0.60
2:B:71:PHE:CZ	2:B:73:PRO:HG3	2.37	0.60
1:A:15:PRO:HA	1:A:16:ASP:CB	2.21	0.59
1:A:178:THR:O	1:A:181:ARG:HG3	2.05	0.56
1:A:98:MET:HG2	1:A:113:TYR:HE1	1.72	0.54
1:A:202:ARG:HD3	1:A:244:TRP:CE3	2.42	0.54
1:A:117:ALA:HB2	2:B:61:TRP:CE2	2.42	0.54
2:B:19:ASN:HA	2:B:73:PRO:O	2.09	0.53
1:A:170:ARG:HD3	4:A:286:HOH:O	2.08	0.53
1:A:81:LEU:HD12	1:A:118:TYR:CD1	2.44	0.52
1:A:199:VAL:HG12	1:A:249:VAL:O	2.09	0.52
1:A:15:PRO:C	1:A:17:ARG:HG3	2.30	0.51
2:B:85:HIS:ND1	2:B:87:THR:HG22	2.25	0.51
1:A:16:ASP:CA	1:A:18:GLY:HA2	2.31	0.51
2:B:7:PRO:HD3	2:B:87:THR:HG21	1.93	0.51
3:C:7:GLU:HA	4:C:227:HOH:O	2.11	0.51
1:A:114:ARG:HH12	3:C:6:THR:HB	1.76	0.50
2:B:35:PRO:HG3	2:B:63:PHE:CE2	2.47	0.49
1:A:44:ARG:HD2	1:A:64:THR:HG21	1.96	0.47
1:A:150:ALA:O	1:A:151:ASP:HB2	2.14	0.47
1:A:63:GLU:O	1:A:67:VAL:HG22	2.15	0.47
1:A:33:PHE:HA	1:A:49:VAL:HG22	1.96	0.47
2:B:90:LYS:HB2	2:B:91:PRO:HD2	1.95	0.47
2:B:7:PRO:HD3	2:B:87:THR:CG2	2.45	0.47
2:B:88:LEU:HD13	2:B:92:LYS:HB2	1.98	0.46
1:A:15:PRO:O	1:A:17:ARG:HG3	2.16	0.45
1:A:98:MET:HE2	2:B:61:TRP:CH2	2.51	0.45
1:A:16:ASP:CG	1:A:18:GLY:HA3	2.36	0.45
1:A:231:VAL:HG11	1:A:244:TRP:CZ2	2.52	0.45
2:B:27:CYS:HB2	2:B:41:LEU:HD21	1.98	0.44
1:A:219:ARG:HB2	1:A:224:GLN:CD	2.38	0.44
2:B:59:LYS:HD3	2:B:59:LYS:O	2.16	0.44
1:A:37:ASP:OD2	1:A:39:TYR:HB3	2.18	0.44
1:A:49:VAL:O	1:A:52:ILE:HG22	2.18	0.44
1:A:93:HIS:HE1	4:A:279:HOH:O	2.01	0.44
1:A:202:ARG:HG2	1:A:204:TRP:NE1	2.33	0.44
1:A:183:GLU:HA	1:A:184:PRO:HD2	1.87	0.43
2:B:26:ASN:HD22	2:B:68:HIS:HB3	1.83	0.43
1:A:13:SER:O	1:A:92:SER:HB2	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:178:THR:HG23	1:A:179:LEU:N	2.34	0.43
1:A:35:ARG:NH1	1:A:48:ARG:NH2	2.67	0.42
2:B:35:PRO:HG3	2:B:63:PHE:CZ	2.55	0.41
1:A:202:ARG:HD3	1:A:244:TRP:CD2	2.56	0.41
1:A:107:GLY:O	1:A:169:ARG:NH1	2.54	0.41
2:B:21:LYS:HA	2:B:22:PRO:HD2	1.80	0.41
1:A:185:PRO:HB3	1:A:208:PHE:HB3	2.03	0.40
1:A:82:ARG:HD2	1:A:89:GLU:HA	2.04	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:ASP:OD2	4:A:472:HOH:O[2_554]	1.70	0.50
1:A:17:ARG:NH2	4:A:462:HOH:O[3_455]	1.92	0.28

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/275 (99%)	260 (95%)	13 (5%)	0	100 100
2	B	97/100 (97%)	91 (94%)	6 (6%)	0	100 100
3	C	7/9 (78%)	7 (100%)	0	0	100 100
All	All	377/384 (98%)	358 (95%)	19 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	233/234 (100%)	221 (95%)	12 (5%)	29 25
2	B	92/93 (99%)	90 (98%)	2 (2%)	60 64
3	C	4/4 (100%)	4 (100%)	0	100 100
All	All	329/331 (99%)	315 (96%)	14 (4%)	35 34

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	ARG
1	A	35	ARG
1	A	65	ARG
1	A	86	ASN
1	A	138	MET
1	A	154	GLU
1	A	156	ARG
1	A	181	ARG
1	A	212	GLU
1	A	219	ARG
1	A	227	ASP
1	A	275	ASP
2	B	71	PHE
2	B	89	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	72	GLN
1	A	86	ASN
1	A	87	GLN
1	A	93	HIS
1	A	96	GLN
1	A	141	GLN
1	A	192	HIS
1	A	262	GLN
2	B	26	ASN
2	B	49	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.16	23 (8%) 14 19	17, 28, 60, 70	0
2	B	99/100 (99%)	0.00	2 (2%) 68 73	20, 33, 50, 54	0
3	C	9/9 (100%)	-0.08	0 100 100	18, 21, 33, 34	0
All	All	383/384 (99%)	0.12	25 (6%) 22 29	17, 30, 56, 70	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	16	ASP	10.5
1	A	197	LEU	5.4
1	A	196	ASP	4.7
1	A	275	ASP	4.5
1	A	194	SER	4.3
1	A	227	ASP	4.3
1	A	195	SER	4.2
1	A	20	SER	3.5
2	B	2	PHE	3.5
2	B	3	VAL	3.3
1	A	226	GLN	3.0
1	A	181	ARG	2.9
1	A	199	VAL	2.8
1	A	255	GLN	2.5
1	A	222	GLN	2.5
1	A	39	TYR	2.4
1	A	193	PRO	2.3
1	A	248	VAL	2.3
1	A	250	PRO	2.3
1	A	253	GLU	2.3
1	A	17	ARG	2.3
1	A	220	GLU	2.2
1	A	247	LEU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	219	ARG	2.1
1	A	274	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.