



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:21 PM GMT

PDB ID : 3QZ7
Title : T-3 ternary complex of Dpo4
Authors : Pata, J.D.; Wu, Y.; Wilson, R.C.
Deposited on : 2011-03-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	<div> <div>23%</div> <div>82%</div> <div>12%</div> <div>• 5%</div> </div>
2	P	13	<div> <div>77%</div> <div>15%</div> <div>8%</div> </div>
3	T	19	<div> <div>53%</div> <div>21%</div> <div>5%</div> <div>21%</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6635 atoms, of which 3219 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase IV.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	341	Total	C	H	N	O	S	0	0	0
			5634	1760	2891	472	504	7			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	GLY	-	EXPRESSION TAG	UNP Q97W02
A	354	GLY	-	EXPRESSION TAG	UNP Q97W02
A	355	HIS	-	EXPRESSION TAG	UNP Q97W02
A	356	HIS	-	EXPRESSION TAG	UNP Q97W02
A	357	HIS	-	EXPRESSION TAG	UNP Q97W02
A	358	HIS	-	EXPRESSION TAG	UNP Q97W02
A	359	HIS	-	EXPRESSION TAG	UNP Q97W02
A	360	HIS	-	EXPRESSION TAG	UNP Q97W02


- Molecule 2 is a DNA chain called 5'-D(*GP*GP*CP*AP*CP*TP*GP*AP*TP*CP*GP*GP*G)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	P	13	Total	C	H	N	O	P	0	0	0
			415	127	147	53	76	12			

- Molecule 3 is a DNA chain called 5'-D(*TP*TP*AP*CP*GP*CP*CP*TP*CP*GP*AP*TP*CP*AP*GP*TP*GP*CP*C)-3'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	T	15	Total	C	H	N	O	P	0	0	0
			473	144	169	54	91	15			

- Molecule 4 is 2'-DEOXYCYTIDINE-5'-TRIPHOSPHATE (three-letter code: DCP) (formula: C₉H₁₆N₃O₁₃P₃).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	0	0
			40	9	12	3	13	3		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	3	Total Ca 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	56	Total O 56 56	0	0
6	P	6	Total O 6 6	0	0
6	T	8	Total O 8 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DNA polymerase IV



• Molecule 2: 5'-D(*GP*GP*CP*AP*CP*TP*GP*AP*TP*CP*GP*GP*G)-3'



• Molecule 3: 5'-D(*TP*TP*AP*CP*GP*CP*CP*TP*CP*GP*AP*TP*CP*AP*GP*TP*GP*CP*C)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	98.71Å 102.46Å 52.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.54 – 2.00 25.54 – 2.00	Depositor EDS
% Data completeness (in resolution range)	90.0 (25.54-2.00) 90.0 (25.54-2.00)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 1.99Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.227 , 0.267 0.228 , 0.266	Depositor DCC
R_{free} test set	1629 reflections (5.14%)	DCC
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.152	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.45 , 53.9	EDS
Estimated twinning fraction	0.014 for k,h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 33331 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6635	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.92% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, DCP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.37	0/2782	0.52	0/3736
2	P	0.69	0/301	1.43	3/464 (0.6%)
3	T	0.70	0/339	1.47	6/520 (1.2%)
All	All	0.45	0/3422	0.81	9/4720 (0.2%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	T	16	DT	O4'-C4'-C3'	-8.59	100.84	106.00
3	T	16	DT	O4'-C1'-N1	7.09	112.96	108.00
3	T	9	DC	O4'-C1'-N1	7.08	112.96	108.00
2	P	6	DT	O4'-C1'-N1	-6.38	103.53	108.00
3	T	5	DG	O4'-C1'-N9	-6.04	103.78	108.00
2	P	12	DG	O4'-C1'-N9	5.61	111.93	108.00
2	P	12	DG	C1'-O4'-C4'	-5.23	104.87	110.10
3	T	10	DG	C5-C6-O6	-5.18	125.49	128.60
3	T	10	DG	N1-C6-O6	5.00	122.90	119.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2743	2891	2889	29	0
2	P	268	147	147	1	0
3	T	304	169	169	2	0
4	A	28	12	12	1	0
5	A	3	0	0	0	0
6	A	56	0	0	0	0
6	P	6	0	0	0	0
6	T	8	0	0	0	0
All	All	3416	3219	3217	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MET:HB2	1:A:112:SER:OG	1.82	0.79
3:T:16:DT:H2"	3:T:17:DG:O5'	2.03	0.59
1:A:247:ARG:C	1:A:248:ILE:HD12	2.23	0.58
3:T:16:DT:H2'	3:T:17:DG:C8	2.40	0.57
1:A:122:TYR:CE2	1:A:126:LEU:HD11	2.39	0.57
1:A:186:ILE:HD11	1:A:225:LEU:HD21	1.86	0.56
1:A:133:LEU:O	1:A:137:LYS:HD3	2.05	0.56
1:A:108:TYR:C	1:A:109:LEU:HD12	2.26	0.56
1:A:289:VAL:HB	1:A:332:ARG:HB2	1.89	0.53
1:A:185:GLY:O	1:A:221:LYS:CE	2.59	0.50
1:A:251:MET:HG2	1:A:264:TYR:CD2	2.47	0.49
1:A:100:GLU:HB2	1:A:237:ILE:HG23	1.93	0.49
1:A:10:TYR:HA	4:A:361:DCP:PB	2.54	0.48
2:P:6:DT:H2"	2:P:7:DG:C8	2.50	0.47
1:A:260:GLU:O	1:A:263:PRO:HD2	2.15	0.47
1:A:67:ILE:HG22	1:A:68:LEU:HG	1.96	0.46
1:A:9:ASP:O	1:A:10:TYR:C	2.54	0.46
1:A:122:TYR:HE2	1:A:126:LEU:HD11	1.80	0.45
1:A:241:VAL:HG23	1:A:241:VAL:O	2.17	0.45
1:A:248:ILE:HD12	1:A:248:ILE:N	2.32	0.44
1:A:180:ILE:HD13	1:A:199:ILE:HG22	1.99	0.44
1:A:31:CYS:HB3	1:A:61:ILE:HD11	1.99	0.44
1:A:79:GLU:CD	1:A:79:GLU:H	2.21	0.43
1:A:248:ILE:HA	1:A:334:GLY:HA3	2.01	0.42
1:A:109:LEU:HD12	1:A:109:LEU:N	2.35	0.42
1:A:160:PRO:O	1:A:161:ASN:C	2.58	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:197:LEU:HD22	1:A:212:LYS:HZ3	1.84	0.42
1:A:129:LYS:HE3	1:A:129:LYS:HB3	1.89	0.42
1:A:185:GLY:O	1:A:221:LYS:HE2	2.19	0.41
1:A:31:CYS:HB3	1:A:61:ILE:CD1	2.51	0.40
1:A:300:ARG:HD3	1:A:302:PHE:CZ	2.56	0.40
1:A:180:ILE:HD13	1:A:194:LEU:HD13	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	339/360 (94%)	325 (96%)	13 (4%)	1 (0%)	46 41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	10	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	300/315 (95%)	292 (97%)	8 (3%)	52 52

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	10	TYR
1	A	127	GLU
1	A	180	ILE
1	A	234	ASN
1	A	252	LYS
1	A	292	ASP
1	A	294	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 3 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	DCP	A	361	5	21,29,29	2.10	6 (28%)	33,45,45	2.40	11 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DCP	A	361	5	-	0/18/34/34	0/2/2/2

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	361	DCP	C3'-C4'	-2.64	1.45	1.53
4	A	361	DCP	C5-C4	2.73	1.47	1.40
4	A	361	DCP	C4-N4	3.17	1.44	1.35
4	A	361	DCP	C4-N3	3.72	1.42	1.35
4	A	361	DCP	C6-C5	4.00	1.46	1.38
4	A	361	DCP	C6-N1	5.06	1.42	1.35

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	361	DCP	O5'-PA-O1A	-7.51	80.47	109.62
4	A	361	DCP	O2B-PB-O3A	-4.65	84.01	105.09
4	A	361	DCP	O3A-PA-O5'	-3.95	92.45	102.94
4	A	361	DCP	O2A-PA-O5'	-3.93	88.63	108.46
4	A	361	DCP	PB-O3B-PG	-3.63	120.50	132.67
4	A	361	DCP	O3G-PG-O3B	2.39	115.93	105.09
4	A	361	DCP	O4'-C1'-N1	2.46	111.97	107.72
4	A	361	DCP	C2-N3-C4	2.85	119.64	115.61
4	A	361	DCP	O2A-PA-O3A	2.99	118.65	105.09
4	A	361	DCP	O5'-C5'-C4'	3.16	120.76	109.12
4	A	361	DCP	O2B-PB-O3B	3.22	119.71	105.09

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	361	DCP	1	0

5.7 Other polymers

There are no such residues in this entry.









5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/360 (94%)	1.47	84 (24%)  	36, 59, 83, 91	0
2	P	13/13 (100%)	0.44	0  	54, 60, 72, 73	0
3	T	15/19 (78%)	0.20	0  	51, 65, 69, 77	0
All	All	369/392 (94%)	1.38	84 (22%)  	36, 60, 82, 91	0

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	116	ARG	6.9
1	A	335	VAL	6.7
1	A	37	PHE	5.9
1	A	334	GLY	5.7
1	A	287	VAL	5.3
1	A	248	ILE	5.2
1	A	333	ILE	4.9
1	A	5	PHE	4.8
1	A	107	ALA	4.8
1	A	209	GLU	4.7
1	A	48	TYR	4.7
1	A	274	TYR	4.7
1	A	286	VAL	4.7
1	A	337	PHE	4.4
1	A	4	LEU	4.2
1	A	108	TYR	4.2
1	A	216	MET	4.1
1	A	241	VAL	4.0
1	A	113	ASP	3.9
1	A	6	VAL	3.9
1	A	144	ILE	3.8
1	A	115	VAL	3.8
1	A	249	VAL	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	284	ILE	3.7
1	A	288	ALA	3.6
1	A	151	ALA	3.6
1	A	143	GLY	3.5
1	A	178	LEU	3.5
1	A	238	ARG	3.5
1	A	239	THR	3.4
1	A	149	VAL	3.4
1	A	142	VAL	3.3
1	A	289	VAL	3.3
1	A	187	GLY	3.3
1	A	210	PHE	3.2
1	A	197	LEU	3.2
1	A	169	GLU	3.2
1	A	109	LEU	3.1
1	A	306	ILE	3.1
1	A	3	VAL	3.1
1	A	150	PHE	3.1
1	A	246	GLY	3.0
1	A	154	ALA	3.0
1	A	192	GLU	3.0
1	A	265	LEU	2.9
1	A	117	ASP	2.9
1	A	278	LYS	2.9
1	A	8	PHE	2.9
1	A	212	LYS	2.9
1	A	189	ILE	2.8
1	A	253	ARG	2.8
1	A	97	GLU	2.8
1	A	188	ASN	2.8
1	A	233	TYR	2.8
1	A	224	TYR	2.7
1	A	38	GLU	2.7
1	A	193	LYS	2.7
1	A	196	LYS	2.7
1	A	11	PHE	2.7
1	A	176	ARG	2.6
1	A	153	ILE	2.6
1	A	252	LYS	2.6
1	A	315	SER	2.6
1	A	311	ALA	2.6
1	A	13	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	341	ILE	2.5
1	A	326	ASP	2.5
1	A	62	VAL	2.5
1	A	61	ILE	2.4
1	A	179	ASP	2.3
1	A	285	HIS	2.3
1	A	329	LYS	2.3
1	A	336	ARG	2.3
1	A	203	VAL	2.2
1	A	119	ARG	2.2
1	A	245	ILE	2.2
1	A	277	ASP	2.2
1	A	222	ALA	2.2
1	A	215	GLY	2.2
1	A	140	VAL	2.1
1	A	268	ALA	2.0
1	A	36	ARG	2.0
1	A	173	ARG	2.0
1	A	2	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
5	CA	A	363	1/1	0.93	0.17	-0.43	71,71,71,71	0
4	DCP	A	361	28/28	0.98	0.15	-0.58	34,43,52,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	CA	A	362	1/1	0.99	0.16	-0.58	38,38,38,38	0
5	CA	A	364	1/1	0.81	0.08	-	94,94,94,94	0

6.5 Other polymers [i](#)

There are no such residues in this entry.