



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:31 AM GMT

PDB ID : 2R2T  
Title : d(ATTAGTTAACTAAAT) complexed with MMLV RT catalytic fragment  
Authors : Goodwin, K.D.; Lewis, M.A.; Long, E.C.; Georgiadis, M.M.  
Deposited on : 2007-08-27  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	8	<div> <div></div> <div>100%</div> </div>
2	G	8	<div> <div></div> <div>13%</div> <div>88%</div> </div>
3	A	255	<div> <div></div> <div>7%</div> <div>75%</div> <div>24%</div> <div>.</div> </div>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 2542 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(\*DAP\*DTP\*DTP\*DTP\*DAP\*DGP\*DTP\*D T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	8	Total	C	N	O	P	0	0	0
			161	80	25	49	7			

- Molecule 2 is a DNA chain called DNA (5'-D(P\*DAP\*DAP\*DCP\*DTP\*DAP\*DAP\*DAP\*DT)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	8	Total	C	N	O	P	0	0	0
			164	79	32	45	8			

- Molecule 3 is a protein called Reverse transcriptase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	255	Total	C	N	O	S	0	0	0
			2041	1311	356	367	7			

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	171	Total	O	0	0
			171	171		
4	B	2	Total	O	0	0
			2	2		
4	G	3	Total	O	0	0
			3	3		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(\*DAP\*DTP\*DTP\*DTP\*DAP\*DGP\*DTP\*DT)-3')

Chain B: 


A1  
T2  
T3  
T4  
A5  
G6  
T7  
T8

- Molecule 2: DNA (5'-D(P\*DAP\*DAP\*DCP\*DTP\*DAP\*DAP\*DAP\*DT)-3')

Chain G: 

A9  
A10  
C11  
T12  
A13  
A14  
A15  
T16

- Molecule 3: Reverse transcriptase

Chain A: 

T24  
W25  
L26  
S27  
D28  
F29  
P30  
Q31  
I49  
I50  
P51  
T57  
P58  
I61  
M66  
S67  
Q68  
E69  
A70  
R71  
L72  
G73  
I74  
K75  
P76  
P100  
Y101  
K102  
K103  
P104  
G105  
T106  
M107  
D108  
E117  
H126  
P127  
M131  
P132  
Y133  
M134  
L135  
L136  
S137  
G138  
L139  
P140  
W145  
H161  
P162

T163  
R173  
D174  
P175  
E176  
M177  
G178  
I179  
S180  
G181  
Q182  
L183  
D206  
Q213  
I218  
V223  
A229  
A230  
E233  
L234  
D235  
C236  
Q237  
G248  
N249  
L250  
Q263  
V266  
K274  
E275  
G276  
Q277  
R278

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	54.64Å 145.80Å 46.88Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 43.72 – 2.00	Depositor EDS
% Data completeness (in resolution range)	92.3 (50.00-2.00) 92.4 (43.72-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.00Å)	Xtriage
Refinement program	CNS	Depositor
R, $R_{free}$	0.224 , 0.264 0.225 , 0.265	Depositor DCC
$R_{free}$ test set	1166 reflections (4.84%)	DCC
Wilson B-factor (Å <sup>2</sup> )	29.1	Xtriage
Anisotropy	0.276	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 49.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 25116 reflections (0.004%)	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2542	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.23	0/179	0.64	0/275
2	G	0.23	0/184	0.68	0/281
3	A	0.32	0/2097	0.59	0/2858
All	All	0.30	0/2460	0.60	0/3414

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	161	0	95	17	1
2	G	164	0	91	16	1
3	A	2041	0	2056	50	0
4	A	171	0	0	3	0
4	B	2	0	0	0	0
4	G	3	0	0	0	0
All	All	2542	0	2242	83	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:G:14:DA:H2''	2:G:15:DA:H5'	1.40	1.03
3:A:233:GLU:HG2	3:A:237:GLN:HE21	1.32	0.93
1:B:3:DT:H2''	1:B:4:DT:H5'	1.51	0.91
1:B:5:DA:H2''	1:B:6:DG:H5''	1.60	0.83
2:G:12:DT:H2''	2:G:13:DA:H5'	1.67	0.77
1:B:1:DA:H2''	1:B:2:DT:H5''	1.67	0.76
3:A:206:ASP:HB3	3:A:250:LEU:HD13	1.69	0.75
3:A:274:LYS:HG2	3:A:275:GLU:HG3	1.71	0.71
1:B:2:DT:H2''	1:B:3:DT:C5'	2.22	0.69
2:G:14:DA:C2'	2:G:15:DA:H5'	2.21	0.69
3:A:104:PRO:HG2	3:A:107:ASN:HB2	1.74	0.68
1:B:5:DA:C2'	1:B:6:DG:H5''	2.22	0.68
1:B:6:DG:H2''	1:B:7:DT:H5'	1.75	0.68
1:B:1:DA:H2''	1:B:2:DT:C5'	2.24	0.67
1:B:6:DG:H2''	1:B:7:DT:C5'	2.25	0.67
3:A:102:LYS:HD2	3:A:106:THR:HA	1.76	0.66
2:G:11:DC:H2''	2:G:12:DT:C5'	2.27	0.65
1:B:5:DA:H2''	1:B:6:DG:C5'	2.27	0.64
2:G:13:DA:H1'	2:G:14:DA:H5''	1.79	0.64
3:A:174:ASP:HB3	3:A:178:GLY:HA2	1.80	0.64
1:B:1:DA:C2'	1:B:2:DT:H5''	2.27	0.64
1:B:2:DT:H2''	1:B:3:DT:H5'	1.81	0.63
3:A:70:ALA:HB1	3:A:100:PRO:HB3	1.82	0.62
3:A:161:HIS:HD2	3:A:163:THR:H	1.49	0.61
3:A:173:ARG:HD2	4:A:434:HOH:O	2.02	0.59
3:A:68:GLN:O	3:A:72:LEU:HD13	2.02	0.59
3:A:103:LYS:HB2	3:A:104:PRO:HD3	1.84	0.59
1:B:3:DT:H2''	1:B:4:DT:C5'	2.29	0.59
3:A:161:HIS:CD2	3:A:163:THR:H	2.23	0.57
3:A:233:GLU:O	3:A:237:GLN:HG3	2.06	0.56
2:G:10:DA:H1'	2:G:11:DC:H5''	1.89	0.55
2:G:11:DC:H2''	2:G:12:DT:H5''	1.89	0.55
3:A:132:PRO:O	3:A:136:LEU:HD23	2.08	0.54
3:A:61:ILE:HD11	3:A:117:GLU:HG3	1.89	0.53
2:G:13:DA:C2'	2:G:14:DA:H5''	2.39	0.53
1:B:2:DT:H2''	1:B:3:DT:H5''	1.90	0.53
3:A:274:LYS:HE2	3:A:275:GLU:OE1	2.08	0.53
2:G:13:DA:H2''	2:G:14:DA:C5'	2.39	0.53
2:G:11:DC:H2''	2:G:12:DT:H5'	1.92	0.53
3:A:179:ILE:HG22	3:A:180:SER:N	2.24	0.52
3:A:233:GLU:HG2	3:A:237:GLN:NE2	2.13	0.52
3:A:173:ARG:HH11	3:A:173:ARG:HG2	1.74	0.52

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:26:LEU:O	3:A:26:LEU:HD23	2.08	0.52
2:G:13:DA:H2''	2:G:14:DA:H5''	1.93	0.51
3:A:24:THR:HB	3:A:27:SER:OG	2.11	0.51
2:G:13:DA:C1'	2:G:14:DA:H5''	2.41	0.50
1:B:2:DT:C2'	1:B:3:DT:H5''	2.42	0.50
2:G:9:DA:H2''	2:G:10:DA:C8	2.48	0.49
3:A:104:PRO:CG	3:A:107:ASN:HB2	2.42	0.48
3:A:145:TRP:CH2	3:A:233:GLU:HB2	2.48	0.48
3:A:179:ILE:HD13	3:A:183:LEU:HD21	1.95	0.48
3:A:57:THR:HG23	3:A:58:PRO:HD2	1.96	0.47
1:B:2:DT:H1'	1:B:3:DT:H5''	1.97	0.47
3:A:71:ARG:HD3	4:A:428:HOH:O	2.15	0.47
3:A:75:LYS:HB3	3:A:76:PRO:HD3	1.96	0.47
3:A:179:ILE:O	3:A:180:SER:C	2.52	0.46
2:G:11:DC:C2'	2:G:12:DT:H5''	2.46	0.46
3:A:138:GLY:O	3:A:140:PRO:HD3	2.14	0.46
3:A:277:GLN:OE1	3:A:277:GLN:HA	2.15	0.46
3:A:102:LYS:CD	3:A:106:THR:HA	2.43	0.46
3:A:105:GLY:O	3:A:106:THR:HB	2.17	0.45
3:A:31:GLN:HE22	3:A:249:ASN:HD22	1.64	0.45
1:B:4:DT:H2''	1:B:5:DA:C8	2.51	0.45
3:A:277:GLN:O	3:A:278:ARG:HB2	2.16	0.45
3:A:248:GLY:HA3	4:A:444:HOH:O	2.16	0.45
3:A:173:ARG:NH1	3:A:181:GLY:HA2	2.32	0.45
3:A:29:PHE:N	3:A:30:PRO:HD3	2.32	0.45
3:A:263:GLN:HB2	3:A:266:VAL:CG1	2.48	0.44
3:A:173:ARG:NH1	3:A:173:ARG:HG2	2.33	0.44
3:A:131:ASN:HD21	3:A:134:ASN:ND2	2.16	0.43
3:A:49:ILE:O	3:A:51:PRO:HD3	2.18	0.43
3:A:213:GLN:HA	3:A:213:GLN:NE2	2.34	0.43
3:A:218:ILE:HB	3:A:229:ALA:HB3	2.00	0.43
2:G:12:DT:C2'	2:G:13:DA:H5'	2.44	0.42
3:A:26:LEU:C	3:A:26:LEU:HD23	2.40	0.41
2:G:11:DC:H1'	2:G:12:DT:H5''	2.03	0.41
3:A:66:MET:HG2	3:A:70:ALA:HB3	2.02	0.41
1:B:6:DG:H2''	1:B:7:DT:H5''	2.00	0.41
3:A:126:HIS:HA	3:A:127:PRO:HD3	1.95	0.41
3:A:145:TRP:CZ2	3:A:233:GLU:HB2	2.56	0.41
3:A:74:ILE:HD11	3:A:100:PRO:HA	2.02	0.41
3:A:71:ARG:CZ	3:A:175:PRO:HG2	2.51	0.40
3:A:230:ALA:HB3	3:A:236:CYS:HB2	2.04	0.40



All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:8:DT:O3'	2:G:9:DA:P[2_765]	1.61	0.59

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	253/255 (99%)	241 (95%)	7 (3%)	5 (2%)	<b>9</b> <b>3</b>

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	106	THR
3	A	223	VAL
3	A	104	PRO
3	A	177	MET
3	A	181	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	224/224 (100%)	223 (100%)	1 (0%)	<b>93</b> <b>95</b>

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	177	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	31	GLN
3	A	84	GLN
3	A	134	ASN
3	A	144	GLN
3	A	161	HIS
3	A	213	GLN
3	A	214	HIS
3	A	237	GLN
3	A	238	GLN
3	A	245	GLN
3	A	249	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	B	8/8 (100%)	0.48	0	100	100	46, 61, 75, 79	0
2	G	8/8 (100%)	0.79	0	100	100	33, 74, 85, 85	0
3	A	255/255 (100%)	0.56	18 (7%)	19	20	18, 32, 67, 85	0
All	All	271/271 (100%)	0.56	18 (6%)	22	22	18, 33, 74, 85	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	A	106	THR	10.5
3	A	105	GLY	9.7
3	A	179	ILE	8.2
3	A	178	GLY	7.3
3	A	180	SER	7.0
3	A	177	MET	5.3
3	A	103	LYS	4.3
3	A	104	PRO	4.2
3	A	173	ARG	3.8
3	A	234	LEU	3.3
3	A	175	PRO	3.0
3	A	108	ASP	3.0
3	A	176	GLU	2.9
3	A	102	LYS	2.7
3	A	107	ASN	2.5
3	A	237	GLN	2.2
3	A	174	ASP	2.2
3	A	26	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.