



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:55 PM GMT

PDB ID : 1R8B  
Title : Crystal Structures of an Archaeal Class I CCA-Adding Enzyme and Its Nucleotide  
Authors : Xiong, Y.; Li, F.; Wang, J.; Weiner, A.M.; Steitz, T.A.  
Deposited on : 2003-10-23  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	437	<div> <div>9%</div> <div>87%</div> <div>12%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MG	A	602	-	-	-	X

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4027 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA nucleotidyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	0	0
			3630	2333	632	652	13			

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mn	0	0
			2	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Mg	0	0
			2	2		

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Na	0	0
			1	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: C<sub>10</sub>H<sub>16</sub>N<sub>5</sub>O<sub>13</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 31	C 10	N 5	O 13	P 3	0	0
6	A	1	Total 31	C 10	N 5	O 13	P 3	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	329	Total O 329 329	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: tRNA nucleotidyltransferase



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.40Å 79.37Å 78.14Å 90.00° 97.33° 90.00°	Depositor
Resolution (Å)	38.63 – 2.00 33.63 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.63-2.00) 99.7 (33.63-2.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.84 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.181 , 0.228 0.192 , 0.241	Depositor DCC
$R_{free}$ test set	1812 reflections (5.42%)	DCC
Wilson B-factor (Å <sup>2</sup> )	29.5	Xtriage
Anisotropy	0.300	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 52.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 35215 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4027	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.56% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, MG, MN, ATP, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.68	2/3713 (0.1%)	0.83	8/4987 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	314	MET	SD-CE	-9.09	1.26	1.77
1	A	314	MET	CG-SD	-5.09	1.68	1.81

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	373	ARG	NE-CZ-NH2	-11.82	114.39	120.30
1	A	373	ARG	NE-CZ-NH1	10.34	125.47	120.30
1	A	88	ASP	CB-CG-OD2	5.78	123.50	118.30
1	A	340	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	A	221	ASP	CB-CG-OD2	5.49	123.25	118.30
1	A	310	ARG	NE-CZ-NH1	5.22	122.91	120.30
1	A	437	ASP	CB-CG-OD2	5.13	122.92	118.30
1	A	128	ASP	CB-CG-OD2	5.12	122.91	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3630	0	3631	21	0
2	A	2	0	0	0	0
3	A	2	0	0	0	0
4	A	1	0	0	0	0
5	A	1	0	0	0	0
6	A	62	0	24	2	0
7	A	329	0	0	3	1
All	All	4027	0	3655	21	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:ARG:O	7:A:765:HOH:O	2.01	0.79
1:A:143:GLY:N	1:A:145:GLU:OE2	2.21	0.74
1:A:14:VAL:HG13	1:A:149:ARG:HB3	1.69	0.74
1:A:262:ILE:HD11	1:A:266:ARG:NH1	2.11	0.66
1:A:130:THR:OG1	6:A:501:ATP:H1'	2.02	0.58
1:A:14:VAL:CG1	1:A:149:ARG:HB3	2.34	0.56
1:A:321:LYS:NZ	1:A:424:PRO:O	2.40	0.54
1:A:258:HIS:ND1	1:A:258:HIS:C	2.61	0.53
1:A:76:ARG:O	1:A:80:LEU:HG	2.10	0.52
1:A:143:GLY:CA	1:A:145:GLU:OE2	2.58	0.50
1:A:258:HIS:ND1	1:A:258:HIS:O	2.44	0.50
1:A:165:TYR:CD2	1:A:295:PRO:HG3	2.49	0.48
1:A:165:TYR:HE2	1:A:298:GLU:OE1	1.98	0.47
1:A:165:TYR:CE2	1:A:298:GLU:OE1	2.70	0.45
1:A:31:LEU:CD1	1:A:86:VAL:CG1	2.96	0.44
1:A:101:HIS:C	1:A:101:HIS:CD2	2.90	0.43
1:A:165:TYR:CE1	1:A:168:ARG:HD3	2.53	0.43
1:A:64:LEU:HD11	1:A:111:VAL:CG1	2.50	0.42
1:A:95:ALA:HB1	6:A:501:ATP:HN62	1.87	0.40
1:A:340:ARG:HD2	7:A:807:HOH:O	2.22	0.40
1:A:356:LYS:NZ	7:A:915:HOH:O	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:657:HOH:O	7:A:683:HOH:O[3_545]	2.19	0.01

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	435/437 (100%)	422 (97%)	12 (3%)	1 (0%)	52 48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	387/387 (100%)	357 (92%)	30 (8%)	16 10

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LYS
1	A	4	GLU
1	A	29	GLU
1	A	33	ARG
1	A	51	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	69	GLU
1	A	71	SER
1	A	72	LYS
1	A	73	GLU
1	A	101	HIS
1	A	103	VAL
1	A	114	CYS
1	A	118	LYS
1	A	119	GLU
1	A	122	ASN
1	A	123	ILE
1	A	210	LYS
1	A	220	VAL
1	A	247	GLU
1	A	258	HIS
1	A	262	ILE
1	A	263	GLU
1	A	301	SER
1	A	317	ARG
1	A	325	GLU
1	A	340	ARG
1	A	356	LYS
1	A	373	ARG
1	A	398	HIS
1	A	419	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	HIS
1	A	334	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	ATP	A	501	3,2	24,33,33	1.13	2 (8%)	31,52,52	2.30	3 (9%)
6	ATP	A	502	2	24,33,33	1.17	2 (8%)	31,52,52	2.48	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	ATP	A	501	3,2	-	0/18/38/38	0/3/3/3
6	ATP	A	502	2	-	0/18/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	501	ATP	C2-N1	2.61	1.38	1.33
6	A	502	ATP	C2-N1	3.05	1.39	1.33
6	A	502	ATP	C2-N3	3.67	1.38	1.32
6	A	501	ATP	C2-N3	3.91	1.39	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	502	ATP	N3-C2-N1	-12.13	119.61	128.89
6	A	501	ATP	N3-C2-N1	-11.00	120.47	128.89
6	A	502	ATP	PA-O3A-PB	-3.36	123.28	132.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	502	ATP	C2'-C1'-N9	-3.05	109.64	114.29
6	A	501	ATP	O2B-PB-O3B	2.79	117.74	105.09
6	A	501	ATP	O4'-C1'-N9	3.41	115.24	108.10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	501	ATP	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	437/437 (100%)	0.16	39 (8%)	12 13	6, 19, 57, 72	0

All (39) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	118	LYS	11.3
1	A	123	ILE	7.8
1	A	125	SER	5.2
1	A	124	LYS	5.0
1	A	115	TYR	5.0
1	A	72	LYS	4.7
1	A	73	GLU	4.7
1	A	66	PHE	4.6
1	A	95	ALA	4.4
1	A	119	GLU	4.1
1	A	1	MET	3.9
1	A	22	ARG	3.6
1	A	70	PHE	3.6
1	A	126	ALA	3.6
1	A	98	PRO	3.3
1	A	121	LYS	3.2
1	A	120	PRO	3.2
1	A	75	LEU	3.1
1	A	258	HIS	3.0
1	A	96	GLU	3.0
1	A	97	HIS	2.9
1	A	67	PRO	2.9
1	A	80	LEU	2.8
1	A	117	LEU	2.7
1	A	261	GLU	2.6
1	A	19	GLU	2.6
1	A	25	ARG	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	69	GLU	2.6
1	A	77	GLU	2.6
1	A	122	ASN	2.5
1	A	17	ASP	2.4
1	A	2	LYS	2.4
1	A	76	ARG	2.3
1	A	26	GLU	2.3
1	A	105	LYS	2.2
1	A	260	LEU	2.2
1	A	74	GLU	2.1
1	A	114	CYS	2.1
1	A	128	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	MG	A	602	1/1	0.78	0.21	4.34	96,96,96,96	0
6	ATP	A	502	31/31	0.94	0.21	1.27	24,39,61,63	0
4	NA	A	605	1/1	0.97	0.12	0.47	44,44,44,44	0
6	ATP	A	501	31/31	0.94	0.14	-0.12	29,45,60,62	0
3	MG	A	604	1/1	0.58	0.18	-	62,62,62,62	0
2	MN	A	600	1/1	0.95	0.15	-	36,36,36,36	0
2	MN	A	601	1/1	0.98	0.22	-	27,27,27,27	0
5	CL	A	691	1/1	0.96	0.10	-	46,46,46,46	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.