



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 09:55 PM GMT

PDB ID : 1R8E  
Title : Crystal Structure of BmrR Bound to DNA at 2.4Å Resolution  
Authors : Newberry, K.J.; Brennan, R.G.  
Deposited on : 2003-10-23  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	23	<div> <div>4%</div> <div>74%</div> <div>26%</div> </div>
2	A	278	<div> <div>2%</div> <div>67%</div> <div>28%</div> <div>• •</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	P4P	A	400	-	-	-	X
4	IMD	A	302	-	-	-	X
4	IMD	A	303	-	-	-	X
4	IMD	A	304	-	-	-	X
4	IMD	A	310	-	-	-	X
5	GOL	A	401	-	X	-	X
5	GOL	A	402	-	X	-	X
5	GOL	A	403	-	X	-	X
5	GOL	B	203	-	X	-	-
5	GOL	B	205	-	X	-	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2963 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a DNA chain called 5'-D(\*GP\*AP\*CP\*CP\*CP\*TP\*CP\*CP\*CP\*CP\*TP\*TP\*AP\*GP\*GP\*GP\*GP\*AP\*GP\*GP\*GP\*TP\*C)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	23	Total	C	N	O	P	0	0	0
			468	222	87	137	22			

- Molecule 2 is a protein called multidrug-efflux transporter regulator.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	275	Total	C	N	O	S	0	0	0
			2254	1455	358	433	8			

- Molecule 3 is TETRAPHENYLPHOSPHONIUM (three-letter code: P4P) (formula: C<sub>24</sub>H<sub>20</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	P	0	0
			25	24	1		

- Molecule 4 is IMIDAZOLE (three-letter code: IMD) (formula:  $C_3H_5N_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		
4	A	1	Total	C	N	0	0
			5	3	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		



- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	118	Total	O	0	0
			118	118		
6	B	13	Total	O	0	0
			13	13		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(\*GP\*AP\*CP\*CP\*CP\*TP\*CP\*CP\*CP\*CP\*TP\*TP\*AP\*GP\*GP\*GP\*GP\*AP\*GP\*GP\*GP\*TP\*C)-3'



- Molecule 2: multidrug-efflux transporter regulator



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	107.28 Å   107.28 Å   145.70 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	86.39 – 2.40 86.39 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.9 (86.39-2.40) 99.0 (86.39-2.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.37 (at 2.40 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.227   ,   0.267 0.227   ,   0.267	Depositor DCC
$R_{free}$ test set	1694 reflections (5.04%)	DCC
Wilson B-factor (Å <sup>2</sup> )	55.1	Xtriage
Anisotropy	0.367	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 46.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 33613 reflections	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2963	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, IMD, P4P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.34	0/524	0.69	0/807
2	A	0.42	0/2303	0.64	2/3116 (0.1%)
All	All	0.41	0/2827	0.65	2/3923 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	188	ARG	NE-CZ-NH2	-6.35	117.12	120.30
2	A	212	ILE	N-CA-C	-5.43	96.33	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	468	0	259	4	0
2	A	2254	0	2232	75	0
3	A	25	0	20	0	0
4	A	55	0	55	2	0
5	A	18	0	12	0	0
5	B	12	0	8	0	0
6	A	118	0	0	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	13	0	0	0	0
All	All	2963	0	2586	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:156:LYS:HG2	2:A:167:ASN:HD21	1.20	1.02
2:A:196:THR:HG23	2:A:198:LYS:H	1.29	0.95
2:A:158:PHE:HB3	2:A:200:ILE:HD11	1.56	0.86
2:A:156:LYS:HG2	2:A:167:ASN:ND2	1.96	0.79
2:A:258:ILE:HD12	2:A:267:GLU:HG2	1.72	0.71
2:A:269:ARG:HH11	2:A:269:ARG:HB2	1.57	0.69
2:A:153:SER:O	2:A:157:LYS:HG2	1.94	0.67
2:A:143:GLY:O	2:A:147:VAL:HG22	1.95	0.67
2:A:223:ASN:HD21	2:A:269:ARG:NH1	1.92	0.66
2:A:160:GLU:HG2	2:A:166:THR:HG22	1.79	0.65
1:B:5:DG:H2''	1:B:6:DG:O5'	1.97	0.65
2:A:196:THR:HG23	2:A:198:LYS:N	2.09	0.64
2:A:127:LEU:HD22	2:A:127:LEU:H	1.63	0.63
2:A:240:ILE:HD11	2:A:274:ILE:HG21	1.80	0.62
2:A:135:ILE:HD13	2:A:136:GLN:N	2.16	0.60
2:A:135:ILE:HD11	2:A:207:MET:HB3	1.83	0.60
2:A:223:ASN:ND2	2:A:269:ARG:NH1	2.51	0.59
2:A:83:PHE:O	2:A:87:GLN:HG2	2.02	0.58
2:A:20:LYS:HG2	6:A:499:HOH:O	2.02	0.58
2:A:113:LYS:O	2:A:117:GLU:HG2	2.05	0.57
2:A:142:ILE:HG21	2:A:147:VAL:HG13	1.86	0.57
2:A:128:ASP:OD2	2:A:216:ARG:HG2	2.04	0.56
2:A:158:PHE:HE2	2:A:203:ILE:HA	1.69	0.56
2:A:168:ASN:HA	6:A:456:HOH:O	2.04	0.56
2:A:182:ILE:O	2:A:185:MET:HG2	2.07	0.55
2:A:269:ARG:NH1	2:A:269:ARG:HB2	2.22	0.55
2:A:121:LEU:HG	2:A:222:TYR:HA	1.90	0.53
2:A:274:ILE:HD13	2:A:275:ARG:N	2.24	0.53
2:A:154:LYS:HG2	2:A:155:LEU:N	2.24	0.53
2:A:154:LYS:H	2:A:154:LYS:HE3	1.74	0.53
2:A:75:LEU:HB2	2:A:79:GLU:HB2	1.90	0.52
2:A:239:TYR:O	2:A:243:ARG:HD3	2.09	0.52

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:94:LYS:O	2:A:98:LEU:HD23	2.10	0.52
2:A:188:ARG:HD2	2:A:250:ASP:OD2	2.09	0.52
2:A:68:GLU:O	2:A:71:LYS:HG3	2.11	0.51
2:A:218:ALA:HB3	2:A:274:ILE:HG22	1.94	0.50
2:A:154:LYS:HE2	4:A:306:IMD:HN1	1.75	0.50
2:A:147:VAL:HG21	2:A:187:TYR:HE1	1.77	0.50
2:A:158:PHE:CE2	2:A:203:ILE:HA	2.47	0.50
2:A:112:MET:O	2:A:116:MET:HG3	2.12	0.50
2:A:168:ASN:HD22	2:A:256:ILE:CD1	2.25	0.49
2:A:75:LEU:HD12	2:A:75:LEU:O	2.12	0.49
2:A:181:SER:O	2:A:184:GLU:HB2	2.14	0.48
2:A:32:LYS:O	2:A:49:GLN:NE2	2.47	0.47
2:A:101:LEU:O	2:A:105:ILE:HG12	2.13	0.47
2:A:162:ALA:O	2:A:196:THR:HG21	2.14	0.47
2:A:168:ASN:HD22	2:A:256:ILE:HD11	1.79	0.47
2:A:127:LEU:N	2:A:127:LEU:HD13	2.31	0.46
2:A:127:LEU:HD21	2:A:219:CYS:SG	2.55	0.46
2:A:154:LYS:HZ2	2:A:155:LEU:H	1.64	0.46
2:A:147:VAL:CG2	2:A:187:TYR:HE1	2.29	0.45
1:B:-7:DT:H2''	1:B:-6:DC:O5'	2.17	0.45
2:A:135:ILE:HD12	2:A:155:LEU:CD2	2.47	0.44
2:A:111:ARG:HG2	2:A:111:ARG:HH11	1.83	0.44
2:A:66:LEU:O	2:A:69:MET:HB2	2.17	0.44
2:A:206:ASP:OD2	2:A:207:MET:HG2	2.17	0.44
2:A:40:THR:O	2:A:41:SER:HB2	2.18	0.44
2:A:111:ARG:HD2	6:A:437:HOH:O	2.17	0.43
2:A:127:LEU:N	2:A:127:LEU:HD22	2.27	0.43
2:A:171:GLY:HA2	2:A:253:GLU:O	2.17	0.43
1:B:-8:DC:H2'	1:B:-7:DT:H71	2.01	0.43
2:A:204:THR:HG23	2:A:206:ASP:OD2	2.18	0.43
2:A:24:TYR:O	2:A:28:ILE:HG12	2.19	0.43
2:A:75:LEU:CD1	2:A:80:LEU:HB2	2.49	0.43
1:B:1:DT:H6	1:B:1:DT:H2'	1.71	0.43
2:A:258:ILE:HD12	2:A:267:GLU:CG	2.46	0.42
2:A:224:PHE:CG	2:A:225:SER:N	2.87	0.42
2:A:28:ILE:O	2:A:28:ILE:HG22	2.19	0.42
2:A:269:ARG:HH11	2:A:269:ARG:CB	2.31	0.42
2:A:135:ILE:C	2:A:135:ILE:HD13	2.40	0.42
2:A:96:ASP:N	2:A:96:ASP:OD2	2.53	0.42
2:A:135:ILE:HG23	2:A:155:LEU:HD22	2.02	0.42
2:A:247:VAL:HB	2:A:274:ILE:HD11	2.01	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:203:ILE:HD11	2:A:207:MET:HB2	2.02	0.41
2:A:239:TYR:O	2:A:243:ARG:HG2	2.20	0.41
2:A:168:ASN:HB3	4:A:309:IMD:N1	2.35	0.41
2:A:194:VAL:O	2:A:194:VAL:HG23	2.21	0.41
2:A:94:LYS:HD3	6:A:490:HOH:O	2.21	0.40
2:A:163:ASP:C	2:A:165:PHE:H	2.25	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	A	273/278 (98%)	270 (99%)	2 (1%)	1 (0%)	39 56

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	A	202	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	A	248/255 (97%)	230 (93%)	18 (7%)	17 27

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	51	ILE
2	A	55	LEU
2	A	71	LYS
2	A	107	LEU
2	A	118	TYR
2	A	127	LEU
2	A	135	ILE
2	A	147	VAL
2	A	154	LYS
2	A	192	THR
2	A	196	THR
2	A	204	THR
2	A	214	LYS
2	A	216	ARG
2	A	243	ARG
2	A	246	THR
2	A	269	ARG
2	A	274	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	73	GLN
2	A	136	GLN
2	A	167	ASN
2	A	168	ASN
2	A	223	ASN
2	A	265	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	IMD	A	302	-	3,5,5	0.68	0	4,5,5	0.56	0
4	IMD	A	303	-	3,5,5	0.65	0	4,5,5	0.57	0
4	IMD	A	304	-	3,5,5	0.69	0	4,5,5	0.58	0
4	IMD	A	305	-	3,5,5	0.64	0	4,5,5	0.56	0
4	IMD	A	306	-	3,5,5	0.61	0	4,5,5	0.56	0
4	IMD	A	307	-	3,5,5	0.66	0	4,5,5	0.57	0
4	IMD	A	308	-	3,5,5	0.62	0	4,5,5	0.55	0
4	IMD	A	309	-	3,5,5	0.59	0	4,5,5	0.60	0
4	IMD	A	310	-	3,5,5	0.57	0	4,5,5	0.59	0
4	IMD	A	311	-	3,5,5	0.64	0	4,5,5	0.59	0
4	IMD	A	312	-	3,5,5	0.64	0	4,5,5	0.58	0
3	P4P	A	400	-	28,28,28	1.91	11 (39%)	38,38,38	0.98	4 (10%)
5	GOL	A	401	-	5,5,5	4.78	5 (100%)	5,5,5	5.70	3 (60%)
5	GOL	A	402	-	5,5,5	4.72	5 (100%)	5,5,5	5.76	3 (60%)
5	GOL	A	403	-	5,5,5	4.75	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	B	203	-	5,5,5	4.78	5 (100%)	5,5,5	5.71	3 (60%)
5	GOL	B	205	-	5,5,5	4.80	5 (100%)	5,5,5	5.69	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	A	302	-	-	0/0/0/0	0/1/1/1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	IMD	A	303	-	-	0/0/0/0	0/1/1/1
4	IMD	A	304	-	-	0/0/0/0	0/1/1/1
4	IMD	A	305	-	-	0/0/0/0	0/1/1/1
4	IMD	A	306	-	-	0/0/0/0	0/1/1/1
4	IMD	A	307	-	-	0/0/0/0	0/1/1/1
4	IMD	A	308	-	-	0/0/0/0	0/1/1/1
4	IMD	A	309	-	-	0/0/0/0	0/1/1/1
4	IMD	A	310	-	-	0/0/0/0	0/1/1/1
4	IMD	A	311	-	-	0/0/0/0	0/1/1/1
4	IMD	A	312	-	-	0/0/0/0	0/1/1/1
3	P4P	A	400	-	-	0/24/24/24	0/4/4/4
5	GOL	A	401	-	-	0/4/4/4	0/0/0/0
5	GOL	A	402	-	-	0/4/4/4	0/0/0/0
5	GOL	A	403	-	-	0/4/4/4	0/0/0/0
5	GOL	B	203	-	-	0/4/4/4	0/0/0/0
5	GOL	B	205	-	-	0/4/4/4	0/0/0/0

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	205	GOL	C3-C2	-8.16	1.21	1.52
5	B	203	GOL	C3-C2	-8.13	1.21	1.52
5	A	403	GOL	C3-C2	-7.97	1.21	1.52
5	A	402	GOL	C3-C2	-7.79	1.22	1.52
5	A	401	GOL	C3-C2	-7.68	1.23	1.52
5	A	401	GOL	O2-C2	-3.19	1.33	1.43
5	A	403	GOL	C1-C2	-3.17	1.40	1.52
5	A	401	GOL	C1-C2	-3.16	1.40	1.52
5	B	203	GOL	C1-C2	-3.09	1.40	1.52
5	A	402	GOL	C1-C2	-3.01	1.40	1.52
5	B	205	GOL	C1-C2	-2.97	1.40	1.52
5	A	402	GOL	O2-C2	-2.95	1.34	1.43
5	B	205	GOL	O2-C2	-2.89	1.34	1.43
5	A	403	GOL	O2-C2	-2.82	1.35	1.43
5	B	203	GOL	O2-C2	-2.73	1.35	1.43
3	A	400	P4P	C4C-C5C	2.04	1.43	1.38
3	A	400	P4P	C3D-C4D	2.08	1.43	1.38
3	A	400	P4P	C6B-C1B	2.10	1.43	1.39
3	A	400	P4P	C5A-C6A	2.24	1.43	1.38
3	A	400	P4P	C6A-C1A	2.46	1.43	1.39
3	A	400	P4P	C2B-C1B	2.55	1.44	1.39
3	A	400	P4P	C3A-C2A	2.55	1.44	1.38

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	400	P4P	C6D-C1D	2.59	1.44	1.39
3	A	400	P4P	C2D-C1D	2.60	1.44	1.39
3	A	400	P4P	C2C-C1C	2.63	1.44	1.39
3	A	400	P4P	C6C-C1C	2.68	1.44	1.39
5	A	402	GOL	O3-C3	3.24	1.56	1.42
5	B	205	GOL	O3-C3	3.26	1.56	1.42
5	B	203	GOL	O3-C3	3.35	1.56	1.42
5	A	403	GOL	O3-C3	3.53	1.57	1.42
5	A	401	GOL	O3-C3	3.74	1.58	1.42
5	A	403	GOL	O1-C1	4.35	1.61	1.42
5	B	203	GOL	O1-C1	4.48	1.61	1.42
5	B	205	GOL	O1-C1	4.57	1.62	1.42
5	A	401	GOL	O1-C1	4.57	1.62	1.42
5	A	402	GOL	O1-C1	4.72	1.62	1.42

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	400	P4P	C1D-P-C1A	-2.38	104.86	109.44
3	A	400	P4P	C1B-P-C1C	-2.13	105.34	109.44
3	A	400	P4P	C1C-P-C1A	2.54	114.32	109.44
3	A	400	P4P	C1B-P-C1A	2.82	114.86	109.44
5	A	401	GOL	O1-C1-C2	2.87	124.11	110.18
5	A	402	GOL	O1-C1-C2	3.10	125.21	110.18
5	A	403	GOL	O1-C1-C2	3.16	125.49	110.18
5	B	203	GOL	O1-C1-C2	3.34	126.38	110.18
5	B	205	GOL	O1-C1-C2	3.36	126.48	110.18
5	A	401	GOL	O2-C2-C3	6.37	137.85	108.65
5	A	402	GOL	O2-C2-C3	6.37	137.88	108.65
5	B	203	GOL	O2-C2-C3	6.48	138.37	108.65
5	B	205	GOL	O2-C2-C3	6.49	138.43	108.65
5	A	403	GOL	O2-C2-C3	6.59	138.88	108.65
5	B	205	GOL	O3-C3-C2	10.40	160.60	110.18
5	A	403	GOL	O3-C3-C2	10.46	160.90	110.18
5	B	203	GOL	O3-C3-C2	10.46	160.91	110.18
5	A	401	GOL	O3-C3-C2	10.64	161.79	110.18
5	A	402	GOL	O3-C3-C2	10.74	162.27	110.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	306	IMD	1	0
4	A	309	IMD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	23/23 (100%)	0.28	1 (4%) 39 40	50, 82, 114, 146	0
2	A	275/278 (98%)	0.38	5 (1%) 71 71	32, 52, 82, 100	0
All	All	298/301 (99%)	0.37	6 (2%) 68 68	32, 53, 88, 146	0

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	167	ASN	3.6
2	A	201	SER	2.7
1	B	12	DC	2.5
2	A	159	ILE	2.2
2	A	200	ILE	2.2
2	A	166	THR	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	IMD	A	304	5/5	0.98	0.28	9.21	49,51,60,64	0
4	IMD	A	310	5/5	0.88	0.42	6.72	100,100,105,108	0
5	GOL	A	402	6/6	0.85	0.33	6.26	41,59,67,72	0
5	GOL	A	401	6/6	0.84	0.24	4.77	41,68,75,84	0
4	IMD	A	302	5/5	0.99	0.25	3.69	41,42,48,53	0
5	GOL	A	403	6/6	0.81	0.29	3.11	98,107,110,110	0
3	P4P	A	400	25/25	0.96	0.25	2.34	45,58,69,86	0
4	IMD	A	303	5/5	0.97	0.22	2.28	48,49,51,61	0
4	IMD	A	309	5/5	0.84	0.35	1.85	103,105,107,110	0
4	IMD	A	306	5/5	0.97	0.26	1.06	71,76,78,78	0
4	IMD	A	305	5/5	0.91	0.17	-0.01	88,90,94,97	0
5	GOL	B	203	6/6	0.81	0.18	-	91,101,106,111	0
4	IMD	A	311	5/5	0.83	0.31	-	99,100,104,105	0
4	IMD	A	307	5/5	0.89	0.39	-	105,106,112,114	0
5	GOL	B	205	6/6	0.74	0.17	-	82,96,99,99	0
4	IMD	A	312	5/5	0.71	0.23	-	125,125,127,127	0
4	IMD	A	308	5/5	0.81	0.22	-	113,113,114,115	0

## 6.5 Other polymers

There are no such residues in this entry.