



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:34 AM GMT

PDB ID : 2RAJ
Title : SO4 bound PX-BAR membrane remodeling unit of Sorting Nexin 9
Authors : Pylypenko, O.; Lundmark, R.; Rasmuson, E.; Carlsson, S.R.; Rak, A.
Deposited on : 2007-09-16
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3116 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sorting nexin-9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	382	3098	1977	523	578	20	0	0	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	13	13	13	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sorting nexin-9



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	61.57Å 147.48Å 115.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.93 – 2.45 19.93 – 2.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.93-2.45) 100.0 (19.93-2.40)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.21 (at 2.41Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.241 , 0.276 0.239 , 0.275	Depositor DCC
R_{free} test set	983 reflections (5.00%)	DCC
Wilson B-factor (Å ²)	71.3	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.1	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	1 of 20902 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3116	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/3168	0.59	1/4268 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	225	LEU	N-CA-C	-5.13	97.14	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3098	0	3085	84	0
2	A	5	0	0	0	0
3	A	13	0	0	0	0
All	All	3116	0	3085	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:455:GLN:NE2	1:A:455:GLN:H	1.74	0.85
1:A:224:GLN:HE22	1:A:381:SER:HB2	1.42	0.85
1:A:314:GLN:HG3	1:A:315:VAL:H	1.52	0.74
1:A:291:ASP:O	1:A:295:GLU:HG3	1.92	0.69
1:A:314:GLN:CG	1:A:315:VAL:H	2.05	0.67
1:A:324:ILE:HD11	1:A:325:LYS:NZ	2.09	0.66
1:A:224:GLN:HB3	1:A:540:ARG:NH2	2.10	0.66
1:A:403:GLY:HA2	1:A:507:ILE:HD13	1.78	0.66
1:A:314:GLN:HG3	1:A:315:VAL:N	2.11	0.65
1:A:455:GLN:NE2	1:A:455:GLN:N	2.45	0.65
1:A:321:GLU:HA	1:A:325:LYS:HG2	1.81	0.62
1:A:455:GLN:HE21	1:A:455:GLN:H	1.45	0.61
1:A:397:GLN:HG3	1:A:398:LYS:N	2.15	0.61
1:A:583:GLU:OE1	1:A:586:ARG:HD2	2.02	0.60
1:A:436:GLN:HE21	1:A:440:LYS:HE3	1.67	0.60
1:A:224:GLN:NE2	1:A:381:SER:HB2	2.16	0.59
1:A:217:GLU:HG2	1:A:516:LYS:NZ	2.18	0.58
1:A:214:PRO:O	1:A:217:GLU:HG3	2.04	0.58
1:A:281:ARG:HH11	1:A:281:ARG:HB3	1.68	0.58
1:A:466:GLU:HG3	1:A:591:ARG:CZ	2.34	0.58
1:A:281:ARG:NH1	1:A:281:ARG:HB3	2.19	0.57
1:A:276:PRO:C	1:A:278:ASN:H	2.09	0.57
1:A:259:LYS:HB3	1:A:268:SER:OG	2.06	0.55
1:A:267:LYS:HG2	1:A:267:LYS:O	2.07	0.55
1:A:503:PHE:HE2	1:A:552:MET:HE3	1.71	0.54
1:A:216:THR:HA	1:A:377:VAL:HG21	1.90	0.54
1:A:533:ASP:O	1:A:537:MET:HG3	2.08	0.53
1:A:361:GLU:HA	1:A:364:THR:HG22	1.88	0.53
1:A:262:LYS:O	1:A:263:MET:HB2	2.09	0.53
1:A:266:LEU:C	1:A:268:SER:H	2.13	0.52
1:A:485:ASP:HB3	1:A:573:GLN:HB2	1.90	0.52
1:A:318:ARG:HB3	1:A:323:PHE:CE2	2.45	0.52
1:A:392:LEU:HD11	1:A:514:ILE:HG23	1.92	0.51
1:A:431:LEU:HB3	1:A:432:PRO:HD3	1.93	0.51
1:A:224:GLN:HB3	1:A:540:ARG:HH22	1.73	0.51
1:A:583:GLU:HA	1:A:583:GLU:OE1	2.11	0.51
1:A:214:PRO:HG2	1:A:516:LYS:HZ1	1.75	0.51
1:A:262:LYS:HG2	1:A:262:LYS:O	2.11	0.51
1:A:217:GLU:HG2	1:A:516:LYS:HZ1	1.74	0.50
1:A:446:ALA:HB2	1:A:464:ILE:CG2	2.42	0.50
1:A:357:ARG:N	1:A:361:GLU:OE1	2.29	0.50
1:A:324:ILE:HD11	1:A:325:LYS:HZ2	1.75	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:254:VAL:HG22	1:A:274:LEU:CD2	2.42	0.49
1:A:241:PRO:HG2	1:A:549:GLN:HB3	1.93	0.49
1:A:255:ALA:HA	1:A:328:MET:SD	2.52	0.49
1:A:350:PHE:O	1:A:354:LEU:HD13	2.12	0.48
1:A:314:GLN:CG	1:A:315:VAL:N	2.72	0.48
1:A:535:GLN:HB3	1:A:539:LYS:HE3	1.94	0.48
1:A:583:GLU:O	1:A:587:GLN:HG3	2.14	0.48
1:A:535:GLN:O	1:A:539:LYS:HG3	2.14	0.48
1:A:361:GLU:HA	1:A:364:THR:CG2	2.43	0.47
1:A:308:PRO:O	1:A:334:TRP:NE1	2.42	0.47
1:A:318:ARG:H	1:A:318:ARG:HG2	1.47	0.46
1:A:569:LEU:O	1:A:573:GLN:HG2	2.16	0.46
1:A:385:PRO:HG3	1:A:543:ILE:HG12	1.96	0.45
1:A:257:PRO:HD3	1:A:328:MET:SD	2.56	0.45
1:A:526:THR:OG1	1:A:528:LYS:HE3	2.17	0.45
1:A:434:GLU:HA	1:A:437:LYS:HE2	1.99	0.45
1:A:430:PRO:O	1:A:434:GLU:HG3	2.17	0.45
1:A:559:ARG:HG2	1:A:559:ARG:HH11	1.82	0.45
1:A:437:LYS:HE3	1:A:437:LYS:HB2	1.84	0.44
1:A:219:TYR:CE2	1:A:540:ARG:HB3	2.53	0.43
1:A:583:GLU:HG3	1:A:587:GLN:HE21	1.83	0.43
1:A:353:PHE:HD2	1:A:354:LEU:HD12	1.83	0.43
1:A:271:GLU:HG2	1:A:284:ASN:HB3	2.00	0.43
1:A:322:GLU:OE1	1:A:322:GLU:HA	2.19	0.43
1:A:316:THR:O	1:A:316:THR:HG22	2.19	0.42
1:A:436:GLN:NE2	1:A:472:GLU:HG2	2.34	0.42
1:A:311:PRO:HG3	1:A:330:ARG:NE	2.35	0.42
1:A:314:GLN:HB3	1:A:323:PHE:CE1	2.54	0.42
1:A:281:ARG:CB	1:A:281:ARG:NH1	2.81	0.42
1:A:357:ARG:HG3	1:A:357:ARG:HH11	1.84	0.42
1:A:218:GLN:O	1:A:222:ALA:HB2	2.20	0.42
1:A:434:GLU:HA	1:A:437:LYS:CE	2.50	0.41
1:A:318:ARG:CB	1:A:323:PHE:CE2	3.02	0.41
1:A:446:ALA:HB2	1:A:464:ILE:HG22	2.02	0.41
1:A:466:GLU:HG3	1:A:591:ARG:NH1	2.34	0.41
1:A:218:GLN:O	1:A:222:ALA:N	2.50	0.41
1:A:403:GLY:HA2	1:A:507:ILE:CD1	2.50	0.41
1:A:258:ARG:HH21	1:A:284:ASN:ND2	2.19	0.40
1:A:591:ARG:HE	1:A:591:ARG:HB3	1.56	0.40
1:A:267:LYS:CG	1:A:267:LYS:O	2.68	0.40
1:A:241:PRO:HG2	1:A:549:GLN:CB	2.52	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:427:CYS:O	1:A:479:ALA:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	380/392 (97%)	345 (91%)	28 (7%)	7 (2%)	11 9

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	224	GLN
1	A	262	LYS
1	A	318	ARG
1	A	321	GLU
1	A	280	ASN
1	A	314	GLN
1	A	226	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	338/347 (97%)	324 (96%)	14 (4%)	37 52

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	224	GLN
1	A	230	GLU
1	A	238	ASP
1	A	239	TYR
1	A	291	ASP
1	A	319	PHE
1	A	326	MET
1	A	329	GLU
1	A	331	LEU
1	A	416	LEU
1	A	455	GLN
1	A	459	ASP
1	A	462	ASP
1	A	528	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	224	GLN
1	A	284	ASN
1	A	436	GLN
1	A	455	GLN
1	A	461	ASN
1	A	493	ASN
1	A	510	HIS
1	A	535	GLN
1	A	549	GLN
1	A	564	ASN
1	A	587	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates

There are no carbohydrates in this entry.

5.6 Ligand geometry

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	600	-	4,4,4	0.27	0	6,6,6	0.19	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	600	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/392 (97%)	0.81	48 (12%) 5 4	47, 74, 103, 114	0

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	263	MET	13.5
1	A	266	LEU	8.4
1	A	262	LYS	7.9
1	A	265	GLY	6.9
1	A	316	THR	6.6
1	A	225	LEU	6.5
1	A	314	GLN	6.3
1	A	315	VAL	6.2
1	A	264	TYR	5.6
1	A	317	GLY	5.3
1	A	318	ARG	4.3
1	A	527	SER	4.2
1	A	324	ILE	3.8
1	A	268	SER	3.8
1	A	313	LYS	3.7
1	A	357	ARG	3.5
1	A	261	SER	3.5
1	A	319	PHE	3.4
1	A	227	LYS	3.4
1	A	321	GLU	3.1
1	A	312	ASP	3.1
1	A	224	GLN	3.1
1	A	267	LYS	3.0
1	A	226	ALA	2.9
1	A	438	ILE	2.9
1	A	238	ASP	2.8
1	A	352	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	595	MET	2.7
1	A	325	LYS	2.7
1	A	428	THR	2.6
1	A	322	GLU	2.6
1	A	539	LYS	2.5
1	A	354	LEU	2.5
1	A	435	TYR	2.5
1	A	237	GLY	2.4
1	A	442	LEU	2.4
1	A	326	MET	2.4
1	A	320	GLU	2.4
1	A	215	GLY	2.3
1	A	272	TYR	2.3
1	A	353	PHE	2.2
1	A	223	LYS	2.2
1	A	218	GLN	2.2
1	A	427	CYS	2.1
1	A	363	LYS	2.1
1	A	445	LEU	2.1
1	A	329	GLU	2.1
1	A	413	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	A	600	5/5	0.94	0.15	-0.95	87,89,90,91	0

6.5 Other polymers [i](#)

There are no such residues in this entry.