



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:17 PM GMT

PDB ID : 4RDZ
Title : Crystal structure of VmoLac in P64 space group
Authors : Hiblot, J.; Bzdrenga, J.; Champion, C.; Gotthard, G.; Gonzalez, D.; Chabriere, E.; Elias, M.
Deposited on : 2014-09-20
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	<div> <div style="width: 85%;"></div> <div style="width: 12%;"></div> <div style="width: 3%;"></div> </div> <div>85% 12% .</div>
1	B	316	<div> <div style="width: 89%;"></div> <div style="width: 9%;"></div> <div style="width: 2%;"></div> </div> <div>89% 9% .</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	B	403	-	-	-	X
4	MYR	A	404	-	-	-	X
4	MYR	B	404	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6244 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Parathion hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	315	Total	C	N	O	S	0	16	0
			2591	1665	451	467	8			
1	B	316	Total	C	N	O	S	0	19	0
			2612	1685	448	470	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	EXPRESSION TAG	UNP F0QXN6
B	0	GLY	-	EXPRESSION TAG	UNP F0QXN6

- Molecule 2 is COBALT (II) ION (three-letter code: CO) (formula: Co).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	2	Total	Co	0	0
			2	2		
2	A	2	Total	Co	0	0
			2	2		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is MYRISTIC ACID (three-letter code: MYR) (formula: $C_{14}H_{28}O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			16	14	2		
4	B	1	Total	C	O	0	0
			16	14	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	486	Total 486	O 486	0	0
5	B	507	Total 507	O 507	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Parathion hydrolase



• Molecule 1: Parathion hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, α , β , γ	174.96 Å 174.96 Å 62.07 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.01 – 1.80 48.01 – 1.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.01-1.80) 100.0 (48.01-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.82 (at 1.79 Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.130 , 0.160 0.143 , 0.169	Depositor DCC
R_{free} test set	5032 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	20.1	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 46.2	EDS
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 100625 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6244	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MYR, CO, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.21	9/2676 (0.3%)	1.19	22/3620 (0.6%)
1	B	1.19	8/2706 (0.3%)	1.15	17/3660 (0.5%)
All	All	1.20	17/5382 (0.3%)	1.17	39/7280 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	45	GLU	CD-OE1	7.24	1.33	1.25
1	B	34	ARG	NE-CZ	7.20	1.42	1.33
1	B	315	ARG	C-O	7.16	1.36	1.23
1	A	5	SER	CB-OG	-7.16	1.32	1.42
1	B	5	SER	CB-OG	-6.95	1.33	1.42
1	A	261	THR	CB-CG2	-6.84	1.29	1.52
1	B	5	SER	CA-CB	6.69	1.62	1.52
1	A	46	GLU	CG-CD	6.48	1.61	1.51
1	A	35	TRP	CD2-CE2	6.31	1.49	1.41
1	A	45	GLU	CG-CD	6.07	1.61	1.51
1	A	297	GLU	CG-CD	5.88	1.60	1.51
1	B	174	VAL	C-O	5.70	1.34	1.23
1	B	60	TYR	CE1-CZ	-5.63	1.31	1.38
1	B	126	GLY	N-CA	-5.62	1.37	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	284	GLU	CD-OE2	-5.12	1.20	1.25
1	A	308	PRO	N-CA	5.11	1.55	1.47
1	B	0	GLY	N-CA	5.09	1.53	1.46

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	228	ASP	CB-CG-OD1	8.35	125.81	118.30
1	A	291	ARG	NE-CZ-NH1	-7.87	116.37	120.30
1	A	215	ARG	NE-CZ-NH1	-7.54	116.53	120.30
1	B	108	LYS	CD-CE-NZ	-7.39	94.70	111.70
1	B	289	ARG	NE-CZ-NH1	6.93	123.76	120.30
1	A	215	ARG	NE-CZ-NH2	6.81	123.70	120.30
1	B	136	PHE	CB-CG-CD2	6.75	125.52	120.80
1	A	49	ARG	NE-CZ-NH1	6.51	123.56	120.30
1	B	110	ARG	NE-CZ-NH2	-6.46	117.07	120.30
1	A	228	ASP	CB-CG-OD2	-6.45	112.49	118.30
1	B	34	ARG	NE-CZ-NH2	6.38	123.49	120.30
1	A	250	ASP	CB-CG-OD2	-6.37	112.57	118.30
1	A	271[A]	ARG	NE-CZ-NH2	-6.27	117.16	120.30
1	A	271[B]	ARG	NE-CZ-NH2	-6.27	117.16	120.30
1	A	3[A]	ARG	NE-CZ-NH1	-6.14	117.23	120.30
1	A	3[B]	ARG	NE-CZ-NH1	-6.14	117.23	120.30
1	B	34	ARG	NE-CZ-NH1	6.13	123.37	120.30
1	A	302[A]	ARG	NE-CZ-NH1	-6.00	117.30	120.30
1	A	302[B]	ARG	NE-CZ-NH1	-6.00	117.30	120.30
1	B	302[A]	ARG	NE-CZ-NH1	-5.92	117.34	120.30
1	B	302[B]	ARG	NE-CZ-NH1	-5.92	117.34	120.30
1	A	5	SER	CB-CA-C	-5.82	99.05	110.10
1	A	165	ASP	CB-CG-OD1	5.78	123.50	118.30
1	B	228	ASP	CB-CG-OD1	5.70	123.43	118.30
1	B	34	ARG	NH1-CZ-NH2	-5.69	113.14	119.40
1	B	291	ARG	NE-CZ-NH2	-5.60	117.50	120.30
1	A	46	GLU	OE1-CD-OE2	5.45	129.84	123.30
1	B	215	ARG	NE-CZ-NH2	5.42	123.01	120.30
1	A	113	ASP	CB-CG-OD2	-5.39	113.44	118.30
1	A	289	ARG	NE-CZ-NH2	-5.39	117.61	120.30
1	B	289	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	B	3	ARG	NE-CZ-NH2	5.09	122.85	120.30
1	A	261	THR	CA-CB-OG1	-5.06	98.38	109.00
1	A	113	ASP	CB-CG-OD1	5.06	122.85	118.30
1	B	184[A]	ARG	NE-CZ-NH2	-5.05	117.77	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	184[B]	ARG	NE-CZ-NH2	-5.05	117.77	120.30
1	A	236	ARG	NE-CZ-NH2	-5.04	117.78	120.30
1	B	250	ASP	CB-CG-OD2	-5.01	113.79	118.30
1	A	257	ASP	CB-CG-OD1	5.01	122.81	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	222	LEU	Peptide
1	B	222	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2591	0	2688	39	0
1	B	2612	0	2724	28	1
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	A	16	0	27	1	0
4	B	16	0	27	0	0
5	A	486	0	0	26	1
5	B	507	0	0	25	0
All	All	6244	0	5482	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:271[B]:ARG:CZ	5:A:722:HOH:O	1.94	1.14

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:LEU:HG	5:A:982:HOH:O	0.98	1.13
1:A:261:THR:HG22	5:A:944:HOH:O	1.45	1.12
1:B:302[B]:ARG:HD2	5:B:642:HOH:O	1.47	1.12
1:A:302[B]:ARG:HD2	5:A:604:HOH:O	1.51	1.05
1:A:271[B]:ARG:NE	5:A:722:HOH:O	1.88	1.04
1:A:261:THR:CG2	5:A:944:HOH:O	1.98	1.04
1:A:131:ASN:HB3	5:A:776:HOH:O	1.60	0.98
1:B:102:GLU:OE1	5:B:972:HOH:O	1.82	0.96
1:A:256:HIS:HE1	1:A:282:ILE:H	1.19	0.88
1:A:234[A]:ASP:OD1	5:A:804:HOH:O	1.95	0.84
1:B:302[B]:ARG:CD	5:B:642:HOH:O	2.15	0.81
1:A:302[B]:ARG:CG	5:A:604:HOH:O	2.27	0.81
1:A:302[B]:ARG:CD	5:A:604:HOH:O	2.15	0.78
1:A:314:GLY:C	5:A:886:HOH:O	2.20	0.78
1:B:234[B]:ASP:OD1	5:B:812:HOH:O	2.00	0.78
1:A:184[B]:ARG:NH1	5:A:974:HOH:O	2.16	0.77
1:A:148[A]:LYS:HD2	5:A:718:HOH:O	1.85	0.76
1:A:313:THR:HG21	5:A:945:HOH:O	1.91	0.71
1:B:302[B]:ARG:CG	5:B:642:HOH:O	2.37	0.70
1:A:302[B]:ARG:HG3	5:A:604:HOH:O	1.90	0.70
1:A:256:HIS:CE1	1:A:282:ILE:H	2.08	0.68
1:A:108[A]:LYS:NZ	5:A:943:HOH:O	2.22	0.66
1:B:250:ASP:O	1:B:310[B]:ARG:NH1	2.29	0.66
1:A:3[B]:ARG:CZ	5:A:951:HOH:O	2.45	0.63
1:B:302[B]:ARG:HG3	5:B:642:HOH:O	2.00	0.62
1:B:0:GLY:N	5:B:984:HOH:O	2.23	0.62
1:B:310[B]:ARG:NH2	5:B:898:HOH:O	2.32	0.61
1:A:0:GLY:N	5:A:947:HOH:O	2.33	0.61
1:A:148[B]:LYS:NZ	5:A:787:HOH:O	2.36	0.58
1:A:307:ASN:OD1	1:A:310[B]:ARG:NH2	2.35	0.56
1:B:245[A]:LYS:HG2	5:B:920:HOH:O	2.06	0.54
1:B:245[B]:LYS:CD	5:B:938:HOH:O	2.57	0.53
1:B:245[B]:LYS:NZ	5:B:938:HOH:O	2.41	0.53
1:B:85[B]:LYS:HG3	5:B:713:HOH:O	2.08	0.53
1:B:47[A]:LEU:HD21	5:B:852:HOH:O	2.09	0.52
1:B:271[B]:ARG:HG2	5:B:738:HOH:O	2.11	0.50
1:B:102:GLU:OE2	5:B:983:HOH:O	2.19	0.49
1:A:45:GLU:HG2	1:A:49:ARG:HH11	1.77	0.49
1:B:302[A]:ARG:NH1	5:B:627:HOH:O	2.46	0.48
1:B:228:ASP:OD2	5:B:823:HOH:O	2.20	0.47
1:A:11:GLU:OE2	1:A:133[B]:ARG:NH2	2.48	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:LEU:CG	5:A:982:HOH:O	1.87	0.46
1:B:245[A]:LYS:HD2	5:B:727:HOH:O	2.15	0.46
1:B:245[B]:LYS:HD3	5:B:938:HOH:O	2.15	0.46
1:A:121:HIS:HD2	5:A:750:HOH:O	1.99	0.45
1:A:103[B]:ILE:HD11	1:A:153:ALA:HB2	1.98	0.45
1:B:195:ARG:HD2	5:B:969:HOH:O	2.17	0.45
1:A:14:PRO:HG2	1:A:313:THR:HG22	1.98	0.44
1:A:34[A]:ARG:HD3	1:A:41:TYR:CG	2.52	0.44
1:A:131:ASN:CB	5:A:776:HOH:O	2.39	0.44
1:B:152:MET:HE2	5:B:809:HOH:O	2.18	0.43
1:A:184[A]:ARG:NH1	5:A:920:HOH:O	2.36	0.43
1:B:138:KCX:HA	1:B:169:ILE:O	2.18	0.43
1:B:121:HIS:HD2	5:B:687:HOH:O	2.02	0.43
1:A:229:ILE:HD13	4:A:404:MYR:H102	2.02	0.42
1:A:42:ASN:ND2	5:A:685:HOH:O	2.52	0.42
1:A:302[B]:ARG:CD	5:A:782:HOH:O	2.68	0.42
1:A:45:GLU:HG2	1:A:49:ARG:NH1	2.34	0.42
1:A:271[B]:ARG:NH1	5:A:735:HOH:O	2.53	0.41
1:A:148[A]:LYS:HA	1:A:148[A]:LYS:HD3	1.74	0.41
1:B:174:VAL:HA	5:B:571:HOH:O	2.21	0.41
1:A:46:GLU:HG2	1:A:261:THR:HG21	2.02	0.41
1:A:34[A]:ARG:HD3	1:A:41:TYR:CD1	2.56	0.41
1:B:1[A]:MET:CE	5:B:579:HOH:O	2.69	0.41
1:B:29[B]:ILE:HD11	5:B:965:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:298:GLU:OE1	5:A:945:HOH:O[6_645]	2.02	0.18

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	328/316 (104%)	325 (99%)	3 (1%)	0	100	100
1	B	332/316 (105%)	328 (99%)	4 (1%)	0	100	100
All	All	660/632 (104%)	653 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/270 (106%)	284 (100%)	1 (0%)	93	92
1	B	289/270 (107%)	288 (100%)	1 (0%)	94	94
All	All	574/540 (106%)	572 (100%)	2 (0%)	93	94

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	255	SER
1	B	255	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	ASN
1	A	90	ASN
1	A	212	GLN
1	A	251	GLN
1	A	256	HIS
1	B	36	ASN
1	B	90	ASN
1	B	121	HIS
1	B	212	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	251	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	KCX	A	138	1,2	7,11,12	0.53	0	7,12,14	1.03	0
1	KCX	B	138	1,2	7,11,12	0.64	0	7,12,14	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	138	1,2	-	0/6/10/12	0/0/0/0
1	KCX	B	138	1,2	-	0/6/10/12	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	138	KCX	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	403	-	5,5,5	1.20	0	5,5,5	1.59	2 (40%)
4	MYR	A	404	2	12,15,15	0.37	0	12,15,15	1.01	1 (8%)
3	GOL	B	403	-	5,5,5	0.66	0	5,5,5	1.32	1 (20%)
4	MYR	B	404	2	12,15,15	0.30	0	12,15,15	0.99	1 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	403	-	-	0/4/4/4	0/0/0/0
4	MYR	A	404	2	-	0/11/13/13	0/0/0/0
3	GOL	B	403	-	-	0/4/4/4	0/0/0/0
4	MYR	B	404	2	-	0/11/13/13	0/0/0/0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	404	MYR	C10-C9-C8	-2.33	102.51	114.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	403	GOL	C3-C2-C1	2.01	118.99	111.12
4	A	404	MYR	C11-C10-C9	2.14	125.60	114.53
3	B	403	GOL	O2-C2-C3	2.27	119.06	108.65
3	A	403	GOL	O1-C1-C2	2.49	122.28	110.18

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	404	MYR	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	314/316 (99%)	-0.67	0 100 100	12, 19, 32, 47	0
1	B	315/316 (99%)	-0.66	0 100 100	13, 19, 31, 51	0
All	All	629/632 (99%)	-0.66	0 100 100	12, 19, 32, 51	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	KCX	A	138	12/13	0.97	0.07	-	12,12,14,14	0
1	KCX	B	138	12/13	0.97	0.07	-	11,12,13,14	0

6.3 Carbohydrates

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	GOL	B	403	6/6	0.90	0.16	15.98	28,42,46,48	0
4	MYR	B	404	16/16	0.84	0.19	12.85	25,41,46,48	0
4	MYR	A	404	16/16	0.81	0.20	10.16	25,38,43,44	0
3	GOL	A	403	6/6	0.93	0.10	1.28	24,36,38,46	0
2	CO	B	402	1/1	1.00	0.04	-4.03	15,15,15,15	0
2	CO	A	402	1/1	1.00	0.04	-5.35	16,16,16,16	0
2	CO	B	401	1/1	1.00	0.04	-	14,14,14,14	0
2	CO	A	401	1/1	1.00	0.06	-	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.