



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 12:38 PM GMT

PDB ID : 3RJ9
Title : Structure of alcohol dehydrogenase from Drosophila lebanonesis T114V mutant complexed with NAD+
Authors : Morgunova, E.; Wuxiuer, Y.; Cols, N.; Popov, A.; Sylte, I.; Karshikoff, A.; Gonzales-Duarte, R.; Ladenstein, R.; Winberg, J.O.
Deposited on : 2011-04-15
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.98 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	8664 (2.00-1.96)
Clashscore	102246	9905 (2.00-1.96)
Ramachandran outliers	100387	9792 (2.00-1.96)
Sidechain outliers	100360	9791 (2.00-1.96)
RSRZ outliers	91569	8679 (2.00-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	254	9%	92% 7% •

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Alcohol dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	254	Total	C	N	O	S	0	0	0
			1963	1260	336	364	3			
1	B	254	Total	C	N	O	S	0	0	0
			1963	1260	336	364	3			
1	C	254	Total	C	N	O	S	0	0	0
			1963	1260	336	364	3			
1	D	254	Total	C	N	O	S	5	0	0
			1963	1260	336	364	3			
1	E	254	Total	C	N	O	S	0	0	0
			1963	1260	336	364	3			
1	F	254	Total	C	N	O	S	0	0	0
			1963	1260	336	364	3			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	VAL	THR	ENGINEERED MUTATION	UNP P10807
B	114	VAL	THR	ENGINEERED MUTATION	UNP P10807
C	114	VAL	THR	ENGINEERED MUTATION	UNP P10807
D	114	VAL	THR	ENGINEERED MUTATION	UNP P10807
E	114	VAL	THR	ENGINEERED MUTATION	UNP P10807
F	114	VAL	THR	ENGINEERED MUTATION	UNP P10807

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	44	21	7	14	2	0	0
2	B	1	44	21	7	14	2	0	0
2	C	1	44	21	7	14	2	0	0
2	D	1	44	21	7	14	2	0	0
2	E	1	44	21	7	14	2	0	0
2	F	1	44	21	7	14	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	182	182	182	0	0
3	B	187	187	187	0	0
3	C	220	220	220	0	0
3	D	200	200	200	0	0
3	E	120	120	120	0	0
3	F	109	109	109	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alcohol dehydrogenase




- Molecule 1: Alcohol dehydrogenase




- Molecule 1: Alcohol dehydrogenase




- Molecule 1: Alcohol dehydrogenase




- Molecule 1: Alcohol dehydrogenase





- Molecule 1: Alcohol dehydrogenase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.32Å 94.83Å 139.01Å 90.00° 98.33° 90.00°	Depositor
Resolution (Å)	49.62 – 1.98 49.62 – 1.98	Depositor EDS
% Data completeness (in resolution range)	93.0 (49.62-1.98) 93.0 (49.62-1.98)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) >$ ¹	3.53 (at 1.98Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.208 , 0.256 0.208 , 0.255	Depositor DCC
R_{free} test set	2066 reflections (2.03%)	DCC
Wilson B-factor (Å ²)	23.0	Xtriage
Anisotropy	0.931	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 53.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 104218 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13060	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.03% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/2002	0.61	0/2727
1	B	0.48	0/2002	0.60	0/2727
1	C	0.53	0/2002	0.62	0/2727
1	D	0.52	0/2002	0.63	0/2727
1	E	0.43	0/2002	0.56	0/2727
1	F	0.44	0/2002	0.57	0/2727
All	All	0.49	0/12012	0.60	0/16362

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	144	ALA	Peptide
1	C	144	ALA	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1963	0	2015	9	0
1	B	1963	0	2015	30	0
1	C	1963	0	2015	12	0
1	D	1963	0	2015	18	0
1	E	1963	0	2015	14	0
1	F	1963	0	2015	9	0
2	A	44	0	26	0	0
2	B	44	0	26	0	0
2	C	44	0	26	2	0
2	D	44	0	26	0	0
2	E	44	0	26	0	0
2	F	44	0	26	0	0
3	A	182	0	0	0	0
3	B	187	0	0	3	0
3	C	220	0	0	0	0
3	D	200	0	0	3	0
3	E	120	0	0	3	0
3	F	109	0	0	0	0
All	All	13060	0	12246	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:96:ASP:O	3:E:976:HOH:O	1.63	1.15
1:B:1:MET:HE3	1:B:6:LYS:HG3	1.52	0.92
1:B:38:ARG:HH12	1:B:102:ARG:NH2	1.76	0.83
1:D:100:ILE:HD11	3:D:320:HOH:O	1.80	0.80
1:B:38:ARG:NH1	1:B:102:ARG:NH2	2.30	0.80
1:D:1:MET:HE3	1:D:6:LYS:HG3	1.66	0.78
1:B:38:ARG:HH12	1:B:102:ARG:HH22	1.44	0.65
1:B:52:ASN:OD1	1:B:54:LYS:HE3	1.95	0.65
1:A:209:HIS:HB3	1:A:210:PRO:HD2	1.79	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:13:ALA:HB3	1:D:35:ILE:HG23	1.81	0.62
1:B:120:ASP:OD1	1:B:126:LYS:NZ	2.32	0.62
1:C:38:ARG:HH11	2:C:850:NAD:C8A	2.13	0.61
1:B:90:ASN:HB3	3:B:1015:HOH:O	2.01	0.60
1:E:1:MET:HB2	1:E:228:ASN:HB2	1.83	0.60
1:C:38:ARG:NH1	2:C:850:NAD:C8A	2.68	0.56
1:E:137:CYS:HB2	1:E:180:ASN:OD1	2.05	0.56
1:B:137:CYS:HB2	1:B:180:ASN:OD1	2.07	0.55
1:D:190:HIS:CE1	3:D:406:HOH:O	2.59	0.55
1:F:136:ILE:HD11	1:F:221:PHE:HE2	1.73	0.54
1:D:1:MET:HE2	1:D:87:ILE:HD11	1.88	0.53
1:C:110:GLY:O	1:C:114:VAL:HG23	2.09	0.53
1:E:163:ASN:HD21	1:E:233:ILE:HD11	1.75	0.52
1:B:1:MET:HE2	1:B:87:ILE:HD11	1.91	0.52
1:D:1:MET:CE	1:D:3:LEU:HD23	2.41	0.50
1:B:38:ARG:NH1	1:B:102:ARG:HH22	2.01	0.49
1:C:68:VAL:CG1	1:C:72:LYS:HE3	2.43	0.49
1:D:119:LEU:O	1:D:123:ASP:HB2	2.13	0.49
1:E:96:ASP:C	3:E:976:HOH:O	2.32	0.49
1:E:98:HIS:CD2	1:E:193:ASN:HB3	2.47	0.49
1:B:91:GLY:HA2	1:B:136:ILE:HD12	1.96	0.48
1:D:110:GLY:O	1:D:114:VAL:HG23	2.14	0.48
1:F:119:LEU:O	1:F:123:ASP:HB2	2.14	0.47
1:C:66:VAL:HB	1:C:67:PRO:HD2	1.95	0.47
1:B:60:HIS:HE1	1:B:77:LYS:NZ	2.13	0.47
1:B:119:LEU:O	1:B:123:ASP:HB2	2.14	0.47
1:B:163:ASN:ND2	3:B:459:HOH:O	2.48	0.47
1:E:209:HIS:HB3	1:E:210:PRO:CD	2.44	0.46
1:E:13:ALA:HB3	1:E:35:ILE:HG23	1.97	0.46
1:D:136:ILE:HD11	1:D:221:PHE:HE2	1.79	0.46
1:D:190:HIS:HE1	3:D:406:HOH:O	1.98	0.46
1:A:98:HIS:CD2	1:A:193:ASN:HB3	2.50	0.46
1:D:137:CYS:HB2	1:D:180:ASN:OD1	2.15	0.46
1:F:13:ALA:HB3	1:F:35:ILE:HG23	1.97	0.46
1:F:40:GLU:HG3	1:F:59:PHE:CE1	2.50	0.46
1:D:45:LEU:HG	1:D:49:LYS:HE3	1.98	0.46
1:A:137:CYS:HB2	1:A:180:ASN:OD1	2.15	0.46
1:F:137:CYS:HB2	1:F:180:ASN:OD1	2.16	0.45
1:D:1:MET:CE	1:D:87:ILE:HD11	2.46	0.45
1:D:1:MET:HE2	1:D:3:LEU:HD23	1.98	0.45
1:B:90:ASN:O	1:B:136:ILE:HG13	2.17	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:99:GLN:N	3:E:976:HOH:O	2.21	0.44
1:B:40:GLU:HG2	1:B:59:PHE:CE1	2.53	0.44
1:C:99:GLN:NE2	1:C:102:ARG:HD3	2.33	0.44
1:C:13:ALA:HB3	1:C:35:ILE:HG23	1.99	0.44
1:B:1:MET:CE	1:B:87:ILE:HD11	2.48	0.44
1:B:229:LYS:HB2	1:B:232:ALA:HB2	1.99	0.44
1:F:1:MET:HB3	1:F:226:GLU:HA	1.99	0.44
1:B:185:ARG:NH1	3:B:655:HOH:O	2.44	0.44
1:D:45:LEU:O	1:D:49:LYS:HG3	2.17	0.43
1:C:163:ASN:HD21	1:C:233:ILE:HG12	1.83	0.43
1:B:98:HIS:CD2	1:B:193:ASN:HB3	2.53	0.43
1:B:209:HIS:HB3	1:B:210:PRO:HD2	2.01	0.43
1:A:196:LEU:HD21	1:B:116:THR:HG23	2.01	0.43
1:B:91:GLY:O	1:B:92:ALA:C	2.56	0.43
1:A:72:LYS:HD2	1:A:121:PHE:CE1	2.53	0.43
1:E:209:HIS:HB3	1:E:210:PRO:HD2	2.01	0.43
1:D:98:HIS:CD2	1:D:193:ASN:HB3	2.54	0.42
1:A:179:ILE:HG13	1:A:221:PHE:CE1	2.55	0.42
1:F:93:GLY:HA2	1:F:107:ASN:OD1	2.20	0.42
1:B:209:HIS:HB3	1:B:210:PRO:CD	2.49	0.42
1:C:1:MET:HB3	1:C:228:ASN:HB2	2.02	0.42
1:C:196:LEU:HD21	1:D:116:THR:HG23	2.00	0.42
1:A:195:TRP:O	1:A:198:VAL:HG13	2.18	0.42
1:D:54:LYS:H	1:D:54:LYS:HE3	1.85	0.42
1:B:14:LEU:HD13	1:B:41:ASN:HB3	2.01	0.41
1:E:119:LEU:O	1:E:123:ASP:HB2	2.20	0.41
1:C:194:SER:OG	1:C:197:ASP:HA	2.21	0.41
1:E:251:ASP:HB3	1:E:253:HIS:CE1	2.55	0.41
1:A:185:ARG:NH1	1:A:211:THR:OG1	2.52	0.41
1:F:137:CYS:O	1:F:138:SER:HB2	2.20	0.41
1:B:137:CYS:O	1:B:138:SER:HB2	2.20	0.41
1:C:68:VAL:HG12	1:C:72:LYS:HE3	2.03	0.41
1:B:45:LEU:HG	1:B:49:LYS:HE2	2.03	0.41
1:E:96:ASP:OD1	1:E:193:ASN:HB2	2.21	0.41
1:B:91:GLY:CA	1:B:136:ILE:HD12	2.51	0.41
1:A:33:PHE:HB3	1:A:57:ILE:HG12	2.02	0.41
1:E:140:THR:HA	1:E:143:ASN:O	2.21	0.41
1:B:60:HIS:HE1	1:B:77:LYS:HZ1	1.69	0.41
1:F:163:ASN:HD21	1:F:233:ILE:HD11	1.87	0.40
1:B:185:ARG:NE	1:B:211:THR:OG1	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	252/254 (99%)	242 (96%)	10 (4%)	0	100 100
1	B	252/254 (99%)	242 (96%)	9 (4%)	1 (0%)	39 31
1	C	252/254 (99%)	243 (96%)	9 (4%)	0	100 100
1	D	252/254 (99%)	243 (96%)	9 (4%)	0	100 100
1	E	252/254 (99%)	242 (96%)	9 (4%)	1 (0%)	39 31
1	F	252/254 (99%)	240 (95%)	10 (4%)	2 (1%)	24 14
All	All	1512/1524 (99%)	1452 (96%)	56 (4%)	4 (0%)	46 39

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	96	ASP
1	B	138	SER
1	E	96	ASP
1	F	138	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	216/216 (100%)	210 (97%)	6 (3%)	51 48
1	B	216/216 (100%)	211 (98%)	5 (2%)	58 57
1	C	216/216 (100%)	212 (98%)	4 (2%)	65 65

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	216/216 (100%)	213 (99%)	3 (1%)	74	75
1	E	216/216 (100%)	213 (99%)	3 (1%)	74	75
1	F	216/216 (100%)	213 (99%)	3 (1%)	74	75
All	All	1296/1296 (100%)	1272 (98%)	24 (2%)	65	65

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	PHE
1	A	54	LYS
1	A	120	ASP
1	A	185	ARG
1	A	207	LEU
1	A	221	PHE
1	B	10	PHE
1	B	120	ASP
1	B	204	GLU
1	B	207	LEU
1	B	221	PHE
1	C	1	MET
1	C	10	PHE
1	C	39	VAL
1	C	221	PHE
1	D	10	PHE
1	D	54	LYS
1	D	221	PHE
1	E	10	PHE
1	E	119	LEU
1	E	221	PHE
1	F	10	PHE
1	F	64	VAL
1	F	221	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	99	GLN
1	A	163	ASN
1	A	228	ASN
1	B	56	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	60	HIS
1	B	81	GLN
1	B	163	ASN
1	B	209	HIS
1	B	219	GLN
1	B	228	ASN
1	C	56	ASN
1	C	99	GLN
1	C	163	ASN
1	C	228	ASN
1	D	81	GLN
1	D	99	GLN
1	D	190	HIS
1	D	228	ASN
1	E	56	ASN
1	E	99	GLN
1	E	163	ASN
1	E	219	GLN
1	E	228	ASN
1	F	163	ASN
1	F	228	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAD	A	850	-	38,48,48	0.80	1 (2%)	47,73,73	1.93	5 (10%)
2	NAD	B	850	-	38,48,48	0.78	1 (2%)	47,73,73	1.96	5 (10%)
2	NAD	C	850	-	38,48,48	0.73	1 (2%)	47,73,73	2.13	5 (10%)
2	NAD	D	850	-	38,48,48	0.82	1 (2%)	47,73,73	2.03	5 (10%)
2	NAD	E	850	-	38,48,48	0.78	1 (2%)	47,73,73	1.94	4 (8%)
2	NAD	F	850	-	38,48,48	0.72	0	47,73,73	2.03	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	850	-	-	0/22/62/62	0/5/5/5
2	NAD	B	850	-	-	0/22/62/62	0/5/5/5
2	NAD	C	850	-	-	0/22/62/62	0/5/5/5
2	NAD	D	850	-	-	0/22/62/62	0/5/5/5
2	NAD	E	850	-	-	0/22/62/62	0/5/5/5
2	NAD	F	850	-	-	0/22/62/62	0/5/5/5

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	850	NAD	O4D-C1D	2.05	1.43	1.41
2	A	850	NAD	O4D-C1D	2.33	1.44	1.41
2	C	850	NAD	O4D-C1D	2.35	1.44	1.41
2	E	850	NAD	O4D-C1D	2.40	1.44	1.41
2	D	850	NAD	O4D-C1D	2.50	1.44	1.41

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	850	NAD	N3A-C2A-N1A	-11.10	120.39	128.89
2	F	850	NAD	N3A-C2A-N1A	-11.08	120.41	128.89
2	C	850	NAD	N3A-C2A-N1A	-11.07	120.42	128.89
2	E	850	NAD	N3A-C2A-N1A	-10.94	120.52	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	850	NAD	N3A-C2A-N1A	-10.43	120.91	128.89
2	B	850	NAD	N3A-C2A-N1A	-10.11	121.16	128.89
2	B	850	NAD	C2B-C1B-N9A	-2.86	109.93	114.29
2	E	850	NAD	PN-O3-PA	-2.80	124.86	132.73
2	F	850	NAD	C1B-N9A-C4A	-2.64	122.96	126.94
2	C	850	NAD	C2B-C1B-N9A	-2.60	110.31	114.29
2	C	850	NAD	C1B-N9A-C4A	-2.59	123.03	126.94
2	A	850	NAD	O7N-C7N-N7N	-2.54	119.02	122.59
2	F	850	NAD	C4A-C5A-N7A	-2.21	107.45	109.48
2	D	850	NAD	C1B-N9A-C4A	-2.15	123.70	126.94
2	A	850	NAD	C2B-C1B-N9A	-2.00	111.23	114.29
2	E	850	NAD	C3N-C7N-N7N	2.16	120.18	117.82
2	D	850	NAD	C3N-C7N-N7N	2.35	120.39	117.82
2	F	850	NAD	C3N-C7N-N7N	2.76	120.84	117.82
2	B	850	NAD	C3N-C7N-N7N	2.85	120.93	117.82
2	B	850	NAD	O4D-C1D-N1N	3.03	111.46	108.13
2	E	850	NAD	O4B-C1B-N9A	3.16	114.70	108.10
2	F	850	NAD	O4B-C1B-N9A	3.16	114.71	108.10
2	D	850	NAD	O4B-C1B-N9A	3.22	114.83	108.10
2	F	850	NAD	O4D-C1D-N1N	3.24	111.69	108.13
2	A	850	NAD	O4D-C1D-N1N	3.51	111.98	108.13
2	C	850	NAD	O4B-C1B-N9A	3.57	115.56	108.10
2	A	850	NAD	C3N-C7N-N7N	3.75	121.92	117.82
2	D	850	NAD	O4D-C1D-N1N	3.87	112.39	108.13
2	B	850	NAD	O4B-C1B-N9A	4.55	117.61	108.10
2	C	850	NAD	O4D-C1D-N1N	5.65	114.34	108.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	850	NAD	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/254 (100%)	0.07	5 (1%) 68 71	13, 24, 38, 45	9 (3%)
1	B	254/254 (100%)	0.02	5 (1%) 68 71	14, 25, 36, 41	11 (4%)
1	C	254/254 (100%)	0.02	6 (2%) 62 66	13, 22, 34, 39	11 (4%)
1	D	254/254 (100%)	-0.04	6 (2%) 62 66	11, 22, 34, 39	10 (3%)
1	E	254/254 (100%)	0.61	18 (7%) 19 22	19, 36, 51, 54	11 (4%)
1	F	254/254 (100%)	0.59	24 (9%) 11 13	20, 35, 50, 54	11 (4%)
All	All	1524/1524 (100%)	0.21	64 (4%) 40 44	11, 27, 44, 54	63 (4%)

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	189	VAL	6.5
1	F	38	ARG	6.4
1	D	38	ARG	5.3
1	C	38	ARG	5.3
1	E	38	ARG	4.7
1	B	90	ASN	4.6
1	E	207	LEU	4.4
1	B	38	ARG	4.0
1	E	238	LEU	4.0
1	A	54	LYS	3.9
1	F	207	LEU	3.9
1	F	191	THR	3.7
1	F	1	MET	3.7
1	E	81	GLN	3.6
1	F	201	ARG	3.6
1	E	56	ASN	3.5
1	F	56	ASN	3.4
1	F	229	LYS	3.4
1	F	173	GLY	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	243	ALA	3.2
1	D	207	LEU	3.1
1	E	248	LYS	3.1
1	A	207	LEU	3.0
1	D	163	ASN	2.9
1	B	89	ILE	2.9
1	F	121	PHE	2.8
1	A	1	MET	2.8
1	F	205	LEU	2.8
1	E	244	ILE	2.8
1	C	102	ARG	2.8
1	E	183	ILE	2.8
1	C	165	LEU	2.7
1	F	165	LEU	2.6
1	F	206	LEU	2.6
1	F	254	ILE	2.6
1	A	191	THR	2.6
1	E	218	GLY	2.5
1	E	127	GLY	2.5
1	E	253	HIS	2.5
1	E	241	LEU	2.5
1	B	165	LEU	2.4
1	F	120	ASP	2.3
1	F	190	HIS	2.3
1	F	171	ILE	2.3
1	D	191	THR	2.2
1	B	81	GLN	2.2
1	E	200	PRO	2.2
1	F	72	LYS	2.2
1	C	90	ASN	2.1
1	A	185	ARG	2.1
1	C	66	VAL	2.1
1	D	190	HIS	2.1
1	E	79	PHE	2.1
1	F	188	LEU	2.1
1	E	219	GLN	2.1
1	E	25	LEU	2.0
1	C	191	THR	2.0
1	E	84	THR	2.0
1	F	172	THR	2.0
1	F	2	ASP	2.0
1	F	127	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	F	215	GLU	2.0
1	D	5	ASN	2.0
1	F	174	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	NAD	F	850	44/44	0.91	0.13	-0.22	39,41,42,43	0
2	NAD	B	850	44/44	0.96	0.09	-0.29	19,24,25,26	0
2	NAD	D	850	44/44	0.96	0.09	-0.31	20,22,24,26	0
2	NAD	A	850	44/44	0.97	0.09	-0.45	18,20,22,24	0
2	NAD	C	850	44/44	0.97	0.09	-0.46	20,22,25,26	0
2	NAD	E	850	44/44	0.96	0.09	-0.50	25,31,33,35	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.