



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:21 PM GMT

PDB ID : 4RNO  
Title : Crystal structure of human polymerase eta extending an abasic site-dA pair  
by inserting dCTP opposite template G  
Authors : Patra, A.; Egli, M.  
Deposited on : 2014-10-24  
Resolution : 2.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2676 (2.84-2.80)
Clashscore	102246	3124 (2.84-2.80)
Ramachandran outliers	100387	3072 (2.84-2.80)
Sidechain outliers	100360	3074 (2.84-2.80)
RSRZ outliers	91569	2690 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	435	<div> <div>4%</div> <div>80%</div> <div>16%</div> <div>..</div> </div>
2	T	12	<div> <div>58%</div> <div>42%</div> </div>
3	P	8	<div> <div>88%</div> <div>13%</div> </div>

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 3801 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase eta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	425	Total	C	N	O	S	0	0	0
			3329	2087	598	621	23			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP Q9Y253
A	-1	PRO	-	EXPRESSION TAG	UNP Q9Y253
A	0	HIS	-	EXPRESSION TAG	UNP Q9Y253


- Molecule 2 is a DNA chain called Nucleic acids Template: CATG(3DR)TGACGCT.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	T	12	Total	C	N	O	P	0	0	1
			218	103	37	67	11			

- Molecule 3 is a DNA chain called Nucleic acids Primar: AGCGTCAA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	P	8	Total	C	N	O	P	0	0	0
			162	78	33	44	7			

- Molecule 4 is 2'-DEOXYCYTIDINE-5'-TRIPHOSPHATE (three-letter code: DCP) (formula: C<sub>9</sub>H<sub>16</sub>N<sub>3</sub>O<sub>13</sub>P<sub>3</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			28	9	3	13	3		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	49	Total	O	0	0
			49	49		
7	T	6	Total	O	0	1
			7	7		
7	P	1	Total	O	0	0
			1	1		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase eta



- Molecule 2: Nucleic acids Template: CATG(3DR)TGACGCT



- Molecule 3: Nucleic acids Primar: AGCGTCAA



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.79 Å 98.79 Å 82.01 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	42.78 – 2.82 42.78 – 2.82	Depositor EDS
% Data completeness (in resolution range)	89.8 (42.78-2.82) 88.8 (42.78-2.82)	Depositor EDS
$R_{merge}$	0.21	Depositor
$R_{sym}$	0.21	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.03 (at 2.81 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1839)	Depositor
R, $R_{free}$	0.241 , 0.282 0.242 , 0.288	Depositor DCC
$R_{free}$ test set	497 reflections (5.34%)	DCC
Wilson B-factor (Å <sup>2</sup> )	69.6	Xtriage
Anisotropy	0.081	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 25.0	EDS
Estimated twinning fraction	0.069 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 9779 reflections (0.010%)	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3801	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, 3DR, DCP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.21	0/3384	0.41	0/4563
2	T	0.52	0/230	0.90	0/352
3	P	0.54	0/182	0.83	0/279
All	All	0.27	0/3796	0.49	0/5194

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3329	0	3372	41	0
2	T	218	0	122	5	0
3	P	162	0	91	1	0
4	A	28	0	12	0	0
5	A	1	0	0	0	0
6	A	6	0	8	0	0
7	A	49	0	0	0	0
7	P	1	0	0	0	0
7	T	7	0	0	0	0
All	All	3801	0	3605	44	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:T:5:3DR:O4'	2:T:5:3DR:C4'	1.67	1.20
1:A:67:ASP:HA	1:A:70:LYS:HE2	1.72	0.71
1:A:366:LEU:HB3	1:A:385:CYS:HB3	1.82	0.61
1:A:166:MET:O	1:A:168:LYS:N	2.34	0.60
1:A:394:LYS:NZ	1:A:398:ASP:OD2	2.37	0.58
1:A:120:ASP:OD1	1:A:122:THR:OG1	2.24	0.56
1:A:371:ARG:HG2	1:A:420:THR:HB	1.86	0.56
2:T:4:DG:H1'	2:T:5:3DR:H1'2	1.89	0.54
1:A:427:THR:OG1	1:A:428:LYS:N	2.39	0.54
1:A:197:VAL:HG21	1:A:216:ILE:HG12	1.88	0.53
1:A:62:SER:HB2	2:T:3:DT:H72	1.91	0.53
1:A:24:ARG:O	1:A:30:ARG:NH2	2.43	0.52
1:A:375:ASP:OD2	1:A:380:SER:OG	2.23	0.51
1:A:86:LYS:NZ	1:A:87:ALA:O	2.41	0.50
1:A:204:ILE:O	1:A:208:THR:OG1	2.25	0.49
1:A:142:LEU:HD23	1:A:198:GLU:HG3	1.95	0.49
1:A:324:ASN:N	2:T:5:3DR:OP2	2.39	0.48
1:A:5:GLN:HB3	1:A:186:PRO:HB3	1.97	0.47
1:A:300:ALA:HB1	1:A:305:ILE:HB	1.98	0.46
1:A:27:PRO:HA	1:A:30:ARG:NE	2.29	0.46
1:A:419:LEU:HD13	1:A:422:LEU:HD21	1.97	0.46
1:A:400:PHE:O	1:A:404:LYS:N	2.48	0.46
1:A:293:LYS:HE2	1:A:293:LYS:HB2	1.85	0.46
1:A:86:LYS:HB3	1:A:323:LYS:HD3	1.97	0.45
1:A:164:GLU:OE2	1:A:167:ARG:NE	2.49	0.45
1:A:224:LYS:NZ	3:P:8:DA:OP1	2.34	0.45
2:T:9:DC:H2''	2:T:10:DG:C8	2.51	0.45
1:A:6:ASP:OD2	1:A:6:ASP:N	2.50	0.45
1:A:333:THR:HA	1:A:414:GLU:HA	1.99	0.45
1:A:7:ARG:NH1	1:A:306:GLU:O	2.49	0.45
1:A:355:ASP:OD1	1:A:359:ASN:ND2	2.49	0.44
1:A:366:LEU:O	1:A:385:CYS:N	2.50	0.44
1:A:185:SER:HA	1:A:186:PRO:HD3	1.87	0.44
1:A:7:ARG:NH2	1:A:309:PRO:HA	2.33	0.44
1:A:226:ALA:HA	1:A:229:LEU:HD12	2.01	0.43
1:A:294:ASN:N	1:A:294:ASN:OD1	2.52	0.43
1:A:48:ILE:O	1:A:61:ARG:HD3	2.20	0.42

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:14:MET:HB2	1:A:17:PHE:HB2	2.02	0.41
1:A:247:PHE:HA	1:A:250:MET:HB3	2.02	0.41
1:A:198:GLU:HG2	1:A:235:GLN:NE2	2.36	0.41
1:A:322:SER:HB3	1:A:423:PHE:HD2	1.85	0.41
1:A:293:LYS:HG3	1:A:293:LYS:H	1.64	0.40
1:A:315:LEU:HA	1:A:316:PRO:HD3	1.89	0.40
1:A:290:PHE:HB2	1:A:294:ASN:HB2	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	419/435 (96%)	386 (92%)	31 (7%)	2 (0%)	34 68

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	166	MET
1	A	167	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/372 (98%)	351 (96%)	13 (4%)	42 75

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	ASP
1	A	74	ASP
1	A	79	GLN
1	A	122	THR
1	A	240	HIS
1	A	242	SER
1	A	294	ASN
1	A	306	GLU
1	A	321	CYS
1	A	334	ARG
1	A	378	LEU
1	A	411	ILE
1	A	413	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	3DR	T	5	2	7,11,12	5.87	5 (71%)	8,14,17	1.02	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3DR	T	5	2	-	0/3/15/16	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	T	5	3DR	C3'-C4'	-6.81	1.33	1.53
2	T	5	3DR	O4'-C1'	-3.78	1.31	1.42
2	T	5	3DR	C2'-C3'	2.61	1.56	1.52
2	T	5	3DR	C2'-C1'	3.15	1.60	1.51
2	T	5	3DR	O4'-C4'	12.69	1.67	1.44

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	T	5	3DR	C1'-O4'-C4'	-2.04	105.50	108.58

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	T	5	3DR	3	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
4	DCP	A	501	5	21,29,29	3.42	10 (47%)	33,45,45	1.40	6 (18%)
6	GOL	A	503	-	5,5,5	0.36	0	5,5,5	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	DCP	A	501	5	-	0/18/34/34	0/2/2/2
6	GOL	A	503	-	-	0/4/4/4	0/0/0/0

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	DCP	C2'-C1'	-7.42	1.31	1.52
4	A	501	DCP	C3'-C4'	-5.97	1.36	1.53
4	A	501	DCP	C5-C4	2.56	1.46	1.40
4	A	501	DCP	C4-N4	2.91	1.43	1.35
4	A	501	DCP	C2'-C3'	2.98	1.61	1.52
4	A	501	DCP	O4'-C4'	3.34	1.52	1.45
4	A	501	DCP	C6-C5	3.88	1.46	1.38
4	A	501	DCP	C4-N3	4.14	1.43	1.35
4	A	501	DCP	C6-N1	5.14	1.43	1.35
4	A	501	DCP	O4'-C1'	7.63	1.60	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	DCP	PB-O3A-PA	-2.91	124.56	132.73
4	A	501	DCP	PB-O3B-PG	-2.48	124.34	132.67
4	A	501	DCP	C3'-C2'-C1'	2.38	108.13	102.40
4	A	501	DCP	C2'-C3'-C4'	2.42	107.79	102.77
4	A	501	DCP	O3A-PA-O5'	2.68	110.05	102.94
4	A	501	DCP	C2-N3-C4	3.18	120.10	115.61

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	425/435 (97%)	0.36	17 (4%) 42 30	45, 55, 79, 96	0
2	T	11/12 (91%)	0.03	0 100 100	52, 55, 101, 104	0
3	P	8/8 (100%)	-0.41	0 100 100	50, 53, 59, 59	0
All	All	444/455 (97%)	0.34	17 (3%) 44 32	45, 55, 80, 104	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	416	SER	4.6
1	A	408	THR	4.2
1	A	369	SER	4.0
1	A	415	TRP	3.6
1	A	419	LEU	3.6
1	A	370	ILE	3.3
1	A	407	ASN	3.0
1	A	368	VAL	3.0
1	A	331	LEU	3.0
1	A	406	CYS	2.8
1	A	378	LEU	2.8
1	A	377	ARG	2.6
1	A	380	SER	2.6
1	A	379	SER	2.3
1	A	258	LEU	2.1
1	A	22	GLU	2.1
1	A	183	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	3DR	T	5	11/12	0.94	0.16	-	59,61,62,63	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	DCP	A	501	28/28	0.94	0.20	-0.20	47,49,57,59	0
5	CA	A	502	1/1	0.93	0.12	-2.28	47,47,47,47	0
6	GOL	A	503	6/6	0.71	0.15	-	58,64,67,68	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.