



wwPDB X-ray Structure Validation Summary Report

Feb 1, 2016 – 08:23 PM GMT

PDB ID : 4RR2
Title : Crystal structure of human primase
Authors : Baranovskiy, A.G.; Gu, J.; Suwa, Y.; Babayeva, N.D.; Tahirov, T.H.
Deposited on : 2014-11-05
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	420	 % 46% 43% 7%
1	C	420	 % 44% 43% 6% 7%
2	B	509	 6% 14% 23% 5% 58%
2	D	509	 4% 40% 37% 6% 16%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA primase small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	389	Total 3267	C 2101	N 566	O 585	S 15	0	0	0
1	C	389	Total 3263	C 2099	N 566	O 583	S 15	0	0	0


- Molecule 2 is a protein called DNA primase large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	213	Total 1755	C 1133	N 299	O 321	S 2	0	0	0
2	D	429	Total 3513	C 2249	N 608	O 643	S 13	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0
3	C	1	Total 1	Zn 1	0	0

- Molecule 4 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	1	Total	Fe S	0	0
			8	4 4		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	23	Total	O	0	0
			23	23		
5	B	4	Total	O	0	0
			4	4		
5	C	40	Total	O	0	0
			40	40		
5	D	21	Total	O	0	0
			21	21		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA primase small subunit



- Molecule 1: DNA primase small subunit



GLY	T360	K212
LYS	D961	D213
ASP	Y362	I214
ILE	T363	I218
LYS	F364	L219
LYS	F365	W220
GLU	S366	E221
PRO	K369	F222
ILE	I370	K225
GLN	M288	L226
PRO	S373	S227
GLU	M374	K228
THR	P375	A229
PRO	P376	L230
GLN	S377	A231
PRO	Q378	L232
LYS	H382	T233
PRO	P385	A234
SER	F386	R235
VAL	R387	S236
GLN	E392	L237
LYS	L394	P238
THR	L311	A239
LYS	F312	V240
LYS	I316	Q241
ASP	E320	S242
ALA	L320	D243
SER	E321	R245
SER	Q322	L246
ALA	A323	Q247
LEU	L324	P248
LEU	K325	L249
SER	K326	L250
LEU	F327	W251
ASN	H328	H252
SER	Y329	L253
SER	Q421	S254
LEU	V422	H255
GLU	A423	S256
MET	C424	Y257
ASP	Q425	T258
MET	E429	G259
GLU	M430	D261
GLY	I431	Y262
LEU	H432	S263
GLU	C437	T264
ASP	G438	Q265
TYR	F439	G266
PHE	S440	M267
SER	L441	V268
GLU	N442	G269
ASP	H443	K270
SER	P444	I271
GLU	P445	L272
GLU	N445	L273
ASP	Q446	D274
SER	F447	
	N456	
	GLY	

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	86.19Å 88.90Å 94.68Å 93.82° 96.57° 111.72°	Depositor
Resolution (Å)	48.48 – 2.65 48.47 – 2.60	Depositor EDS
% Data completeness (in resolution range)	88.5 (48.48-2.65) 84.6 (48.47-2.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.89 (at 2.61Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.229 , 0.271 0.216 , 0.251	Depositor DCC
R_{free} test set	3493 reflections (5.56%)	DCC
Wilson B-factor (Å ²)	71.2	Xtriage
Anisotropy	0.392	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 51.9	EDS
Estimated twinning fraction	0.006 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 72404 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11896	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/3349	0.70	0/4514
1	C	0.49	0/3346	0.74	0/4512
2	B	0.39	0/1787	0.63	0/2402
2	D	0.44	0/3595	0.67	1/4838 (0.0%)
All	All	0.45	0/12077	0.69	1/16266 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	252	HIS	N-CA-C	5.33	125.39	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3267	0	3249	206	0
1	C	3263	0	3243	198	0
2	B	1755	0	1801	184	0
2	D	3513	0	3494	232	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	8	0	0	0	0
5	A	23	0	0	11	0
5	B	4	0	0	0	0
5	C	40	0	0	11	0
5	D	21	0	0	3	0
All	All	11896	0	11787	815	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

The worst 5 of 815 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:82:LYS:HE2	2:D:98:ARG:HD3	1.36	1.07
2:D:117:ARG:HG2	2:D:230:LEU:HD13	1.38	1.05
2:B:68:GLU:HA	2:B:71:GLN:HB3	1.42	1.02
2:B:82:LYS:HG2	2:B:98:ARG:HD3	1.42	1.01
1:C:50:LYS:HD2	1:C:50:LYS:H	1.25	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	383/420 (91%)	331 (86%)	45 (12%)	7 (2%)	11	23
1	C	383/420 (91%)	327 (85%)	36 (9%)	20 (5%)	2	4
2	B	209/509 (41%)	156 (75%)	37 (18%)	16 (8%)	1	1
2	D	425/509 (84%)	346 (81%)	58 (14%)	21 (5%)	3	4
All	All	1400/1858 (75%)	1160 (83%)	176 (13%)	64 (5%)	3	5

5 of 64 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	37	LYS
1	A	50	LYS
1	A	95	LYS
1	A	196	GLN
1	A	380	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/393 (93%)	347 (95%)	17 (5%)	32 60
1	C	363/393 (92%)	341 (94%)	22 (6%)	23 46
2	B	194/459 (42%)	171 (88%)	23 (12%)	6 13
2	D	389/459 (85%)	363 (93%)	26 (7%)	20 41
All	All	1310/1704 (77%)	1222 (93%)	88 (7%)	20 41

5 of 88 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	247	GLN
1	C	119	ARG
2	D	288	MET
1	C	2	GLU
1	C	55	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	86	HIS
1	C	257	GLN
2	D	329	GLN
1	C	89	ASN
1	C	102	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SF4	D	1000	2	0,12,12	0.00	-	0,24,24	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	SF4	D	1000	2	-	0/0/48/48	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	389/420 (92%)	-0.01	5 (1%) 79 79	13, 42, 78, 100	0
1	C	389/420 (92%)	-0.03	5 (1%) 79 79	14, 37, 75, 95	0
2	B	213/509 (41%)	0.64	29 (13%) 4 3	23, 77, 100, 109	0
2	D	429/509 (84%)	0.14	20 (4%) 35 33	15, 61, 95, 107	0
All	All	1420/1858 (76%)	0.13	59 (4%) 40 38	13, 50, 93, 109	0

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	29	TYR	9.2
1	C	1	MET	6.9
2	D	361	ASP	6.9
1	A	1	MET	6.9
2	B	74	LEU	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ZN	C	900	1/1	1.00	0.21	1.98	26,26,26,26	0
3	ZN	A	800	1/1	0.98	0.20	1.24	34,34,34,34	0
4	SF4	D	1000	8/8	0.98	0.21	1.02	17,26,32,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.