



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:42 PM GMT

PDB ID : 3RT9
Title : Crystal structure of tm0922, a fusion of a domain of unknown function and ADP/ATP-dependent NAD(P)H-hydrate dehydratase from *Thermotoga maritima* soaked with Coenzyme A
Authors : Shumilin, I.A.; Cymborowski, M.; Lesley, S.A.; Minor, W.
Deposited on : 2011-05-03
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	502	
2	B	6	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	COA	A	494	-	-	-	X
4	COA	A	495	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	489	3742	2388	631	712	11	0	3	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-11	MET	-	EXPRESSION TAG	UNP Q9X024
A	-10	GLY	-	EXPRESSION TAG	UNP Q9X024
A	-9	SER	-	EXPRESSION TAG	UNP Q9X024
A	-8	ASP	-	EXPRESSION TAG	UNP Q9X024
A	-7	LYS	-	EXPRESSION TAG	UNP Q9X024
A	-6	ILE	-	EXPRESSION TAG	UNP Q9X024
A	-5	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-4	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-3	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-2	HIS	-	EXPRESSION TAG	UNP Q9X024
A	-1	HIS	-	EXPRESSION TAG	UNP Q9X024
A	0	HIS	-	EXPRESSION TAG	UNP Q9X024

- Molecule 2 is a protein called Unknown peptide, probably from expression host.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	6	52	37	7	8	0	0	0

- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		

- Molecule 4 is COENZYME A (three-letter code: COA) (formula: C₂₁H₃₆N₇O₁₆P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
4	A	1	Total 48	C 21	N 7	O 16	P 3	S 1	0	0
4	A	1	Total 31	C 10	N 5	O 13	P 3		0	0
4	A	1	Total 27	C 10	N 5	O 10	P 2		0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	181	Total	O	0	0
			181	181		
5	B	2	Total	O	0	0
			2	2		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein



- Molecule 2: Unknown peptide, probably from expression host



There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, α , β , γ	122.09Å 122.09Å 155.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 1.95 37.59 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-1.95) 99.4 (37.59-1.95)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.157 , 0.192 0.166 , 0.198	Depositor DCC
R_{free} test set	2150 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 55.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 42640 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4132	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/3803	0.84	1/5145 (0.0%)
2	B	1.18	0/54	0.83	0/73
All	All	0.90	0/3857	0.84	1/5218 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	ASP	CB-CG-OD1	5.62	123.36	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3742	0	3856	30	0
2	B	52	0	45	0	0
3	A	1	0	0	0	0
4	A	154	0	86	12	0
5	A	181	0	0	5	0
5	B	2	0	0	0	0
All	All	4132	0	3987	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:LEU:HD23	5:A:662:HOH:O	1.70	0.91
1:A:243:LEU:CD2	5:A:662:HOH:O	2.20	0.89
1:A:117:PHE:CE1	4:A:492:COA:H1B	2.22	0.74
1:A:152:ILE:HD11	1:A:187:LEU:HD12	1.73	0.71
1:A:201:HIS:HB3	5:A:653:HOH:O	1.96	0.66
1:A:2:LYS:HG3	4:A:492:COA:H132	1.81	0.62
1:A:320:ASN:OD1	1:A:320:ASN:C	2.40	0.59
1:A:117:PHE:HE1	4:A:492:COA:H1B	1.65	0.59
1:A:158:LYS:HE3	4:A:494:COA:C2A	2.34	0.57
1:A:243:LEU:HD21	5:A:662:HOH:O	1.96	0.57
1:A:457:LEU:HD21	1:A:488:LEU:HD22	1.88	0.55
4:A:492:COA:O5A	4:A:492:COA:O9P	2.24	0.54
1:A:121:LEU:HD12	4:A:492:COA:N7A	2.23	0.53
4:A:492:COA:O5A	4:A:492:COA:H143	2.10	0.52
4:A:492:COA:O5A	4:A:492:COA:C9P	2.60	0.49
1:A:5:ASP:OD2	4:A:492:COA:H62	2.12	0.49
1:A:358:LYS:HD3	1:A:358:LYS:N	2.28	0.49
1:A:257:LYS:NZ	5:A:665:HOH:O	2.47	0.47
4:A:493:COA:H131	4:A:493:COA:O9P	2.16	0.46
1:A:216:VAL:HG12	1:A:488:LEU:HD21	1.97	0.45
1:A:367:PRO:HD3	1:A:402:LYS:O	2.17	0.45
1:A:1:MET:HE1	1:A:203:ILE:HD13	1.99	0.45
1:A:262:LEU:HD11	1:A:283:ILE:CD1	2.47	0.44
1:A:253:MET:HG3	1:A:279:PHE:CD2	2.53	0.43
1:A:253:MET:HG3	1:A:279:PHE:CE2	2.54	0.43
1:A:251:SER:O	1:A:435:GLY:HA3	2.19	0.42
4:A:493:COA:O9P	4:A:493:COA:CDP	2.67	0.42
1:A:7:LEU:HD22	1:A:11[A]:GLU:HG3	2.01	0.42
1:A:106:LEU:HD13	1:A:139:SER:HB3	2.02	0.41
1:A:50:GLY:HA2	1:A:80:THR:HG23	2.02	0.41
1:A:177:ILE:HD13	1:A:385:TYR:CZ	2.56	0.41
1:A:379:GLY:HA2	4:A:494:COA:H4B	2.03	0.40
1:A:262:LEU:HD11	1:A:283:ILE:HD12	2.03	0.40
1:A:121:LEU:HD23	1:A:160:LEU:HD12	2.02	0.40
1:A:264:LYS:HE2	1:A:309:VAL:CG1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	490/502 (98%)	479 (98%)	11 (2%)	0	100	100
2	B	4/6 (67%)	4 (100%)	0	0	100	100
All	All	494/508 (97%)	483 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	413/422 (98%)	408 (99%)	5 (1%)	78	75
2	B	4/4 (100%)	4 (100%)	0	100	100
All	All	417/426 (98%)	412 (99%)	5 (1%)	78	75

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	186	ASP
1	A	213	ARG
1	A	220	LEU
1	A	358	LYS
1	A	413	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	COA	A	492	-	40,50,50	1.88	3 (7%)	50,75,75	2.64	14 (28%)
4	COA	A	493	-	40,50,50	1.63	4 (10%)	50,75,75	2.02	11 (22%)
4	COA	A	494	-	26,33,50	1.44	3 (11%)	33,52,75	2.31	5 (15%)
4	COA	A	495	-	24,29,50	1.38	3 (12%)	28,45,75	2.37	4 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	COA	A	492	-	-	1/44/64/64	0/3/3/3
4	COA	A	493	-	-	1/44/64/64	0/3/3/3
4	COA	A	494	-	-	0/17/37/64	0/3/3/3
4	COA	A	495	-	-	0/11/31/64	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	493	COA	O4B-C1B	2.22	1.44	1.41
4	A	495	COA	C2A-N1A	2.67	1.39	1.33
4	A	492	COA	C2A-N1A	2.81	1.39	1.33
4	A	493	COA	C2A-N1A	2.81	1.39	1.33
4	A	493	COA	C2A-N3A	2.99	1.37	1.32
4	A	494	COA	P2A-O4A	3.10	1.61	1.51
4	A	494	COA	C2A-N1A	3.20	1.40	1.33
4	A	495	COA	P1A-O1A	3.42	1.62	1.51
4	A	492	COA	C2A-N3A	3.87	1.39	1.32
4	A	495	COA	C2A-N3A	3.88	1.39	1.32
4	A	494	COA	C2A-N3A	4.08	1.39	1.32
4	A	493	COA	O9P-C9P	8.05	1.39	1.23
4	A	492	COA	O9P-C9P	10.14	1.43	1.23

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	494	COA	N3A-C2A-N1A	-11.13	120.37	128.89
4	A	495	COA	N3A-C2A-N1A	-11.09	120.41	128.89
4	A	492	COA	N3A-C2A-N1A	-10.75	120.66	128.89
4	A	493	COA	N3A-C2A-N1A	-8.64	122.28	128.89
4	A	492	COA	C2B-C1B-N9A	-7.31	103.13	114.29
4	A	492	COA	C1B-N9A-C4A	-4.15	120.69	126.94
4	A	493	COA	C2B-C1B-N9A	-4.12	108.00	114.29
4	A	492	COA	C7P-N8P-C9P	-3.37	115.86	122.53
4	A	492	COA	P3B-O3B-C3B	-3.37	113.49	121.56
4	A	494	COA	P1A-O3A-P2A	-3.23	121.83	132.67
4	A	493	COA	P2A-O3A-P1A	-2.96	124.42	132.73
4	A	492	COA	O4B-C4B-C3B	-2.87	98.24	104.86
4	A	492	COA	CEP-CBP-CDP	-2.80	103.66	109.28
4	A	493	COA	OAP-CAP-C9P	-2.70	104.18	110.38
4	A	493	COA	O5P-C5P-C6P	-2.66	117.40	121.98
4	A	492	COA	O5P-C5P-C6P	-2.62	117.47	121.98
4	A	494	COA	C4A-C5A-N7A	-2.37	107.30	109.48
4	A	493	COA	C1B-N9A-C4A	-2.32	123.44	126.94
4	A	493	COA	CDP-CBP-CCP	-2.22	105.63	108.50
4	A	493	COA	O3A-P2A-O6A	-2.20	97.10	102.94
4	A	493	COA	C4A-C5A-N7A	-2.19	107.46	109.48
4	A	492	COA	C4A-C5A-N7A	-2.14	107.51	109.48
4	A	495	COA	C2B-C1B-N9A	-2.14	111.02	114.29
4	A	495	COA	O4B-C1B-N9A	2.00	112.29	108.10
4	A	494	COA	O2A-P1A-O3A	2.07	114.47	105.09

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	492	COA	O2B-C2B-C3B	2.32	117.85	111.16
4	A	495	COA	P3B-O3B-C3B	2.42	127.36	121.56
4	A	492	COA	O3B-C3B-C4B	2.49	119.77	109.99
4	A	494	COA	C4B-O4B-C1B	2.64	112.62	109.72
4	A	492	COA	O6A-CCP-CBP	2.98	115.33	110.55
4	A	493	COA	C6P-C5P-N4P	3.61	122.74	116.46
4	A	493	COA	CAP-C9P-N8P	3.78	124.84	116.47
4	A	492	COA	O4B-C1B-N9A	4.99	118.54	108.10
4	A	492	COA	CEP-CBP-CCP	6.10	116.41	108.50

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	492	COA	CAP-C9P-N8P-C7P
4	A	493	COA	CAP-C9P-N8P-C7P

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	492	COA	8	0
4	A	493	COA	2	0
4	A	494	COA	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	489/502 (97%)	0.36	37 (7%) 17 26	20, 30, 53, 90	1 (0%)
2	B	6/6 (100%)	0.18	0 100 100	29, 32, 47, 47	0
All	All	495/508 (97%)	0.35	37 (7%) 17 26	20, 30, 53, 90	1 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	100	GLN	5.3
1	A	122	ARG	5.0
1	A	1	MET	4.5
1	A	101	PHE	3.8
1	A	206	ILE	3.7
1	A	105	ILE	3.6
1	A	290	GLU	3.6
1	A	107	ASN	3.4
1	A	358	LYS	3.3
1	A	77	LYS	3.3
1	A	102	GLU	3.3
1	A	433	LEU	3.2
1	A	394	GLU	3.2
1	A	128	GLU	2.9
1	A	356	GLU	2.8
1	A	222	GLU	2.8
1	A	365	PRO	2.6
1	A	308	ASP	2.6
1	A	138	LYS	2.6
1	A	412	GLU	2.6
1	A	201	HIS	2.5
1	A	2	LYS	2.5
1	A	400	VAL	2.5
1	A	401	LEU	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	140	GLY	2.4
1	A	35	LEU	2.3
1	A	390	GLU	2.2
1	A	297	GLN	2.2
1	A	19	LEU	2.2
1	A	108	GLU	2.2
1	A	40	ASP	2.2
1	A	402	LYS	2.1
1	A	406	THR	2.1
1	A	207[A]	ASN	2.1
1	A	213	ARG	2.1
1	A	300	GLN	2.1
1	A	393	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	COA	A	494	31/48	0.80	0.24	4.56	33,73,94,94	0
4	COA	A	495	27/48	0.64	0.32	2.80	80,92,104,104	0
4	COA	A	492	48/48	0.80	0.26	1.59	34,70,87,88	0
4	COA	A	493	48/48	0.93	0.14	-0.17	28,45,81,83	0
3	K	A	491	1/1	1.00	0.05	-1.88	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.