



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 2, 2016 – 12:03 AM GMT

PDB ID : 9RUB
Title : CRYSTAL STRUCTURE OF ACTIVATED RIBULOSE-1,5-BISPHOSPHATE CARBOXYLASE COMPLEXED WITH ITS SUBSTRATE, RIBULOSE-1,5-BISPHOSPHATE
Authors : Lundqvist, T.; Schneider, G.
Deposited on : 1990-11-28
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	466	
1	B	466	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FMT	A	601	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called RIBULOSE-1,5-BISPHOSPHATE CARBOXYLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	8	0	0
			3502	2213	613	658	18			
1	B	458	Total	C	N	O	S	1	0	0
			3498	2211	612	657	18			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	91	ASP	HIS	CONFLICT	UNP P04718
B	91	ASP	HIS	CONFLICT	UNP P04718

- Molecule 2 is SUGAR (RIBULOSE-1,5-DIPHOSPHATE) (three-letter code: RUB) (formula: $C_5H_{12}O_{11}P_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			18	5	11	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	O	P	0	0
			18	5	11	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			3	1	2		
4	B	1	Total	C	O	0	0
			3	1	2		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: RIBULOSE-1,5-BISPHOSPHATE CARBOXYLASE



• Molecule 1: RIBULOSE-1,5-BISPHOSPHATE CARBOXYLASE



ARG	A396	G328	D263	P200	P139
SER	F397	K329	G264	F201	S140
ALA	G398	M330	Y265	A202	V141
LEU	H399			P203	I142
PRO	D401	E333	G268	L204	I143
ALA	G402	S334	A269	R205	S144
	P403	S335	A270	D206	A145
		I336	A271	T207	L146
		R337	I272	I208	H147
	G406	A338	T273	A209	K148
	A407	I339	T274	L210	V149
	R408	A340	A275	V211	L150
	S409	Y341	R276	A212	G151
		M342	R277	D213	A152
		L343	R278	A214	P153
		T344		M215	E154
	Q412	Q345		R216	V155
	A413		D281	R217	D156
	N414		N282	A218	G157
	Q415	Q349	F283	Q219	G158
	A416	G350	L284	D220	L159
	A417	P351	H285	E221	V160
	R418		Y286	T222	V161
	D419	R354	R287	G223	G162
		Q355	R288	E224	T163
	P422	S356	A289	A225	I164
	V423	W357	G290	R226	I165
	L424			L227	K166
	D425	M360	V294	F228	P167
	Y426	K361	T295	S229	K168
	A427	A362	S296	A230	L168
	R428	G363	P297	N231	G170
	E429	T364	Q298		
	H430	P365	S299	A234	L171
	K431	I366	R300	D235	P172
	E432	I367	R301	P173	
		S368	G302	K174	P175
	R435		Y303	P237	
	A436	M371	T304		F176
	F437	N372	A305	I240	A177
	E438	A373	F306		E178
	S439	L374	V307	R243	A179
	P441	R375	H308	G244	C180
	G442	M376	G309	E245	H181
	D443	P377	R310	Y246	A182
	A444		K311	V247	F183
	D445	F380	A312	L248	H184
	Q446	E381	R313	E249	L185
	L447	N382	L314	T250	G186
	Y448		Q315	F251	G187
		N385	G316	G252	D188
	W451	A386	A317	E253	F189
	R452	N387	S318	N254	I190
	K453	V388	G319	A255	K191
	A454	I389	I320	S256	N192
	L455	L390	H321	H257	D193
	G456	T391	T322	V258	F194
	W457	A392	G323	A259	P195
	E458	G393		L260	Q196
	D459	G394	G326	L261	
	THR	G395	F327	V262	Q199

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	65.50Å 70.60Å 104.10Å 90.00° 92.10° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.60	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.60)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.199 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	7044	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: RUB, FMT, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	4/3586 (0.1%)	2.90	360/4859 (7.4%)
1	B	1.02	1/3582 (0.0%)	2.84	300/4854 (6.2%)
All	All	1.08	5/7168 (0.1%)	2.87	660/9713 (6.8%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	458	GLU	CG-CD	21.57	1.84	1.51
1	A	459	ASP	CG-OD2	13.53	1.56	1.25
1	A	460	THR	N-CA	-10.23	1.25	1.46
1	B	457	VAL	C-O	-9.22	1.05	1.23
1	A	459	ASP	CB-CG	-5.24	1.40	1.51

The worst 5 of 660 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	428	ARG	CD-NE-CZ	38.62	177.67	123.60
1	B	65	ARG	CD-NE-CZ	26.96	161.35	123.60
1	A	78	ARG	CD-NE-CZ	23.14	156.00	123.60
1	A	96	ASP	CB-CG-OD1	20.82	137.03	118.30
1	A	459	ASP	C-N-CA	20.41	172.74	121.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	457	VAL	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3502	0	3368	276	0
1	B	3498	0	3371	244	0
2	A	18	0	8	3	0
2	B	18	0	7	4	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	3	0	0	2	0
4	B	3	0	0	0	0
All	All	7044	0	6754	495	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 36.

The worst 5 of 495 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:234:ALA:HB3	1:B:240:ILE:HG13	1.25	1.11
1:B:321:HIS:HB3	2:B:700:RUB:O6P	1.56	1.05
1:A:167:PRO:HD3	1:B:59:THR:HG21	1.41	1.01
1:A:285:HIS:HE2	1:A:321:HIS:HE2	1.05	0.94
1:B:322:THR:HG22	1:B:323:GLY:H	1.34	0.92

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/466 (98%)	330 (72%)	82 (18%)	45 (10%)	1	0
1	B	456/466 (98%)	360 (79%)	60 (13%)	36 (8%)	1	1
All	All	913/932 (98%)	690 (76%)	142 (16%)	81 (9%)	1	1

5 of 81 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	11	ALA
1	A	38	TYR
1	A	129	PRO
1	A	136	PHE
1	A	144	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/354 (98%)	266 (76%)	82 (24%)	1	1
1	B	348/354 (98%)	268 (77%)	80 (23%)	1	2
All	All	696/708 (98%)	534 (77%)	162 (23%)	1	2

5 of 162 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	423	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	28	TYR
1	B	409	SER
1	A	432	GLU
1	B	2	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	387	ASN
1	B	3	GLN
1	B	382	ASN
1	A	385	ASN
1	B	385	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	RUB	A	600	3	17,17,17	2.92	10 (58%)	15,25,25	1.40	4 (26%)
4	FMT	A	601	1,3	0,2,2	0.00	-	0,1,1	0.00	-
2	RUB	B	700	3,4	17,17,17	1.46	2 (11%)	15,25,25	0.93	0
4	FMT	B	701	1,3,2	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	RUB	A	600	3	-	0/20/20/20	0/0/0/0
4	FMT	A	601	1,3	-	0/0/0/0	0/0/0/0
2	RUB	B	700	3,4	-	0/20/20/20	0/0/0/0
4	FMT	B	701	1,3,2	-	0/0/0/0	0/0/0/0

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600	RUB	O2-C2	2.03	1.25	1.21
2	A	600	RUB	O1-C1	2.23	1.44	1.43
2	A	600	RUB	P2-O5P	2.39	1.63	1.54
2	A	600	RUB	C3-C2	2.52	1.61	1.52
2	A	600	RUB	P2-O5	2.70	1.69	1.60

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	RUB	O6P-P2-O4P	-2.27	103.28	110.58
2	A	600	RUB	O3P-P1-O1	2.16	112.79	106.56
2	A	600	RUB	O5-P2-O4P	2.23	112.82	107.14
2	A	600	RUB	O6P-P2-O5	2.35	113.34	106.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600	RUB	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	601	FMT	2	0
2	B	700	RUB	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.