



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:43 PM GMT

PDB ID : 3RV3
Title : Crystal structure of E.coli biotin carboxylase in complex with two ADP and one Mg ion
Authors : Chou, C.Y.; Tong, L.
Deposited on : 2011-05-05
Resolution : 1.91 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	5832 (1.94-1.90)
Clashscore	102246	6540 (1.94-1.90)
Ramachandran outliers	100387	6464 (1.94-1.90)
Sidechain outliers	100360	6465 (1.94-1.90)
RSRZ outliers	91569	5846 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	<div> <div>4%</div> <div>88%</div> <div>11%</div> </div>
1	B	452	<div> <div>10%</div> <div>87%</div> <div>12%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7740 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Biotin carboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	447	Total	C	N	O	S	0	0	0
			3441	2166	616	637	22			
1	B	447	Total	C	N	O	S	0	0	0
			3442	2168	615	637	22			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1C	GLY	-	EXPRESSION TAG	UNP P24182
A	1B	SER	-	EXPRESSION TAG	UNP P24182
A	1A	HIS	-	EXPRESSION TAG	UNP P24182
B	1C	GLY	-	EXPRESSION TAG	UNP P24182
B	1B	SER	-	EXPRESSION TAG	UNP P24182
B	1A	HIS	-	EXPRESSION TAG	UNP P24182

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Mg	0	0
			1	1		
3	A	1	Total	Mg	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	367	Total	O	0	0
			367	367		
4	B	380	Total	O	0	0
			380	380		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Biotin carboxylase



• Molecule 1: Biotin carboxylase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	171.36Å 58.01Å 85.24Å 90.00° 94.64° 90.00°	Depositor
Resolution (Å)	30.00 – 1.91 29.43 – 1.91	Depositor EDS
% Data completeness (in resolution range)	99.4 (30.00-1.91) 99.4 (29.43-1.91)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.81 (at 1.91Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.208 , 0.263 0.208 , 0.262	Depositor DCC
R_{free} test set	3280 reflections (5.34%)	DCC
Wilson B-factor (Å ²)	18.2	Xtriage
Anisotropy	0.525	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	1 of 64727 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7740	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/3505	0.62	1/4730 (0.0%)
1	B	0.51	0/3505	0.61	0/4730
All	All	0.52	0/7010	0.61	1/9460 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	154	TYR	N-CA-C	5.75	126.54	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	154	TYR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3441	0	3460	46	1
1	B	3442	0	3467	46	1
2	A	54	0	24	1	0
2	B	54	0	24	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	367	0	0	17	0
4	B	380	0	0	15	0
All	All	7740	0	6975	94	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:288:GLU:HG2	4:A:684:HOH:O	1.38	1.21
1:A:157:ILE:HD13	4:A:668:HOH:O	1.41	1.18
1:B:397:VAL:HB	4:B:820:HOH:O	1.46	1.14
1:B:407:GLN:HG2	4:B:466:HOH:O	1.66	0.94
1:B:190:LYS:HA	1:B:194:SER:O	1.68	0.93
1:A:18:LEU:HD23	4:A:743:HOH:O	1.77	0.84
1:A:342:GLU:HG3	4:A:752:HOH:O	1.76	0.84
1:B:158:ILE:CG2	1:B:198:VAL:HG11	2.09	0.82
1:A:108:GLU:HG2	4:A:689:HOH:O	1.80	0.80
1:B:234:ARG:NH1	1:B:234:ARG:HB3	1.96	0.79
1:A:156:VAL:HB	4:A:710:HOH:O	1.82	0.79
1:A:200:MET:HG2	4:A:710:HOH:O	1.82	0.79
1:A:19:ARG:HG3	4:A:743:HOH:O	1.84	0.77
1:B:181:SER:O	1:B:185:THR:HG23	1.87	0.74
1:B:234:ARG:NH2	4:B:591:HOH:O	2.18	0.72
2:B:1003:ADP:O2B	4:B:771:HOH:O	2.07	0.72
1:B:234:ARG:HH11	1:B:234:ARG:HB3	1.55	0.70
2:A:1002:ADP:O2A	4:A:684:HOH:O	2.11	0.68
1:A:200:MET:CG	4:A:710:HOH:O	2.41	0.67
1:A:116:LYS:O	1:A:120:ILE:HG12	1.94	0.67
1:A:22:LYS:HE2	4:B:619:HOH:O	1.95	0.65
1:A:18:LEU:CD2	1:A:22:LYS:HE3	2.26	0.64
1:A:27:LYS:NZ	4:A:734:HOH:O	2.30	0.63
1:A:363:PHE:HD1	4:A:596:HOH:O	1.81	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:205:GLU:O	1:B:438:HIS:HD2	1.83	0.61
1:A:370:HIS:CD2	1:A:370:HIS:H	2.18	0.61
1:B:147:ALA:O	1:B:151:ARG:HG3	2.00	0.61
1:A:23:GLU:OE1	1:A:310:LYS:HE3	1.99	0.61
1:B:158:ILE:HG23	1:B:198:VAL:HG11	1.83	0.61
1:A:18:LEU:CD2	4:A:743:HOH:O	2.40	0.60
1:B:252:LEU:HD12	4:B:565:HOH:O	2.01	0.60
1:B:255:TYR:O	1:B:259:ARG:HG2	2.04	0.58
1:B:403:LYS:O	1:B:407:GLN:HG3	2.03	0.58
1:B:158:ILE:CG2	1:B:198:VAL:CG1	2.79	0.58
1:A:205:GLU:O	1:A:438:HIS:CD2	2.56	0.58
1:A:155:PRO:HD2	1:A:202:LYS:HE3	1.87	0.57
1:A:234:ARG:HB3	1:A:234:ARG:HH11	1.70	0.57
1:B:158:ILE:HG23	1:B:198:VAL:CG1	2.34	0.57
1:B:254:ARG:HG3	4:B:648:HOH:O	2.05	0.56
1:A:228:ARG:HH21	1:A:294:GLN:NE2	2.03	0.56
1:A:180:GLN:O	1:A:184:MET:HG3	2.06	0.56
1:A:93:GLU:OE1	1:A:111:ARG:NH1	2.39	0.55
1:A:18:LEU:HD21	1:A:22:LYS:HE3	1.88	0.54
1:B:394:ASN:N	4:B:820:HOH:O	2.17	0.53
1:B:158:ILE:HG22	1:B:198:VAL:HG11	1.88	0.53
1:A:346:THR:OG1	1:A:348:LEU:HG	2.10	0.52
1:A:18:LEU:HD23	1:A:22:LYS:HE3	1.91	0.51
1:B:234:ARG:CB	1:B:234:ARG:HH11	2.22	0.51
1:B:228:ARG:HH21	1:B:294:GLN:NE2	2.09	0.51
1:A:234:ARG:HB3	1:A:234:ARG:NH1	2.26	0.51
1:A:205:GLU:O	1:A:438:HIS:HD2	1.93	0.50
1:B:251:GLU:HG3	4:B:468:HOH:O	2.12	0.50
1:A:141:ASP:HB2	4:A:495:HOH:O	2.13	0.49
1:A:23:GLU:OE1	1:A:310:LYS:CE	2.60	0.49
1:B:295:VAL:HG23	4:B:771:HOH:O	2.13	0.48
1:B:57:VAL:HG13	1:B:58:LYS:HG3	1.95	0.48
1:B:250:PRO:O	1:B:254:ARG:HG2	2.14	0.48
1:A:169:MET:HE3	4:A:668:HOH:O	2.13	0.47
1:B:307:ASP:O	1:B:311:GLU:HG2	2.14	0.47
1:B:344:PRO:HG3	1:B:416:THR:O	2.15	0.47
1:B:228:ARG:HH21	1:B:294:GLN:HE21	1.62	0.47
1:B:347:PHE:HE1	1:B:420:LEU:HD22	1.79	0.47
1:B:113:MET:CE	1:B:267:ILE:HG21	2.44	0.47
1:B:206:ASN:N	1:B:207:PRO:HD3	2.30	0.46
1:B:292:ARG:NH1	4:B:771:HOH:O	2.48	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:258:GLU:HG2	4:B:488:HOH:O	2.16	0.46
1:A:263:ALA:O	1:A:267:ILE:HG12	2.16	0.45
1:A:157:ILE:HD12	1:A:171:VAL:HG22	1.98	0.45
1:A:82:TYR:CZ	1:A:295:VAL:HG22	2.52	0.45
1:B:206:ASN:N	1:B:207:PRO:CD	2.80	0.45
1:A:93:GLU:HG3	1:A:107:ALA:CB	2.46	0.45
1:B:249:THR:OG1	1:B:251:GLU:HG2	2.16	0.44
1:A:131:VAL:CG1	1:A:204:LEU:HD11	2.47	0.44
1:B:253:ARG:NH1	4:B:813:HOH:O	2.51	0.44
1:B:259:ARG:NE	1:B:259:ARG:HA	2.33	0.44
1:B:82:TYR:CZ	1:B:295:VAL:HG22	2.52	0.44
1:A:442:LYS:HG3	4:A:548:HOH:O	2.18	0.44
1:A:296:GLU:O	1:A:299:VAL:HG22	2.18	0.43
1:B:2:LEU:HA	1:B:76:VAL:HG21	1.99	0.43
1:B:113:MET:HE2	1:B:267:ILE:HG21	1.98	0.43
1:B:205:GLU:HG3	4:B:531:HOH:O	2.18	0.42
1:B:228:ARG:NH2	1:B:294:GLN:HE21	2.17	0.42
1:A:105:PRO:HG2	1:A:291:THR:HB	2.01	0.42
1:B:13:ILE:HB	1:B:82:TYR:CE2	2.55	0.42
1:A:228:ARG:NH2	1:A:294:GLN:HE21	2.17	0.42
1:A:131:VAL:HG13	1:A:204:LEU:HD11	2.01	0.42
1:A:13:ILE:HB	1:A:82:TYR:CE2	2.55	0.42
1:A:157:ILE:HG13	1:A:170:ARG:O	2.19	0.41
1:A:440:LEU:O	1:A:444:LEU:HG	2.20	0.41
1:A:417:ASN:HB3	4:A:752:HOH:O	2.20	0.41
1:B:353:LYS:HA	1:B:376:THR:HA	2.03	0.41
1:B:234:ARG:HG2	1:B:235:ARG:HD2	2.02	0.41
1:A:27:LYS:HB2	1:A:27:LYS:HE3	1.74	0.41
1:B:205:GLU:N	4:B:756:HOH:O	2.53	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:98:SER:CB	1:B:251:GLU:OE2[4_546]	1.95	0.25

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	445/452 (98%)	434 (98%)	10 (2%)	1 (0%)	52	42
1	B	445/452 (98%)	429 (96%)	15 (3%)	1 (0%)	52	42
All	All	890/904 (98%)	863 (97%)	25 (3%)	2 (0%)	52	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	TYR
1	B	195	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/363 (99%)	357 (99%)	2 (1%)	90	90
1	B	359/363 (99%)	352 (98%)	7 (2%)	65	58
All	All	718/726 (99%)	709 (99%)	9 (1%)	76	73

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	154	TYR
1	A	363	PHE
1	B	5	ILE
1	B	278	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	345	ASN
1	B	428	GLU
1	B	432	HIS
1	B	442	LYS
1	B	444	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	290	ASN
1	A	294	GLN
1	A	370	HIS
1	A	407	GLN
1	A	426	ASN
1	A	432	HIS
1	A	438	HIS
1	B	206	ASN
1	B	236	HIS
1	B	237	GLN
1	B	294	GLN
1	B	340	ASN
1	B	404	ASN
1	B	431	GLN
1	B	438	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ADP	A	1000	-	22,29,29	1.08	2 (9%)	27,45,45	1.76	2 (7%)
2	ADP	A	1002	3	22,29,29	1.08	1 (4%)	27,45,45	2.31	6 (22%)
2	ADP	B	1001	3	22,29,29	0.94	1 (4%)	27,45,45	2.14	5 (18%)
2	ADP	B	1003	-	22,29,29	1.05	1 (4%)	27,45,45	1.96	5 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	A	1000	-	-	0/12/32/32	0/3/3/3
2	ADP	A	1002	3	-	0/12/32/32	0/3/3/3
2	ADP	B	1001	3	-	0/12/32/32	0/3/3/3
2	ADP	B	1003	-	-	0/12/32/32	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1000	ADP	C2-N3	2.57	1.36	1.32
2	B	1001	ADP	C5-C4	2.85	1.46	1.40
2	B	1003	ADP	C5-C4	2.96	1.47	1.40
2	A	1000	ADP	C5-C4	2.98	1.47	1.40
2	A	1002	ADP	C5-C4	3.16	1.47	1.40

All (18) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1003	ADP	N3-C2-N1	-8.20	122.61	128.89
2	B	1001	ADP	N3-C2-N1	-7.99	122.77	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1002	ADP	N3-C2-N1	-7.70	123.00	128.89
2	A	1000	ADP	N3-C2-N1	-7.28	123.32	128.89
2	A	1002	ADP	C2'-C1'-N9	-6.90	103.75	114.29
2	B	1001	ADP	C2'-C1'-N9	-4.68	107.14	114.29
2	A	1002	ADP	C1'-N9-C4	-3.02	122.39	126.94
2	A	1000	ADP	C4-C5-N7	-3.00	106.72	109.48
2	B	1001	ADP	C1'-N9-C4	-2.76	122.77	126.94
2	B	1001	ADP	C4-C5-N7	-2.70	106.99	109.48
2	B	1003	ADP	C2'-C1'-N9	-2.35	110.70	114.29
2	B	1003	ADP	C1'-N9-C4	-2.29	123.49	126.94
2	B	1003	ADP	C4-C5-N7	-2.20	107.46	109.48
2	A	1002	ADP	C4-C5-N7	-2.02	107.62	109.48
2	A	1002	ADP	O4'-C1'-N9	2.02	112.34	108.10
2	B	1003	ADP	C2-N1-C6	2.09	122.50	118.77
2	B	1001	ADP	C2-N1-C6	2.16	122.62	118.77
2	A	1002	ADP	C2-N1-C6	2.35	122.97	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1002	ADP	1	0
2	B	1003	ADP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	447/452 (98%)	0.49	19 (4%)	39 43	7, 16, 29, 35	1 (0%)
1	B	447/452 (98%)	0.71	44 (9%)	10 11	9, 18, 34, 43	1 (0%)
All	All	894/904 (98%)	0.60	63 (7%)	19 22	7, 17, 32, 43	2 (0%)

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	447	GLN	7.8
1	B	446	LEU	6.6
1	B	445	GLY	5.9
1	B	194	SER	4.4
1	A	154	TYR	4.4
1	A	348	LEU	4.3
1	B	193	PHE	4.2
1	A	346	THR	4.0
1	A	345	ASN	3.9
1	B	141	ASP	3.7
1	A	347	PHE	3.2
1	A	192	ALA	3.2
1	B	322	SER	3.2
1	A	344	PRO	3.2
1	B	1	MET	3.1
1	B	205	GLU	3.1
1	B	191	ALA	3.0
1	B	206	ASN	3.0
1	B	140	ASP	2.9
1	B	189	ALA	2.9
1	B	143	ASP	2.9
1	B	127	GLY	2.8
1	B	432	HIS	2.8
1	B	138	LEU	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	255	TYR	2.7
1	B	139	GLY	2.7
1	B	185	THR	2.7
1	B	254	ARG	2.7
1	A	206	ASN	2.6
1	B	281	ASN	2.6
1	A	204	LEU	2.6
1	B	345	ASN	2.5
1	B	128	VAL	2.5
1	B	190	LYS	2.5
1	B	142	MET	2.5
1	A	191	ALA	2.5
1	B	150	LYS	2.5
1	B	160	ALA	2.4
1	B	347	PHE	2.4
1	B	187	ALA	2.4
1	B	148	ILE	2.4
1	A	322	SER	2.4
1	A	326	GLU	2.4
1	B	173	ARG	2.4
1	B	152	ILE	2.3
1	A	151	ARG	2.2
1	B	323	ILE	2.2
1	B	195	ASN	2.2
1	B	126	ALA	2.2
1	B	287	ILE	2.2
1	A	203	TYR	2.2
1	A	376	THR	2.2
1	A	141	ASP	2.1
1	A	125	LYS	2.1
1	B	72	ILE	2.1
1	B	329	HIS	2.1
1	B	259	ARG	2.1
1	B	151	ARG	2.1
1	A	1(A)	HIS	2.1
1	B	125	LYS	2.0
1	B	348	LEU	2.0
1	B	331	ARG	2.0
1	A	176	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ADP	A	1002	27/27	0.93	0.14	-0.08	13,21,22,25	0
2	ADP	B	1003	27/27	0.96	0.12	-0.70	12,17,19,20	0
2	ADP	A	1000	27/27	0.96	0.12	-0.78	12,16,18,19	0
2	ADP	B	1001	27/27	0.96	0.12	-0.78	14,17,20,20	0
3	MG	B	1005	1/1	0.96	0.07	-4.14	18,18,18,18	0
3	MG	A	1004	1/1	0.94	0.07	-	15,15,15,15	0

6.5 Other polymers [i](#)

There are no such residues in this entry.