



wwPDB X-ray Structure Validation Summary Report

Feb 1, 2016 – 08:28 PM GMT

PDB ID : 4RX2
Title : A triple mutant in the omega-loop of TEM-1 beta-lactamase changes the substrate profile via a large conformational change and an altered general base for catalysis
Authors : Stojanoski, V.; Chow, D.; Hu, L.; Sankaran, B.; Gilbert, H.; Prasad, B.V.V.; Palzkill, T.
Deposited on : 2014-12-08
Resolution : 2.31 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4425 (2.34-2.30)
Clashscore	102246	5057 (2.34-2.30)
Ramachandran outliers	100387	5008 (2.34-2.30)
Sidechain outliers	100360	5007 (2.34-2.30)
RSRZ outliers	91569	4432 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	263	 83% 13% ..
1	B	263	 87% 9% ..
1	C	263	 85% 11% ..
1	D	263	 83% 13% ..
1	E	263	 85% 9% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	263	 86% 9% . .
1	G	263	 3% 86% 9% . .
1	H	263	 2% 84% 10% . .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 16545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-lactamase TEM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	255	1967	1227	350	380	10	0	1	0
1	B	256	1972	1229	351	382	10	0	2	0
1	C	258	1986	1240	353	383	10	0	1	0
1	D	256	1971	1229	351	381	10	0	1	0
1	E	252	1944	1212	346	376	10	0	1	0
1	F	254	1959	1223	348	378	10	0	1	0
1	G	252	1937	1212	346	369	10	0	0	0
1	H	253	1952	1219	347	376	10	0	2	0

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
A	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
A	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
A	182	THR	MET	ENGINEERED MUTATION	UNP P62593
B	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
B	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
B	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
B	182	THR	MET	ENGINEERED MUTATION	UNP P62593
C	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
C	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
C	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
C	182	THR	MET	ENGINEERED MUTATION	UNP P62593
D	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
D	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
D	182	THR	MET	ENGINEERED MUTATION	UNP P62593
E	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
E	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
E	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
E	182	THR	MET	ENGINEERED MUTATION	UNP P62593
F	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
F	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
F	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
F	182	THR	MET	ENGINEERED MUTATION	UNP P62593
G	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
G	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
G	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
G	182	THR	MET	ENGINEERED MUTATION	UNP P62593
H	165	TYR	TRP	ENGINEERED MUTATION	UNP P62593
H	166	TYR	GLU	ENGINEERED MUTATION	UNP P62593
H	167	GLY	PRO	ENGINEERED MUTATION	UNP P62593
H	182	THR	MET	ENGINEERED MUTATION	UNP P62593

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf		
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total 5	O 4	S 1	0	0
2	D	1	Total 5	O 4	S 1	0	0
2	E	1	Total 5	O 4	S 1	0	0
2	F	1	Total 5	O 4	S 1	0	0
2	G	1	Total 5	O 4	S 1	0	0
2	H	1	Total 5	O 4	S 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total 124	O 124	0	0
3	B	152	Total 152	O 152	0	0
3	C	112	Total 112	O 112	0	0
3	D	124	Total 124	O 124	0	0
3	E	70	Total 70	O 70	0	0
3	F	98	Total 98	O 98	0	0
3	G	64	Total 64	O 64	0	0
3	H	73	Total 73	O 73	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-lactamase TEM




- Molecule 1: Beta-lactamase TEM



- Molecule 1: Beta-lactamase TEM




- Molecule 1: Beta-lactamase TEM








• Molecule 1: Beta-lactamase TEM




• Molecule 1: Beta-lactamase TEM



• Molecule 1: Beta-lactamase TEM



• Molecule 1: Beta-lactamase TEM



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.32Å 83.30Å 95.92Å 90.06° 89.97° 90.02°	Depositor
Resolution (Å)	62.89 – 2.31 62.93 – 2.32	Depositor EDS
% Data completeness (in resolution range)	99.6 (62.89-2.31) 99.6 (62.93-2.32)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.12 (at 2.32Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
R, R_{free}	0.235 , 0.268 0.249 , 0.279	Depositor DCC
R_{free} test set	4081 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	16.4	Xtriage
Anisotropy	0.045	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 8.0	EDS
Estimated twinning fraction	0.070 for h,-k,-l 0.239 for -h,k,-l 0.074 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 80979 reflections	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	16545	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.30 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.7931e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	3/2003 (0.1%)	0.80	0/2708
1	B	0.79	2/2013 (0.1%)	0.82	2/2721 (0.1%)
1	C	0.71	0/2023	0.77	1/2736 (0.0%)
1	D	0.73	0/2007	0.78	0/2713
1	E	0.57	0/1977	0.74	1/2670 (0.0%)
1	F	0.56	0/1995	0.75	0/2697
1	G	0.52	0/1967	0.74	2/2658 (0.1%)
1	H	0.51	0/1993	0.75	2/2694 (0.1%)
All	All	0.68	5/15978 (0.0%)	0.77	8/21597 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	82[A]	SER	CB-OG	14.77	1.61	1.42
1	A	82[B]	SER	CB-OG	14.77	1.61	1.42
1	A	121	GLU	CD-OE1	-5.70	1.19	1.25
1	B	83	ARG	CZ-NH1	-5.31	1.26	1.33
1	B	83	ARG	CZ-NH2	-5.24	1.26	1.33

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	83	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	G	102	LEU	CA-CB-CG	5.96	129.01	115.30
1	C	277	ARG	NE-CZ-NH1	5.96	123.28	120.30
1	H	277	ARG	NE-CZ-NH1	-5.41	117.60	120.30
1	H	221	LEU	CB-CG-CD1	5.27	119.96	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1967	0	1975	21	0
1	B	1972	0	1979	20	0
1	C	1986	0	1996	25	0
1	D	1971	0	1978	25	0
1	E	1944	0	1955	17	0
1	F	1959	0	1969	20	0
1	G	1937	0	1953	17	0
1	H	1952	0	1966	17	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
2	G	5	0	0	0	0
2	H	5	0	0	1	0
3	A	124	0	0	8	0
3	B	152	0	0	13	0
3	C	112	0	0	9	0
3	D	124	0	0	7	0
3	E	70	0	0	4	0
3	F	98	0	0	9	0
3	G	64	0	0	8	0
3	H	73	0	0	8	0
All	All	16545	0	15771	157	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 157 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:207:LEU:HD12	3:F:478:HOH:O	1.46	1.13
1:C:111:LYS:HB3	3:C:428:HOH:O	1.65	0.94
1:B:85:ASP:HA	3:B:495:HOH:O	1.74	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:240:GLU:HG2	3:G:453:HOH:O	1.76	0.85
3:B:494:HOH:O	1:D:111:LYS:HE3	1.76	0.83

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/263 (96%)	243 (96%)	9 (4%)	0	100	100
1	B	254/263 (97%)	245 (96%)	9 (4%)	0	100	100
1	C	255/263 (97%)	246 (96%)	8 (3%)	1 (0%)	39	48
1	D	253/263 (96%)	246 (97%)	7 (3%)	0	100	100
1	E	245/263 (93%)	239 (98%)	6 (2%)	0	100	100
1	F	251/263 (95%)	246 (98%)	5 (2%)	0	100	100
1	G	246/263 (94%)	240 (98%)	6 (2%)	0	100	100
1	H	251/263 (95%)	245 (98%)	6 (2%)	0	100	100
All	All	2007/2104 (95%)	1950 (97%)	56 (3%)	1 (0%)	56	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	175	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/216 (98%)	204 (97%)	7 (3%)	45	61
1	B	212/216 (98%)	204 (96%)	8 (4%)	40	54
1	C	213/216 (99%)	202 (95%)	11 (5%)	29	38
1	D	211/216 (98%)	203 (96%)	8 (4%)	40	54
1	E	209/216 (97%)	201 (96%)	8 (4%)	40	54
1	F	210/216 (97%)	200 (95%)	10 (5%)	31	42
1	G	206/216 (95%)	196 (95%)	10 (5%)	31	41
1	H	210/216 (97%)	197 (94%)	13 (6%)	23	30
All	All	1682/1728 (97%)	1607 (96%)	75 (4%)	34	46

5 of 75 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	273	ASP
1	E	273	ASP
1	H	148	LEU
1	E	30	LEU
1	E	98	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	206	GLN
1	H	88	GLN
1	F	289	HIS
1	D	26	HIS
1	G	96	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	301	-	4,4,4	0.61	0	6,6,6	0.72	0
2	SO4	B	301	-	4,4,4	1.17	0	6,6,6	0.67	0
2	SO4	C	301	-	4,4,4	0.71	0	6,6,6	1.40	1 (16%)
2	SO4	D	301	-	4,4,4	0.66	0	6,6,6	1.69	2 (33%)
2	SO4	E	301	-	4,4,4	0.44	0	6,6,6	0.33	0
2	SO4	F	301	-	4,4,4	0.23	0	6,6,6	0.22	0
2	SO4	G	301	-	4,4,4	0.58	0	6,6,6	0.28	0
2	SO4	H	301	-	4,4,4	0.62	0	6,6,6	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	301	-	-	0/0/0/0	0/0/0/0
2	SO4	B	301	-	-	0/0/0/0	0/0/0/0
2	SO4	C	301	-	-	0/0/0/0	0/0/0/0
2	SO4	D	301	-	-	0/0/0/0	0/0/0/0
2	SO4	E	301	-	-	0/0/0/0	0/0/0/0
2	SO4	F	301	-	-	0/0/0/0	0/0/0/0
2	SO4	G	301	-	-	0/0/0/0	0/0/0/0
2	SO4	H	301	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	C	301	SO4	O2-S-O1	-3.35	98.88	109.50
2	D	301	SO4	O4-S-O3	2.04	117.29	108.98
2	D	301	SO4	O2-S-O1	3.30	119.94	109.50

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	301	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/263 (96%)	-0.12	3 (1%) 81 85	3, 11, 30, 65	0
1	B	256/263 (97%)	-0.18	0 100 100	3, 11, 28, 48	0
1	C	258/263 (98%)	-0.12	2 (0%) 87 91	3, 14, 31, 44	0
1	D	256/263 (97%)	-0.13	1 (0%) 93 95	3, 13, 30, 55	0
1	E	252/263 (95%)	0.34	6 (2%) 62 71	15, 28, 47, 55	0
1	F	254/263 (96%)	0.27	1 (0%) 93 95	15, 28, 43, 55	0
1	G	252/263 (95%)	0.35	8 (3%) 51 60	17, 28, 44, 71	0
1	H	253/263 (96%)	0.33	4 (1%) 74 80	14, 28, 46, 61	0
All	All	2036/2104 (96%)	0.09	25 (1%) 81 85	3, 22, 42, 71	0

The worst 5 of 25 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	166	TYR	5.8
1	C	100	ASN	3.9
1	A	241	ARG	3.3
1	H	52	ASN	3.1
1	E	53	SER	2.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	SO4	F	301	5/5	0.97	0.13	-0.59	36,37,38,41	0
2	SO4	E	301	5/5	0.97	0.12	-0.66	41,42,43,43	0
2	SO4	D	301	5/5	0.98	0.09	-1.31	11,11,11,12	0
2	SO4	B	301	5/5	0.97	0.08	-1.86	9,10,10,12	0
2	SO4	G	301	5/5	0.99	0.07	-1.94	23,24,26,27	0
2	SO4	A	301	5/5	0.98	0.07	-1.99	16,18,19,22	0
2	SO4	C	301	5/5	0.98	0.08	-3.07	12,13,14,14	0
2	SO4	H	301	5/5	0.98	0.07	-3.44	15,16,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.