



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:43 PM GMT

PDB ID : 3RZE  
Title : Structure of the human histamine H1 receptor in complex with doxepin  
Authors : Shimamura, T.; Han, G.W.; Shiroishi, M.; Weyand, S.; Tsujimoto, H.; Winter, G.; Katritch, V.; Abagyan, R.; Cherezov, V.; Liu, W.; Kobayashi, T.; Stevens, R.; Iwata, S.; GPCR Network (GPCR); Joint Center for Innovative Membrane Protein Technologies (JCIMPT)  
Deposited on : 2011-05-11  
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	452	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PO4	A	1202	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	OLC	A	1205	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3555 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histamine H1 receptor, Lysozyme chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	428	3481	2279	591	586	25	0	0	0

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1054	THR	CYS	ENGINEERED MUTATION	UNP P00720
A	1097	ALA	CYS	ENGINEERED MUTATION	UNP P00720
A	488	GLY	-	EXPRESSION TAG	UNP P35367
A	489	GLU	-	EXPRESSION TAG	UNP P35367
A	490	ASN	-	EXPRESSION TAG	UNP P35367
A	491	LEU	-	EXPRESSION TAG	UNP P35367
A	492	TYR	-	EXPRESSION TAG	UNP P35367
A	493	PHE	-	EXPRESSION TAG	UNP P35367
A	494	GLN	-	EXPRESSION TAG	UNP P35367

- Molecule 2 is (3E)-3-(DIBENZO[B,E]OXEPIN-11(6H)-YLIDENE)-N,N-DIMETHYLPROPAN-1-AMINE (three-letter code: 5EH) (formula: C<sub>19</sub>H<sub>21</sub>NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	21	19	1	1	0	1

- Molecule 3 is (3Z)-3-(DIBENZO[B,E]OXEPIN-11(6H)-YLIDENE)-N,N-DIMETHYLPROPAN-1-AMINE (three-letter code: D7V) (formula: C<sub>19</sub>H<sub>21</sub>NO).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	21	19	1	1	0	1

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0
4	A	1	Total O P 5 4 1	0	0

- Molecule 5 is (2R)-2,3-DIHYDROXYPROPYL (9Z)-OCTADEC-9-ENOATE (three-letter code: OLC) (formula: C<sub>21</sub>H<sub>40</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			15	11	4		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	O	0	0
			2	2		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histamine H1 receptor, Lysozyme chimera



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.14Å 88.14Å 331.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.50 – 3.10 34.52 – 3.10	Depositor EDS
% Data completeness (in resolution range)	(Not available) (34.50-3.10) 96.5 (34.52-3.10)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.07 (at 3.12Å)	Xtrriage
Refinement program	BUSTER 2.8.0	Depositor
R, $R_{free}$	0.214 , 0.249 0.230 , 0.278	Depositor DCC
$R_{free}$ test set	797 reflections (7.12%)	DCC
Wilson B-factor (Å <sup>2</sup> )	52.2	Xtrriage
Anisotropy	0.138	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 77.3	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Outliers	0 of 11996 reflections	Xtrriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3555	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.79% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, OLC, D7V, 5EH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.49	0/3566	0.72	1/4835 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	1112	ALA	CB-CA-C	-5.52	101.82	110.10

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3481	0	3566	67	0
2	A	21	0	21	3	0
3	A	21	0	21	2	0
4	A	15	0	0	1	0
5	A	15	0	19	0	0
6	A	2	0	0	0	0
All	All	3555	0	3627	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:429:ILE:HG22	1:A:430:PRO:CD	1.80	1.11
1:A:429:ILE:HG22	1:A:430:PRO:HD3	1.14	1.07
1:A:442:LYS:O	1:A:443:ASN:HB2	1.52	1.07
1:A:145:SER:HA	1:A:148:ILE:HG12	1.55	0.86
1:A:429:ILE:CG2	1:A:430:PRO:CD	2.60	0.77
1:A:1081:ASN:HB3	1:A:1084:LEU:HB2	1.69	0.74
1:A:429:ILE:CG2	1:A:430:PRO:N	2.52	0.72
1:A:429:ILE:CG2	1:A:430:PRO:HD3	2.07	0.72
1:A:219:GLN:C	1:A:219:GLN:OE1	2.30	0.70
1:A:220:HIS:O	1:A:221:CYS:O	2.11	0.68
1:A:425:ILE:CG2	1:A:429:ILE:HD13	2.23	0.68
1:A:219:GLN:OE1	1:A:219:GLN:CA	2.42	0.67
1:A:425:ILE:O	1:A:429:ILE:HB	1.95	0.66
1:A:442:LYS:O	1:A:443:ASN:CB	2.36	0.64
1:A:145:SER:HA	1:A:148:ILE:CG1	2.29	0.63
1:A:136:LEU:O	1:A:137:LYS:HB2	1.98	0.63
1:A:1094:VAL:HG21	1:A:1156:GLY:C	2.19	0.61
1:A:220:HIS:NE2	1:A:411:ARG:NH1	2.48	0.61
1:A:441:CYS:O	1:A:442:LYS:CB	2.50	0.59
1:A:196:ILE:HA	1:A:200:TYR:HB2	1.85	0.59
1:A:425:ILE:HG23	1:A:429:ILE:HD13	1.85	0.58
1:A:429:ILE:HG23	1:A:430:PRO:N	2.20	0.57
1:A:185:TYR:HA	1:A:191:LYS:HE2	1.88	0.56
1:A:148:ILE:O	1:A:152:TRP:HD1	1.90	0.55
1:A:1143:PRO:HD2	4:A:1204:PO4:O4	2.06	0.55
1:A:175:ARG:N	1:A:185:TYR:HH	2.05	0.54
1:A:219:GLN:OE1	1:A:219:GLN:HA	2.09	0.53
1:A:127:ARG:HH22	1:A:139:ARG:HD3	1.73	0.53
1:A:1053:ASN:C	1:A:1053:ASN:HD22	2.12	0.53
1:A:1111:VAL:O	1:A:1114:PHE:HB2	2.08	0.53
1:A:425:ILE:HG23	1:A:429:ILE:CD1	2.38	0.53
1:A:65:TYR:HD1	1:A:120:ILE:HG23	1.75	0.51
1:A:1050:ILE:HG22	1:A:1052:ARG:HG2	1.92	0.51
1:A:1027:ILE:HG21	1:A:1046:LEU:HD13	1.94	0.49
1:A:219:GLN:OE1	1:A:219:GLN:O	2.30	0.49
1:A:431:TYR:HE2	2:A:1200[A]:5EH:H09	1.77	0.49
1:A:1083:LYS:HD3	1:A:1112:ALA:O	2.13	0.48
1:A:61:VAL:HG11	1:A:139:ARG:HG3	1.95	0.48
1:A:429:ILE:N	1:A:430:PRO:HD2	2.29	0.48
1:A:428:TRP:HH2	2:A:1200[A]:5EH:H01	1.79	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1091:LEU:HD22	1:A:1095:ARG:HB3	1.97	0.47
1:A:425:ILE:HG22	1:A:429:ILE:HD13	1.97	0.47
1:A:220:HIS:CD2	1:A:411:ARG:NH1	2.84	0.46
1:A:428:TRP:CZ3	2:A:1200[A]:5EH:H20	2.51	0.46
1:A:36:THR:HA	1:A:39:LEU:HD12	1.96	0.46
1:A:87:TYR:CZ	1:A:93:TRP:HB3	2.52	0.45
1:A:134:ARG:C	1:A:136:LEU:H	2.20	0.45
1:A:220:HIS:N	1:A:220:HIS:CD2	2.83	0.44
1:A:218:ARG:O	1:A:218:ARG:HG2	2.16	0.44
1:A:1016:LYS:HA	1:A:1056:GLY:O	2.16	0.44
1:A:1059:THR:OG1	1:A:1062:GLU:HG3	2.18	0.44
1:A:1094:VAL:CG2	1:A:1095:ARG:N	2.81	0.43
1:A:82:PRO:HA	1:A:85:ILE:HD12	2.01	0.42
1:A:435:PHE:CZ	3:A:1201[B]:D7V:H18	2.55	0.42
1:A:182:THR:HG22	1:A:185:TYR:HB3	2.01	0.42
1:A:65:TYR:CD1	1:A:120:ILE:HG23	2.55	0.42
1:A:456:LEU:HA	1:A:459:ILE:HD12	2.02	0.42
1:A:127:ARG:HH11	1:A:134:ARG:HG3	1.84	0.42
1:A:422:ALA:O	1:A:426:LEU:HB2	2.20	0.42
1:A:1094:VAL:HG21	1:A:1156:GLY:O	2.19	0.41
1:A:431:TYR:OH	3:A:1201[B]:D7V:H3	2.20	0.41
1:A:48:VAL:HG22	1:A:479:PHE:HZ	1.85	0.41
1:A:83:MET:HA	1:A:86:LEU:HD12	2.01	0.41
1:A:1086:PRO:HB2	1:A:1122:GLN:OE1	2.19	0.41
1:A:148:ILE:O	1:A:152:TRP:CD1	2.72	0.41
1:A:32:VAL:HA	1:A:35:SER:HB2	2.03	0.41
1:A:61:VAL:HG13	1:A:139:ARG:NH1	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	424/452 (94%)	401 (95%)	18 (4%)	5 (1%)	16 52

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	221	CYS
1	A	443	ASN
1	A	442	LYS
1	A	181	GLU
1	A	1143	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	375/399 (94%)	333 (89%)	42 (11%)	7 29

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	VAL
1	A	47	LEU
1	A	61	VAL
1	A	81	MET
1	A	89	LEU
1	A	100	CYS
1	A	125	ARG
1	A	127	ARG
1	A	136	LEU
1	A	137	LYS
1	A	138	TYR
1	A	149	LEU
1	A	154	LEU
1	A	165	TRP
1	A	175	ARG
1	A	178	ASP

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	181	GLU
1	A	186	ASP
1	A	204	LEU
1	A	205	LEU
1	A	218	ARG
1	A	219	GLN
1	A	221	CYS
1	A	1007	LEU
1	A	1013	LEU
1	A	1027	ILE
1	A	1053	ASN
1	A	1090	SER
1	A	1094	VAL
1	A	1140	ASN
1	A	1159	ASP
1	A	412	LYS
1	A	419	PHE
1	A	426	LEU
1	A	429	ILE
1	A	433	ILE
1	A	435	PHE
1	A	441	CYS
1	A	445	CYS
1	A	446	ASN
1	A	463	LEU
1	A	482	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	1053	ASN
1	A	1055	ASN
1	A	474	ASN

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	5EH	A	1200[A]	-	23,23,23	0.67	0	25,31,31	1.08	1 (4%)
3	D7V	A	1201[B]	-	23,23,23	0.71	0	25,31,31	1.10	1 (4%)
4	PO4	A	1202	-	4,4,4	1.43	0	6,6,6	0.28	0
4	PO4	A	1203	-	4,4,4	1.49	0	6,6,6	0.28	0
4	PO4	A	1204	-	4,4,4	1.62	1 (25%)	6,6,6	0.30	0
5	OLC	A	1205	-	14,14,24	0.64	0	15,15,25	1.01	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5EH	A	1200[A]	-	-	0/6/19/19	0/2/3/3
3	D7V	A	1201[B]	-	-	0/6/19/19	0/2/3/3
4	PO4	A	1202	-	-	0/0/0/0	0/0/0/0
4	PO4	A	1203	-	-	0/0/0/0	0/0/0/0
4	PO4	A	1204	-	-	0/0/0/0	0/0/0/0
5	OLC	A	1205	-	-	0/14/14/24	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1204	PO4	P-O1	2.01	1.61	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1205	OLC	C21-O20-C1	2.91	124.98	116.85
3	A	1201[B]	D7V	C13-O1-C14	3.83	125.35	115.95
2	A	1200[A]	5EH	C14-O15-C16	3.93	125.58	115.95

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1200[A]	5EH	3	0
3	A	1201[B]	D7V	2	0
4	A	1204	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	428/452 (94%)	0.08	24 (5%) <span style="border: 1px solid red; padding: 2px;">28</span> <span style="border: 1px solid red; padding: 2px;">11</span>	10, 93, 162, 179	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	138	TYR	6.4
1	A	483	LEU	5.2
1	A	137	LYS	4.9
1	A	482	ILE	4.6
1	A	479	PHE	3.9
1	A	439	ALA	3.8
1	A	54	SER	3.3
1	A	478	THR	3.2
1	A	473	GLU	3.1
1	A	167	HIS	3.1
1	A	57	LYS	2.9
1	A	99	LEU	2.8
1	A	134	ARG	2.8
1	A	477	LYS	2.7
1	A	218	ARG	2.7
1	A	139	ARG	2.6
1	A	94	SER	2.6
1	A	136	LEU	2.5
1	A	178	ASP	2.5
1	A	161	PRO	2.4
1	A	32	VAL	2.1
1	A	443	ASN	2.1
1	A	55	GLU	2.1
1	A	444	CYS	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	PO4	A	1202	5/5	0.60	0.55	2.96	172,178,178,178	0
5	OLC	A	1205	15/25	0.91	0.24	2.87	10,39,208,209	0
4	PO4	A	1204	5/5	0.94	0.21	1.18	69,74,75,76	0
3	D7V	A	1201[B]	21/21	0.90	0.29	0.41	50,105,151,159	21
2	5EH	A	1200[A]	21/21	0.90	0.28	0.28	23,123,153,165	21
4	PO4	A	1203	5/5	0.78	0.47	-	143,149,150,150	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.