



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:07 PM GMT

PDB ID : 1S47  
Title : Crystal structure analysis of the DNA quadruplex d(TGGGGT)S2  
Authors : Caceres, C.; Wright, G.; Gouyette, C.; Parkinson, G.; Subirana, J.A.  
Deposited on : 2004-01-15  
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*










The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4242 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	6	
1	B	6	
1	C	6	
1	D	6	
1	E	6	
1	F	6	
1	G	6	
1	H	6	
1	I	6	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	J	6	 83%17%
1	K	6	 17%83%
1	L	6	 33%67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NA	A	1515	-	-	-	X
2	NA	E	1505	-	-	-	X
2	NA	E	1507	-	-	-	X
2	NA	H	1506	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 1531 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	6	Total	C	N	O	P	0	0	0
			108	50	22	31	5			
1	B	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	C	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	D	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	E	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	F	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	G	6	Total	C	N	O	P	0	0	0
			108	50	22	31	5			
1	H	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	I	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	J	6	Total	C	N	O	P	0	0	0
			108	50	22	31	5			
1	K	6	Total	C	N	O	P	0	0	0
			125	60	24	36	5			
1	L	6	Total	C	N	O	P	0	0	0
			108	50	22	31	5			

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	1	Total	Na	0	0
			1	1		
2	A	2	Total	Na	0	0
			2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	K	1	Total 1	Na 1	0	0
2	F	1	Total 1	Na 1	0	0
2	E	2	Total 2	Na 2	0	0

- Molecule 3 is THALLIUM (I) ION (three-letter code: Tl) (formula: Tl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total 1	Tl 1	0	0
3	I	3	Total 3	Tl 3	0	0
3	D	1	Total 1	Tl 1	0	0
3	E	1	Total 1	Tl 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total 2	O 2	0	0
4	B	11	Total 11	O 11	0	0
4	C	6	Total 6	O 6	0	0
4	D	10	Total 10	O 10	0	0
4	E	6	Total 6	O 6	0	0
4	F	5	Total 5	O 5	0	0
4	G	5	Total 5	O 5	0	0
4	H	12	Total 12	O 12	0	0
4	I	12	Total 12	O 12	0	0
4	J	5	Total 5	O 5	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	K	6	Total	O	0	0
			6	6		
4	L	6	Total	O	0	0
			6	6		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'



- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'



- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'




- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'



- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'



- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'



- Molecule 1: 5'-D(\*TP\*GP\*GP\*GP\*GP\*T)-3'





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	28.32Å 56.34Å 51.75Å 90.00° 99.17° 90.00°	Depositor
Resolution (Å)	19.84 – 2.50 19.84 – 2.51	Depositor EDS
% Data completeness (in resolution range)	79.0 (19.84-2.50) 96.2 (19.84-2.51)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.50Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.224 , 0.311 0.246 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	DCC
Wilson B-factor (Å <sup>2</sup> )	21.3	Xtriage
Anisotropy	1.336	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 41.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	4 of 5472 reflections (0.073%)	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	1531	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.98 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.1103e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, TL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.39	0/121	0.65	0/186
1	B	0.42	0/140	0.69	0/216
1	C	0.44	0/140	0.68	0/216
1	D	0.49	0/140	0.61	0/216
1	E	0.41	0/140	0.68	0/216
1	F	0.43	0/140	0.75	0/216
1	G	0.42	0/121	0.63	0/186
1	H	0.50	0/140	0.70	0/216
1	I	0.47	0/140	0.77	0/216
1	J	0.46	0/121	0.58	0/186
1	K	0.47	0/140	0.62	0/216
1	L	0.47	0/121	0.71	0/186
All	All	0.45	0/1604	0.68	0/2472

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	108	0	57	5	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	125	0	70	2	0
1	C	125	0	70	1	0
1	D	125	0	70	1	0
1	E	125	0	70	7	0
1	F	125	0	70	7	0
1	G	108	0	57	8	0
1	H	125	0	70	5	0
1	I	125	0	70	8	0
1	J	108	0	57	2	0
1	K	125	0	70	14	0
1	L	108	0	57	5	0
2	A	2	0	0	0	0
2	E	2	0	0	0	0
2	F	1	0	0	0	0
2	H	1	0	0	0	0
2	K	1	0	0	0	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	I	3	0	0	0	0
4	A	2	0	0	1	0
4	B	11	0	0	0	0
4	C	6	0	0	0	0
4	D	10	0	0	0	0
4	E	6	0	0	0	0
4	F	5	0	0	0	0
4	G	5	0	0	0	0
4	H	12	0	0	1	0
4	I	12	0	0	1	0
4	J	5	0	0	0	0
4	K	6	0	0	0	0
4	L	6	0	0	0	0
All	All	1531	0	788	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:801:DT:H5''	1:K:1242:DG:H1'	1.44	0.97

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:911:DT:H2''	4:H:1990:HOH:O	1.69	0.93
1:K:1246:DT:H6	1:K:1246:DT:H5''	1.37	0.87
1:K:1245:DG:H4'	1:K:1246:DT:O5'	1.75	0.83
1:G:801:DT:H5''	1:K:1242:DG:C1'	2.10	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

### 5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 13 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	6/6 (100%)	1.33	1 (16%) 2 2	17, 19, 34, 39	0
1	B	6/6 (100%)	1.01	1 (16%) 2 2	14, 29, 30, 42	0
1	C	6/6 (100%)	0.93	0 100 100	13, 19, 24, 32	0
1	D	6/6 (100%)	1.67	1 (16%) 2 2	17, 20, 22, 37	0
1	E	6/6 (100%)	1.06	1 (16%) 2 2	13, 20, 26, 34	0
1	F	6/6 (100%)	0.90	1 (16%) 2 2	13, 19, 20, 36	0
1	G	6/6 (100%)	0.70	0 100 100	17, 22, 24, 34	0
1	H	6/6 (100%)	1.61	1 (16%) 2 2	13, 17, 32, 39	0
1	I	6/6 (100%)	0.76	1 (16%) 2 2	11, 13, 23, 31	0
1	J	6/6 (100%)	0.04	0 100 100	13, 17, 18, 18	0
1	K	6/6 (100%)	0.52	0 100 100	13, 16, 19, 27	0
1	L	6/6 (100%)	0.24	0 100 100	9, 13, 20, 23	0
All	All	72/72 (100%)	0.90	7 (9%) 10 10	9, 19, 36, 42	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	436	DT	5.7
1	H	911	DT	4.7
1	E	581	DT	3.0
1	A	106	DT	3.0
1	F	691	DT	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	NA	E	1505	1/1	0.96	0.50	22.26	3,3,3,3	0
2	NA	H	1506	1/1	0.90	0.48	20.64	3,3,3,3	0
2	NA	E	1507	1/1	0.87	0.27	4.98	11,11,11,11	0
2	NA	A	1515	1/1	0.85	0.39	3.86	24,24,24,24	0
2	NA	K	1512	1/1	0.95	0.29	0.95	4,4,4,4	0
2	NA	F	1510	1/1	0.98	0.14	-0.74	3,3,3,3	0
2	NA	A	1514	1/1	0.96	0.20	-0.96	4,4,4,4	0
3	TL	B	1703	1/1	0.99	0.08	-3.90	47,47,47,47	1
3	TL	I	1705	1/1	0.96	0.10	-4.75	25,25,25,25	1
3	TL	D	1704	1/1	0.96	0.09	-5.65	28,28,28,28	1
3	TL	I	1700	1/1	0.98	0.09	-5.91	55,55,55,55	1
3	TL	I	1702	1/1	0.98	0.08	-6.69	59,59,59,59	1
3	TL	E	1701	1/1	0.97	0.09	-9.71	57,57,57,57	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.