



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 10:09 PM GMT

PDB ID : 1S9Q  
Title : crystal structure of the ligand-binding domain of the estrogen-related receptor gamma in complex with 4-hydroxytamoxifen  
Authors : Greschik, H.; Flaig, R.; Renaud, J.P.; Moras, D.  
Deposited on : 2004-02-05  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	251	 4% 80% 9% 11%
1	B	251	 4% 73% 12% • 14%

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Estrogen-related receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	224	Total	C	N	O	S	0	0	0
			1743	1118	284	330	11			
1	B	216	Total	C	N	O	S	0	0	0
			1677	1076	272	317	12			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	208	MET	-	cloning artifacts	UNP P62509
A	209	GLY	-	cloning artifacts	UNP P62509
A	210	SER	-	cloning artifacts	UNP P62509
A	211	SER	-	cloning artifacts	UNP P62509
A	212	HIS	-	EXPRESSION TAG	UNP P62509
A	213	HIS	-	EXPRESSION TAG	UNP P62509
A	214	HIS	-	EXPRESSION TAG	UNP P62509
A	215	HIS	-	EXPRESSION TAG	UNP P62509
A	216	HIS	-	EXPRESSION TAG	UNP P62509
A	217	HIS	-	EXPRESSION TAG	UNP P62509
A	218	SER	-	cloning artifacts	UNP P62509
A	219	SER	-	cloning artifacts	UNP P62509
A	220	GLY	-	cloning artifacts	UNP P62509
A	221	LEU	-	cloning artifacts	UNP P62509
A	222	VAL	-	cloning artifacts	UNP P62509
A	223	PRO	-	cloning artifacts	UNP P62509
A	224	ARG	-	cloning artifacts	UNP P62509
A	225	GLY	-	cloning artifacts	UNP P62509
A	226	SER	-	cloning artifacts	UNP P62509
A	227	HIS	-	cloning artifacts	UNP P62509
A	228	MET	-	cloning artifacts	UNP P62509
B	208	MET	-	cloning artifacts	UNP P62509
B	209	GLY	-	cloning artifacts	UNP P62509
B	210	SER	-	cloning artifacts	UNP P62509
B	211	SER	-	cloning artifacts	UNP P62509

*Continued on next page...*

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	212	HIS	-	EXPRESSION TAG	UNP P62509
B	213	HIS	-	EXPRESSION TAG	UNP P62509
B	214	HIS	-	EXPRESSION TAG	UNP P62509
B	215	HIS	-	EXPRESSION TAG	UNP P62509
B	216	HIS	-	EXPRESSION TAG	UNP P62509
B	217	HIS	-	EXPRESSION TAG	UNP P62509
B	218	SER	-	cloning artifacts	UNP P62509
B	219	SER	-	cloning artifacts	UNP P62509
B	220	GLY	-	cloning artifacts	UNP P62509
B	221	LEU	-	cloning artifacts	UNP P62509
B	222	VAL	-	cloning artifacts	UNP P62509
B	223	PRO	-	cloning artifacts	UNP P62509
B	224	ARG	-	cloning artifacts	UNP P62509
B	225	GLY	-	cloning artifacts	UNP P62509
B	226	SER	-	cloning artifacts	UNP P62509
B	227	HIS	-	cloning artifacts	UNP P62509
B	228	MET	-	cloning artifacts	UNP P62509

- Molecule 2 is 4-HYDROXYTAMOXIFEN (three-letter code: OHT) (formula:  $C_{26}H_{29}NO_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			29	26	1	2		
2	B	1	Total	C	N	O	0	0
			29	26	1	2		

- Molecule 3 is CHOLIC ACID (three-letter code: CHD) (formula:  $C_{24}H_{40}O_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			29	24	5		
3	B	1	Total	C	O	0	0
			29	24	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	64	Total	O	0	0
			64	64		
4	B	60	Total	O	0	0
			60	60		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Estrogen-related receptor gamma



- Molecule 1: Estrogen-related receptor gamma



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	89.48 Å 89.48 Å 161.76 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.83 – 2.20 29.83 – 2.20	Depositor EDS
% Data completeness (in resolution range)	98.3 (29.83-2.20) 99.9 (29.83-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.08	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	6.55 (at 2.20 Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.220 , 0.244 0.222 , 0.245	Depositor DCC
$R_{free}$ test set	1728 reflections (5.07%)	DCC
Wilson B-factor (Å <sup>2</sup> )	41.8	Xtriage
Anisotropy	0.359	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 42.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 34096 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3660	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OHT, CHD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.34	0/1774	0.53	0/2405
1	B	0.32	0/1705	0.51	1/2311 (0.0%)
All	All	0.33	0/3479	0.52	1/4716 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	445	PRO	N-CA-CB	5.49	109.88	103.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1743	0	1739	17	0
1	B	1677	0	1669	24	0
2	A	29	0	28	2	0
2	B	29	0	28	2	0
3	A	29	0	38	1	0
3	B	29	0	38	0	0
4	A	64	0	0	0	0
4	B	60	0	0	2	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3660	0	3540	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:304:ALA:HB2	1:B:377:SER:HB3	1.72	0.70
1:B:246:PRO:HB3	1:B:278:VAL:CG1	2.25	0.67
1:B:431:ALA:O	2:B:459:OHT:H15	1.97	0.65
1:B:407:HIS:HA	1:B:409:GLU:OE2	1.99	0.63
1:B:400:GLN:HG3	4:B:103:HOH:O	2.01	0.61
1:B:246:PRO:HB3	1:B:278:VAL:HG11	1.84	0.60
1:A:433:GLN:HE22	1:B:433:GLN:HE22	1.51	0.59
1:A:304:ALA:HB2	1:A:377:SER:HB3	1.85	0.58
1:B:449:LEU:O	1:B:453:MET:HG3	2.09	0.52
1:A:447:HIS:HB3	1:A:450:PHE:HB2	1.93	0.51
1:A:429:THR:O	1:A:433:GLN:HG3	2.11	0.50
1:A:431:ALA:O	2:A:200:OHT:H15	2.12	0.50
1:B:403:GLU:HG3	1:B:414:ALA:HB2	1.94	0.49
1:A:435:PHE:CZ	1:A:439:LYS:HD2	2.47	0.49
1:A:400:GLN:HG3	1:B:412:ARG:HD2	1.95	0.49
1:B:333:ASP:OD1	1:B:336:GLN:HG3	2.13	0.48
1:A:310:ILE:HD12	1:A:428:SER:HA	1.96	0.47
1:A:310:ILE:CD1	1:A:428:SER:HA	2.45	0.47
1:A:448:LYS:H	1:A:448:LYS:HG2	1.24	0.46
1:A:333:ASP:OD1	1:A:336:GLN:HG3	2.16	0.46
1:B:437:ASN:N	1:B:437:ASN:HD22	2.14	0.46
1:A:307:GLU:HA	1:A:428:SER:OG	2.18	0.44
1:B:249:ILE:HD11	1:B:278:VAL:HG21	1.99	0.44
1:B:400:GLN:CG	4:B:103:HOH:O	2.63	0.44
1:B:315:TYR:O	1:B:318:LEU:HG	2.18	0.44
1:A:353:VAL:HG12	1:A:357:LYS:HD2	1.99	0.42
1:A:380:MET:HG2	1:B:351:GLN:OE1	2.20	0.42
3:A:459:CHD:H182	3:A:459:CHD:H8	1.80	0.42
1:B:366:PHE:CZ	1:B:370:LYS:HE2	2.55	0.42
1:B:245:GLU:HA	1:B:245:GLU:OE2	2.19	0.42
1:B:234:TYR:HD1	1:B:239:SER:HG	1.64	0.42
1:A:326:TYR:OH	2:A:200:OHT:H101	2.19	0.41
1:B:407:HIS:CA	1:B:409:GLU:OE2	2.68	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:412:ARG:HB3	1:B:396:HIS:CE1	2.55	0.41
1:B:326:TYR:OH	2:B:459:OHT:H101	2.22	0.40
1:B:306:MET:O	1:B:310:ILE:HG12	2.21	0.40
1:A:449:LEU:O	1:A:453:MET:HG3	2.21	0.40
1:B:445:PRO:O	1:B:446:MET:C	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	220/251 (88%)	213 (97%)	7 (3%)	0	100	100
1	B	212/251 (84%)	209 (99%)	3 (1%)	0	100	100
All	All	432/502 (86%)	422 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/221 (85%)	187 (100%)	1 (0%)	92	96
1	B	180/221 (81%)	178 (99%)	2 (1%)	80	89
All	All	368/442 (83%)	365 (99%)	3 (1%)	86	93

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	448	LYS
1	B	400	GLN
1	B	409	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	336	GLN
1	A	400	GLN
1	A	433	GLN
1	A	437	ASN
1	B	336	GLN
1	B	396	HIS
1	B	433	GLN
1	B	437	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	OHT	A	200	-	31,31,31	3.47	9 (29%)	41,41,41	1.60	5 (12%)
3	CHD	A	459	-	29,32,32	2.91	11 (37%)	48,51,51	2.53	15 (31%)
2	OHT	B	459	-	31,31,31	3.44	11 (35%)	41,41,41	1.60	5 (12%)
3	CHD	B	500	-	29,32,32	2.91	11 (37%)	48,51,51	2.54	17 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OHT	A	200	-	-	0/24/24/24	0/3/3/3
3	CHD	A	459	-	-	0/7/74/74	0/4/4/4
2	OHT	B	459	-	-	0/24/24/24	0/3/3/3
3	CHD	B	500	-	-	0/7/74/74	0/4/4/4

All (42) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	459	CHD	C13-C14	-8.12	1.41	1.55
3	B	500	CHD	C13-C14	-8.03	1.41	1.55
3	B	500	CHD	C13-C12	-8.01	1.41	1.54
3	A	459	CHD	C13-C12	-7.82	1.42	1.54
3	A	459	CHD	C16-C17	-5.91	1.40	1.54
3	B	500	CHD	C16-C17	-5.88	1.40	1.54
3	A	459	CHD	O12-C12	-3.51	1.37	1.43
3	B	500	CHD	O12-C12	-3.30	1.38	1.43
3	B	500	CHD	C10-C9	-3.21	1.49	1.56
3	A	459	CHD	C10-C9	-3.14	1.50	1.56
3	B	500	CHD	C6-C5	2.02	1.57	1.53
2	B	459	OHT	C5-C4	2.11	1.43	1.38
3	A	459	CHD	C6-C5	2.13	1.57	1.53
2	B	459	OHT	C12-C11	2.13	1.42	1.39
2	B	459	OHT	C7-C8	2.19	1.40	1.34
3	A	459	CHD	O7-C7	2.22	1.48	1.43
3	A	459	CHD	C19-C10	2.22	1.58	1.54
3	B	500	CHD	O7-C7	2.23	1.48	1.43
3	B	500	CHD	C19-C10	2.24	1.58	1.54
2	A	200	OHT	C5-C4	2.27	1.43	1.38
2	A	200	OHT	C1-C7	2.43	1.53	1.49
2	B	459	OHT	C1-C7	2.72	1.53	1.49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	459	OHT	C16-C11	2.73	1.44	1.39
2	A	200	OHT	C16-C11	2.76	1.44	1.39
3	A	459	CHD	C18-C13	2.78	1.58	1.54
3	A	459	CHD	C11-C9	2.78	1.58	1.53
3	B	500	CHD	C11-C9	2.82	1.58	1.53
3	B	500	CHD	C18-C13	2.84	1.59	1.54
3	B	500	CHD	C8-C14	3.19	1.60	1.53
3	A	459	CHD	C8-C14	3.28	1.60	1.53
2	B	459	OHT	C2-C1	3.82	1.45	1.39
2	A	200	OHT	C2-C1	3.87	1.45	1.39
2	A	200	OHT	C13-C14	5.73	1.52	1.38
2	B	459	OHT	C13-C14	5.87	1.53	1.38
2	B	459	OHT	C15-C16	6.87	1.53	1.38
2	A	200	OHT	C14-C15	6.87	1.55	1.38
2	A	200	OHT	C15-C16	7.00	1.53	1.38
2	B	459	OHT	C14-C15	7.05	1.56	1.38
2	B	459	OHT	C3-C4	7.57	1.54	1.38
2	A	200	OHT	C3-C4	7.65	1.54	1.38
2	B	459	OHT	C18-C17	10.66	1.57	1.39
2	A	200	OHT	C18-C17	10.95	1.58	1.39

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	459	OHT	C19-C18-C17	-6.00	113.81	120.76
2	A	200	OHT	C19-C18-C17	-5.92	113.89	120.76
3	B	500	CHD	C9-C11-C12	-5.72	107.14	114.36
3	A	459	CHD	C9-C11-C12	-5.67	107.20	114.36
3	A	459	CHD	C18-C13-C12	-3.95	105.23	109.09
3	B	500	CHD	C18-C13-C12	-3.89	105.30	109.09
3	A	459	CHD	C18-C13-C14	-3.77	105.27	111.22
3	B	500	CHD	C18-C13-C14	-3.65	105.47	111.22
2	A	200	OHT	C2-C1-C7	-3.56	115.74	120.92
2	B	459	OHT	C2-C1-C7	-3.42	115.93	120.92
3	B	500	CHD	C6-C5-C10	-3.37	108.94	112.66
3	A	459	CHD	C19-C10-C1	-3.36	102.55	108.20
3	A	459	CHD	C6-C5-C10	-3.28	109.04	112.66
3	B	500	CHD	C19-C10-C1	-3.11	102.97	108.20
3	A	459	CHD	C16-C15-C14	-2.85	99.39	105.12
3	B	500	CHD	C16-C15-C14	-2.78	99.52	105.12
3	B	500	CHD	C11-C9-C8	-2.46	107.23	110.73
3	A	459	CHD	C11-C9-C8	-2.38	107.34	110.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	500	CHD	C9-C8-C7	-2.29	109.22	111.92
3	A	459	CHD	C6-C7-C8	-2.24	109.09	111.47
3	B	500	CHD	C6-C7-C8	-2.21	109.13	111.47
2	A	200	OHT	C13-C14-C15	-2.13	116.19	119.93
2	B	459	OHT	C13-C14-C15	-2.13	116.19	119.93
3	B	500	CHD	C15-C16-C17	2.06	109.28	105.12
3	B	500	CHD	C14-C8-C9	2.11	112.53	109.62
3	A	459	CHD	C14-C8-C9	2.12	112.53	109.62
3	B	500	CHD	C1-C10-C5	2.51	111.94	107.81
3	A	459	CHD	C1-C10-C5	2.57	112.03	107.81
3	B	500	CHD	C10-C9-C8	2.97	115.14	111.88
3	A	459	CHD	C10-C9-C8	2.98	115.15	111.88
2	A	200	OHT	C18-C19-C20	3.41	124.05	119.74
2	B	459	OHT	C13-C12-C11	3.44	124.66	120.33
2	A	200	OHT	C13-C12-C11	3.51	124.75	120.33
2	B	459	OHT	C18-C19-C20	3.57	124.24	119.74
3	A	459	CHD	C17-C13-C12	4.31	121.50	117.68
3	B	500	CHD	C17-C13-C12	4.55	121.72	117.68
3	B	500	CHD	C11-C12-C13	6.05	117.34	111.20
3	A	459	CHD	C11-C12-C13	6.21	117.51	111.20
3	A	459	CHD	O12-C12-C13	6.44	121.56	111.11
3	B	500	CHD	O12-C12-C13	6.48	121.61	111.11
3	A	459	CHD	C6-C5-C4	6.86	118.71	111.05
3	B	500	CHD	C6-C5-C4	6.99	118.85	111.05

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	200	OHT	2	0
3	A	459	CHD	1	0
2	B	459	OHT	2	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	224/251 (89%)	0.00	10 (4%) 37 36	28, 43, 66, 82	0
1	B	216/251 (86%)	0.15	11 (5%) 32 31	30, 47, 79, 89	0
All	All	440/502 (87%)	0.08	21 (4%) 34 34	28, 44, 74, 89	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	436	TYR	5.2
1	A	424	LEU	3.5
1	B	449	LEU	3.2
1	A	233	PRO	3.1
1	A	421	LEU	3.0
1	B	451	LEU	2.9
1	B	422	PRO	2.8
1	B	421	LEU	2.8
1	A	422	PRO	2.8
1	A	259	ASP	2.5
1	B	408	MET	2.3
1	A	451	LEU	2.3
1	B	423	LEU	2.2
1	B	445	PRO	2.2
1	A	308	ILE	2.2
1	B	450	PHE	2.2
1	A	455	GLU	2.2
1	B	438	ILE	2.1
1	B	234	TYR	2.1
1	A	304	ALA	2.1
1	A	456	ALA	2.1



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	OHT	A	200	29/29	0.94	0.19	1.55	36,40,59,61	0
2	OHT	B	459	29/29	0.90	0.19	1.20	46,50,65,67	0
3	CHD	B	500	29/29	0.81	0.27	0.80	58,60,71,73	0
3	CHD	A	459	29/29	0.86	0.14	0.12	44,48,61,62	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.