Full wwPDB NMR Structure Validation Report (i)

Apr 26, 2016 – 04:55 PM BST

PDB ID	:	1SB0
Title	:	Solution structure of the KIX domain of CBP bound to the transactivation domain of c-Myb
Authors	:	Zor, T.; De Guzman, R.N.; Dyson, H.J.; Wright, P.E.
Deposited on	:	2004-02-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 81%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	87	82% • • 14%
2	B	25	56% 8% 36%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:4-A:78, B:89-B:104 (91)	0.35	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 5 single-model clusters were found.

Cluster number	Models
1	1, 4, 6, 7, 9, 13, 14, 16, 18, 20
2	2, 3, 5, 15, 19
Single-model clusters	8; 10; 11; 12; 17

3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1894 atoms, of which 960 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called protein CBP.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	87	1464	459	735	136	132	2	0

- Molecule 2 is a protein called protein c-Myb.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	25	430	128	225	35	41	1	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: protein CBP

Chain A:

- Molecule 2: protein c-Myb

Chain B:



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: protein CBP

Chain A:

- Molecule 2: protein c-Myb

Chain B:

4.2.2 Score per residue for model 2

- Molecule 1: protein CBP

Chain A:

- Molecule 2: protein c-Myb

Chain B:

4.2.3 Score per residue for model 3

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.4 Score per residue for model 4 (medoid)

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.5 Score per residue for model 5

- Molecule 1: protein CBP

Chain A: 78% 7% • 14%

- Molecule 2: protein c-Myb

Chain B: 56% 8% 36%

4.2.6 Score per residue for model 6

- Molecule 1: protein CBP

Chain A: 78% 7% • 14%



- Molecule 2: protein c-Myb

Chain B: 60% • 36%

4.2.7 Score per residue for model 7

- Molecule 1: protein CBP

Chain A: 77% 8% • 14%

- Molecule 2: protein c-Myb

Chain B: 56% 8% 36%



4.2.8 Score per residue for model 8

- Molecule 1: protein CBP

Chain A: 82% 5% 14%



- Molecule 2: protein c-Myb

Chain B: 48% 16% 36%

4.2.9 Score per residue for model 9

- Molecule 1: protein CBP

Chain A: 80% 5% • 14%

- Molecule 2: protein c-Myb

Chain B: 56% 8% 36%

4.2.10 Score per residue for model 10

- Molecule 1: protein CBP

Chain A: 79% 6% • 14%



- Molecule 2: protein c-Myb

Chain B: 56% 8% 36%

4.2.11 Score per residue for model 11

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.12 Score per residue for model 12

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.13 Score per residue for model 13

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.14 Score per residue for model 14

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.15 Score per residue for model 15

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.16 Score per residue for model 16

- Molecule 1: protein CBP

Chain A:



- Molecule 2: protein c-Myb

Chain B:

4.2.17 Score per residue for model 17

- Molecule 1: protein CBP

Chain A: 76% 9% • 14%

- Molecule 2: protein c-Myb

Chain B: 60% • 36%

4.2.18 Score per residue for model 18

- Molecule 1: protein CBP

Chain A: 77% 8% • 14%



- Molecule 2: protein c-Myb

Chain B: 60% • 36%

4.2.19 Score per residue for model 19

- Molecule 1: protein CBP

Chain A: 82% • • 14%

- Molecule 2: protein c-Myb

Chain B: 64% 36%



4.2.20 Score per residue for model 20

- Molecule 1: protein CBP

Chain A:

- Molecule 2: protein c-Myb

Chain B:

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *initial structures from DYANA and refined in AMBER.*

Of the 200 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	**Classification**	**Version**
DYANA	structure solution	1.5
Amber	refinement	7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6095
Number of chemical shift lists	2
Total number of shifts	1266
Number of shifts mapped to atoms	1266
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	81%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.55±0.00	0±0/640 (0.0±0.0%)	0.87±0.02	0±0/864 (0.0±0.0%)
2	B	0.43±0.01	0±0/134 (0.0±0.0%)	0.80±0.06	0±0/178 (0.0±0.0%)
All	All	0.53	0/15480 (0.0%)	0.86	4/20840 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	2.6±1.0
2	B	0.0±0.0	0.1±0.2
All	All	0	54

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	73	TYR	CB-CG-CD1	-6.77	116.94	121.00	1	1
1	A	73	TYR	CB-CG-CD2	-5.58	117.65	121.00	14	2
1	A	61	ARG	NE-CZ-NH1	5.28	122.94	120.30	18	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	73	TYR	Sidechain	19
1	A	64	TYR	Sidechain	15
1	A	55	TYR	Sidechain	6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group	Models (Total)
1	A	65	TYR	Sidechain	5
1	A	27	PHE	Sidechain	3
2	B	91	ARG	Sidechain	1
1	A	38	ARG	Sidechain	1
1	A	39	ARG	Sidechain	1
1	A	46	TYR	Sidechain	1
1	A	61	ARG	Sidechain	1
1	A	15	ARG	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	624	617	617	0±0
2	B	134	145	145	0±0
All	All	15160	15240	15240	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:17:HIS:CD2	2:B:102:THR:HG23	0.45	2.46	11	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	75/87 (86%)	71±1 (95±2%)	4±1 (5±2%)	0±0 (0±0%)	59 88

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	B	16/25 (64%)	15±1 (96±5%)	0±0 (2±3%)	0±1 (2±3%)	14 56
All	All	1820/2240 (81%)	1725 (95%)	87 (5%)	8 (0%)	43 81

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	89	GLU	4
2	B	102	THR	2
1	A	28	PRO	1
1	A	60	SER	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	66/77 (86%)	62±2 (94±3%)	4±2 (6±3%)	27 72
2	B	16/23 (70%)	15±1 (91±5%)	2±1 (9±5%)	15 60
All	All	1640/2000 (82%)	1527 (93%)	113 (7%)	24 69

All 22 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	67	LEU	20
1	A	50	VAL	19
1	A	73	TYR	15
2	B	91	ARG	14
2	B	89	GLU	7
1	A	59	ASN	6
1	A	38	ARG	5
2	B	104	ASN	4
1	A	77	LYS	3
1	A	4	LYS	3
1	A	42	ASN	3
2	B	99	LEU	2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	12	GLN	2
1	A	20	HIS	2
1	A	39	ARG	1
1	A	21	LYS	1
1	A	36	LYS	1
1	A	8	GLU	1
2	B	92	ILE	1
2	B	93	LYS	1
2	B	90	LYS	1
1	A	40	MET	1

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 81% for the well-defined parts and 80% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 6095

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	963
Number of shifts mapped to atoms	963
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	86	1.72 ± 0.17	Should be applied
$^{13}C_{\beta}$	83	2.13 ± 0.07	Should be applied
$^{13}C'$	0	—	—
^{15}N	83	0.45 ± 0.22	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 67%, i.e. 802 atoms were assigned a chemical shift out of a possible 1203. 0 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	294/449 (65%)	147/179 (82%)	75/182 (41%)	72/88 (82%)
Sidechain	437/658 (66%)	272/387 (70%)	159/240 (66%)	6/31 (19%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	71/96 (74%)	36/51 (71%)	34/39 (87%)	1/6 (17%)
Overall	802/1203 (67%)	455/617 (74%)	268/461 (58%)	79/125 (63%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 62%, i.e. 934 atoms were assigned a chemical shift out of a possible 1507. 0 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	339/554 (61%)	170/221 (77%)	86/224 (38%)	83/109 (76%)
Sidechain	524/857 (61%)	327/505 (65%)	189/304 (62%)	8/48 (17%)
Aromatic	71/96 (74%)	36/51 (71%)	34/39 (87%)	1/6 (17%)
Overall	934/1507 (62%)	533/777 (69%)	309/567 (54%)	92/163 (56%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 6095

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	303
Number of shifts mapped to atoms	303
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	25	-0.39 ± 0.18	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	24	0.17 ± 0.18	None needed (< 0.5 ppm)
$^{13}C'$	0	—	—
^{15}N	23	-0.47 ± 0.65	None needed (< 0.5 ppm)

7.2.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 15%, i.e. 175 atoms were assigned a chemical shift out of a possible 1203. 4 out of 16 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	62/449 (14%)	31/179 (17%)	16/182 (9%)	15/88 (17%)
Sidechain	113/658 (17%)	67/387 (17%)	45/240 (19%)	1/31 (3%)
Aromatic	0/96 (0%)	0/51 (0%)	0/39 (0%)	0/6 (0%)
Overall	175/1203 (15%)	98/617 (16%)	61/461 (13%)	16/125 (13%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 18%, i.e. 266 atoms were assigned a chemical shift out of a possible 1507. 6 out of 21 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	95/554 (17%)	47/221 (21%)	25/224 (11%)	23/109 (21%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Sidechain	171/857 (20%)	100/505 (20%)	68/304 (22%)	3/48 (6%)
Aromatic	0/96 (0%)	0/51 (0%)	0/39 (0%)	0/6 (0%)
Overall	266/1507 (18%)	147/777 (19%)	93/567 (16%)	26/163 (16%)

7.2.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain B: