W O R L D W I D E
wwPDB
PROTEIN DATA BANK

wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 10:19 PM GMT

|		
PDB ID	:	1T2A
Title	:	Crystal structure of human GDP-D-mannose 4,6-dehydratase
Authors	:	Walker, J.R.; Vedadi, M.; Sharma, S.; Houston, S.; Wasney, G.; Loppnau, P.; Sundstrom, M.; Arrowsmith, C.; Edwards, A.; Oppermann, U.
Deposited on	:	2004-04-20
Resolution	:	1.84 Å(reported)

1   Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2634 (1.86-1.82)
Clashscore	102246	2862 (1.86-1.82)
Ramachandran outliers	100387	2831 (1.86-1.82)
Sidechain outliers	100360	2832 (1.86-1.82)
RSRZ outliers	91569	2639 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	2% / 76% / 12% • 10%
1	B	375	3% / 77% / 11% • 10%
1	C	375	6% / 74% / 15% • 10%
1	D	375	2% / 79% / 9% • 10%

2   Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 12166 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GDP-mannose 4,6 dehydratase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	338	Total	C	N	O	S	0	0	0
			2718	1735	467	506	10			
1	B	338	Total	C	N	O	S	0	0	0
			2718	1735	467	506	10			
1	C	338	Total	C	N	O	S	0	0	0
			2718	1735	467	506	10			
1	D	337	Total	C	N	O	S	0	0	0
			2710	1731	465	504	10			

There are 100 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	CLONING ARTIFACT	UNP O60547
A	1	GLY	-	CLONING ARTIFACT	UNP O60547
A	3	SER	-	CLONING ARTIFACT	UNP O60547
A	3	SER	-	CLONING ARTIFACT	UNP O60547
A	5	HIS	-	CLONING ARTIFACT	UNP O60547
A	6	HIS	-	CLONING ARTIFACT	UNP O60547
A	7	HIS	-	CLONING ARTIFACT	UNP O60547
A	8	HIS	-	CLONING ARTIFACT	UNP O60547
A	9	HIS	-	CLONING ARTIFACT	UNP O60547
A	9	HIS	-	CLONING ARTIFACT	UNP O60547
A	11	SER	-	CLONING ARTIFACT	UNP O60547
A	11	SER	-	CLONING ARTIFACT	UNP O60547
A	12	GLY	-	CLONING ARTIFACT	UNP O60547
A	13	ARG	-	CLONING ARTIFACT	UNP O60547
A	14	GLU	-	CLONING ARTIFACT	UNP O60547
A	15	ASN	-	CLONING ARTIFACT	UNP O60547
A	16	LYS	-	CLONING ARTIFACT	UNP O60547
A	17	TYR	-	CLONING ARTIFACT	UNP O60547
A	18	PHE	-	CLONING ARTIFACT	UNP O60547
A	19	GLN	-	CLONING ARTIFACT	UNP O60547
A	20	GLY	-	CLONING ARTIFACT	UNP O60547

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	21	HIS	-	CLONING ARTIFACT	UNP O60547
A	22	MET	-	CLONING ARTIFACT	UNP O60547
A	373	GLY	-	CLONING ARTIFACT	UNP O60547
A	374	SER	-	CLONING ARTIFACT	UNP O60547
B	0	MET	-	CLONING ARTIFACT	UNP O60547
B	1	GLY	-	CLONING ARTIFACT	UNP O60547
B	3	SER	-	CLONING ARTIFACT	UNP O60547
B	3	SER	-	CLONING ARTIFACT	UNP O60547
B	5	HIS	-	CLONING ARTIFACT	UNP O60547
B	6	HIS	-	CLONING ARTIFACT	UNP O60547
B	7	HIS	-	CLONING ARTIFACT	UNP O60547
B	8	HIS	-	CLONING ARTIFACT	UNP O60547
B	9	HIS	-	CLONING ARTIFACT	UNP O60547
B	9	HIS	-	CLONING ARTIFACT	UNP O60547
B	11	SER	-	CLONING ARTIFACT	UNP O60547
B	11	SER	-	CLONING ARTIFACT	UNP O60547
B	12	GLY	-	CLONING ARTIFACT	UNP O60547
B	13	ARG	-	CLONING ARTIFACT	UNP O60547
B	14	GLU	-	CLONING ARTIFACT	UNP O60547
B	15	ASN	-	CLONING ARTIFACT	UNP O60547
B	16	LYS	-	CLONING ARTIFACT	UNP O60547
B	17	TYR	-	CLONING ARTIFACT	UNP O60547
B	18	PHE	-	CLONING ARTIFACT	UNP O60547
B	19	GLN	-	CLONING ARTIFACT	UNP O60547
B	20	GLY	-	CLONING ARTIFACT	UNP O60547
B	21	HIS	-	CLONING ARTIFACT	UNP O60547
B	22	MET	-	CLONING ARTIFACT	UNP O60547
B	373	GLY	-	CLONING ARTIFACT	UNP O60547
B	374	SER	-	CLONING ARTIFACT	UNP O60547
C	0	MET	-	CLONING ARTIFACT	UNP O60547
C	1	GLY	-	CLONING ARTIFACT	UNP O60547
C	3	SER	-	CLONING ARTIFACT	UNP O60547
C	3	SER	-	CLONING ARTIFACT	UNP O60547
C	5	HIS	-	CLONING ARTIFACT	UNP O60547
C	6	HIS	-	CLONING ARTIFACT	UNP O60547
C	7	HIS	-	CLONING ARTIFACT	UNP O60547
C	8	HIS	-	CLONING ARTIFACT	UNP O60547
C	9	HIS	-	CLONING ARTIFACT	UNP O60547
C	9	HIS	-	CLONING ARTIFACT	UNP O60547
C	11	SER	-	CLONING ARTIFACT	UNP O60547
C	11	SER	-	CLONING ARTIFACT	UNP O60547
C	12	GLY	-	CLONING ARTIFACT	UNP O60547

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	13	ARG	-	CLONING ARTIFACT	UNP O60547
C	14	GLU	-	CLONING ARTIFACT	UNP O60547
C	15	ASN	-	CLONING ARTIFACT	UNP O60547
C	16	LYS	-	CLONING ARTIFACT	UNP O60547
C	17	TYR	-	CLONING ARTIFACT	UNP O60547
C	18	PHE	-	CLONING ARTIFACT	UNP O60547
C	19	GLN	-	CLONING ARTIFACT	UNP O60547
C	20	GLY	-	CLONING ARTIFACT	UNP O60547
C	21	HIS	-	CLONING ARTIFACT	UNP O60547
C	22	MET	-	CLONING ARTIFACT	UNP O60547
C	373	GLY	-	CLONING ARTIFACT	UNP O60547
C	374	SER	-	CLONING ARTIFACT	UNP O60547
D	0	MET	-	CLONING ARTIFACT	UNP O60547
D	1	GLY	-	CLONING ARTIFACT	UNP O60547
D	3	SER	-	CLONING ARTIFACT	UNP O60547
D	3	SER	-	CLONING ARTIFACT	UNP O60547
D	5	HIS	-	CLONING ARTIFACT	UNP O60547
D	6	HIS	-	CLONING ARTIFACT	UNP O60547
D	7	HIS	-	CLONING ARTIFACT	UNP O60547
D	8	HIS	-	CLONING ARTIFACT	UNP O60547
D	9	HIS	-	CLONING ARTIFACT	UNP O60547
D	9	HIS	-	CLONING ARTIFACT	UNP O60547
D	11	SER	-	CLONING ARTIFACT	UNP O60547
D	11	SER	-	CLONING ARTIFACT	UNP O60547
D	12	GLY	-	CLONING ARTIFACT	UNP O60547
D	13	ARG	-	CLONING ARTIFACT	UNP O60547
D	14	GLU	-	CLONING ARTIFACT	UNP O60547
D	15	ASN	-	CLONING ARTIFACT	UNP O60547
D	16	LYS	-	CLONING ARTIFACT	UNP O60547
D	17	TYR	-	CLONING ARTIFACT	UNP O60547
D	18	PHE	-	CLONING ARTIFACT	UNP O60547
D	19	GLN	-	CLONING ARTIFACT	UNP O60547
D	20	GLY	-	CLONING ARTIFACT	UNP O60547
D	21	HIS	-	CLONING ARTIFACT	UNP O60547
D	22	MET	-	CLONING ARTIFACT	UNP O60547
D	373	GLY	-	CLONING ARTIFACT	UNP O60547
D	374	SER	-	CLONING ARTIFACT	UNP O60547

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).

| Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 28	C 10	N 5	O 11	P 2	0	0
3	B	1	Total 28	C 10	N 5	O 11	P 2	0	0
3	C	1	Total 28	C 10	N 5	O 11	P 2	0	0
3	D	1	Total 28	C 10	N 5	O 11	P 2	0	0

- Molecule 4 is water.

| Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	285	Total 285	O 285	0	0
4	B	256	Total 256	O 256	0	0
4	C	199	Total 199	O 199	0	0
4	D	258	Total 258	O 258	0	0

3 Residue-property plots ⓘ

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GDP-mannose 4,6 dehydratase


- Molecule 1: GDP-mannose 4,6 dehydratase


- Molecule 1: GDP-mannose 4,6 dehydratase

● Molecule 1: GDP-mannose 4,6 dehydratase

Chain D:

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	88.59Å 122.06Å 139.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.93 – 1.84 35.75 – 1.84	Depositor EDS
% Data completeness (in resolution range)	98.2 (24.93-1.84) 98.3 (35.75-1.84)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.65 (at 1.84Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.173 , 0.212 0.173 , 0.213	Depositor DCC
R_{free} test set	6496 reflections (5.03%)	DCC
Wilson B-factor (Å^2)	20.1	Xtriage
Anisotropy	0.361	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.38 , 59.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< |L| > = 0.49, < L^2 > = 0.33$	Xtriage
Outliers	1 of 129273 reflections (0.001%)	Xtriage
F_o,F_c correlation	0.96	EDS
Total number of atoms	12166	wwPDB-VP
Average B, all atoms (Å^2)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

---

[1]Intensities estimated from amplitudes.
[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, NDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$|Z|$>5	RMSZ	#$|Z|$>5
1	A	1.16	3/2777 (0.1%)	1.21	11/3755 (0.3%)
1	B	1.09	3/2777 (0.1%)	1.20	6/3755 (0.2%)
1	C	0.97	2/2777 (0.1%)	0.90	2/3755 (0.1%)
1	D	1.15	4/2769 (0.1%)	0.96	4/3744 (0.1%)
All	All	1.10	12/11100 (0.1%)	1.08	23/15009 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	45	GLU	CD-OE1	6.46	1.32	1.25
1	D	357	GLU	CD-OE2	-6.34	1.18	1.25
1	A	260	TRP	CE3-CZ3	6.07	1.48	1.38
1	D	311	GLU	CG-CD	5.91	1.60	1.51
1	A	134	ARG	CD-NE	-5.89	1.36	1.46

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	247	ARG	NE-CZ-NH2	-32.81	103.90	120.30
1	A	134	ARG	NE-CZ-NH1	31.96	136.28	120.30
1	B	247	ARG	NE-CZ-NH1	30.76	135.68	120.30
1	A	134	ARG	NE-CZ-NH2	-29.80	105.40	120.30

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	247	ARG	CD-NE-CZ	10.43	138.21	123.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	231	TYR	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2718	0	2687	35	0
1	B	2718	0	2687	30	0
1	C	2718	0	2687	42	0
1	D	2710	0	2681	29	0
2	A	48	0	26	0	0
2	B	48	0	26	2	0
2	C	48	0	26	0	0
2	D	48	0	26	2	0
3	A	28	0	12	0	0
3	B	28	0	12	0	0
3	C	28	0	12	0	0
3	D	28	0	12	0	0
4	A	285	0	0	5	0
4	B	256	0	0	7	0
4	C	199	0	0	1	0
4	D	258	0	0	7	0
All	All	12166	0	10894	134	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 134 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:327:THR:HG21	1:D:192:ASN:HD21	1.29	0.95
1:D:145:ILE:HD12	1:D:145:ILE:H	1.40	0.87
1:A:244:ASP:OD1	1:A:282:ARG:NH1	2.07	0.87
1:D:167:GLN:HE22	1:D:333:GLN:H	1.21	0.85
1:C:27:LEU:HD11	1:C:53:ILE:HD11	1.58	0.84

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/375 (89%)	327 (98%)	5 (2%)	2 (1%)	30	14
1	B	334/375 (89%)	325 (97%)	8 (2%)	1 (0%)	46	29
1	C	334/375 (89%)	325 (97%)	8 (2%)	1 (0%)	46	29
1	D	333/375 (89%)	326 (98%)	7 (2%)	0	100	100
All	All	1335/1500 (89%)	1303 (98%)	28 (2%)	4 (0%)	46	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	368	THR
1	A	160	GLY
1	C	160	GLY
1	B	160	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	293/323 (91%)	282 (96%)	11 (4%)	40	19
1	B	293/323 (91%)	287 (98%)	6 (2%)	63	47
1	C	293/323 (91%)	281 (96%)	12 (4%)	37	16
1	D	292/323 (90%)	282 (97%)	10 (3%)	44	23
All	All	1171/1292 (91%)	1132 (97%)	39 (3%)	45	25

5 of 39 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	68	LEU
1	C	111	GLN
1	D	344	ASN
1	C	79	ASN
1	C	80	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	333	GLN
1	B	369	ASN
1	D	333	GLN
1	B	344	ASN
1	C	167	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#|Z| > 2$	Counts	RMSZ	$\#|Z| > 2$
2	NDP	A	1001	-	42,52,52	0.78	2 (4%)	55,80,80	1.01	4 (7%)
3	GDP	A	1002	-	23,30,30	1.04	1 (4%)	30,47,47	2.27	4 (13%)
2	NDP	B	1101	-	42,52,52	0.75	2 (4%)	55,80,80	1.01	4 (7%)
3	GDP	B	1102	-	23,30,30	1.04	1 (4%)	30,47,47	2.27	4 (13%)
2	NDP	C	1201	-	42,52,52	0.78	2 (4%)	55,80,80	1.01	4 (7%)
3	GDP	C	1202	-	23,30,30	1.03	1 (4%)	30,47,47	2.28	4 (13%)
2	NDP	D	1301	-	42,52,52	0.75	2 (4%)	55,80,80	1.01	4 (7%)
3	GDP	D	1302	-	23,30,30	1.05	1 (4%)	30,47,47	2.28	4 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	1001	-	-	0/30/77/77	0/5/5/5
3	GDP	A	1002	-	-	0/12/32/32	0/3/3/3
2	NDP	B	1101	-	-	0/30/77/77	0/5/5/5
3	GDP	B	1102	-	-	0/12/32/32	0/3/3/3
2	NDP	C	1201	-	-	0/30/77/77	0/5/5/5
3	GDP	C	1202	-	-	0/12/32/32	0/3/3/3
2	NDP	D	1301	-	-	0/30/77/77	0/5/5/5
3	GDP	D	1302	-	-	0/12/32/32	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1001	NDP	P2B-O2B	-2.51	1.52	1.60
2	C	1201	NDP	P2B-O2B	-2.47	1.52	1.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	NDP	P2B-O2B	-2.30	1.53	1.60
2	D	1301	NDP	P2B-O2B	-2.27	1.53	1.60
2	D	1301	NDP	C4N-C5N	2.11	1.53	1.49

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1302	GDP	C5-C6-N1	-8.82	111.53	123.59
3	B	1102	GDP	C5-C6-N1	-8.79	111.57	123.59
3	C	1202	GDP	C5-C6-N1	-8.75	111.63	123.59
3	A	1002	GDP	C5-C6-N1	-8.74	111.63	123.59
2	D	1301	NDP	O2B-P2B-O1X	-2.91	99.83	107.11

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1101	NDP	2	0
2	D	1301	NDP	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	338/375 (90%)	-0.28	6 (1%)	71	69	7, 18, 35, 62	1 (0%)
1	B	338/375 (90%)	-0.12	12 (3%)	46	42	7, 21, 43, 64	1 (0%)
1	C	338/375 (90%)	0.20	24 (7%)	19	17	9, 26, 47, 63	1 (0%)
1	D	337/375 (89%)	-0.10	7 (2%)	67	64	11, 21, 37, 55	1 (0%)
All	All	1351/1500 (90%)	-0.08	49 (3%)	46	42	7, 21, 43, 64	4 (0%)

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	69	TYR	8.5
1	D	188	TRP	8.4
1	B	188	TRP	8.1
1	C	188	TRP	7.9
1	A	368	THR	7.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
2	NDP	D	1301	48/48	0.94	0.16	1.64	11,13,17,17	0
2	NDP	A	1001	48/48	0.95	0.12	1.46	10,13,17,17	0
3	GDP	B	1102	28/28	0.92	0.12	1.37	16,18,23,25	0
2	NDP	B	1101	48/48	0.92	0.12	1.34	11,13,17,17	0
3	GDP	A	1002	28/28	0.94	0.10	0.99	14,15,21,23	0
3	GDP	D	1302	28/28	0.97	0.09	-0.11	15,17,22,24	0
2	NDP	C	1201	48/48	0.96	0.10	-0.44	10,13,17,17	0
3	GDP	C	1202	28/28	0.95	0.09	-0.65	14,15,21,23	0

6.5 Other polymers (i)

There are no such residues in this entry.