



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:09 PM GMT

PDB ID : 3T3O
Title : Molecular basis for the recognition and cleavage of RNA (CUGG) by the bi-functional 5'-3' exo/endoribonuclease RNase J
Authors : Dorleans, A.; Li de la Sierra-Gallay, I.; Piton, J.; Zig, L.; Gilet, L.; Putzer, H.; Condon, C.
Deposited on : 2011-07-25
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)
RNA backbone	2183	1172 (3.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	562	<div> <div>11%</div> <div>81%</div> <div>16%</div> <div>..</div> </div>
2	B	5	<div> <div>20%</div> <div>20%</div> <div>40%</div> <div>40%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4545 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Metal dependent hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	553	Total	C	N	O	S	0	0	0
			4330	2778	761	777	14			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	EXPRESSION TAG	UNP Q72JJ7
A	-6	SER	-	EXPRESSION TAG	UNP Q72JJ7
A	-5	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-4	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-3	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-2	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	-1	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	0	HIS	-	EXPRESSION TAG	UNP Q72JJ7
A	1	SER	-	EXPRESSION TAG	UNP Q72JJ7
A	77	ALA	HIS	ENGINEERED MUTATION	UNP Q72JJ7

- Molecule 2 is a RNA chain called O2'methyl-RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	5	Total	C	N	O	P	0	0	0
			111	51	17	38	5			

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	96	Total	O	0	0
			96	96		
5	B	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Metal dependent hydrolase



- Molecule 2: O2'methyl-RNA



4 Data and refinement statistics

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	99.35Å 117.94Å 229.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.76 – 2.50 26.95 – 2.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (27.76-2.50) 98.3 (26.95-2.50)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 2.50Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.227 , 0.285 0.237 , 0.299	Depositor DCC
R_{free} test set	1182 reflections (5.40%)	DCC
Wilson B-factor (Å ²)	52.5	Xtriage
Anisotropy	0.690	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 65.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	1 of 23081 reflections (0.004%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4545	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OMC, GOL, ZN, OMG, OMU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/4427	0.65	0/5993
2	B	1.12	0/22	1.63	1/34 (2.9%)
All	All	0.41	0/4449	0.66	1/6027 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	905	U	O4'-C1'-N1	5.88	112.90	108.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4330	0	4428	39	0
2	B	111	0	62	2	0
3	A	1	0	0	0	0
4	A	6	0	8	0	0
5	A	96	0	0	0	0
5	B	1	0	0	0	0
All	All	4545	0	4498	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:GLN:HE22	1:A:282:LYS:H	1.22	0.84
1:A:268:MET:HE3	1:A:304:LEU:HB3	1.75	0.68
1:A:54:LEU:HD23	1:A:447:VAL:HG21	1.80	0.63
1:A:487:VAL:HG11	1:A:509:ALA:HB3	1.88	0.55
1:A:241:ALA:HA	1:A:268:MET:HG3	1.88	0.55
1:A:150:HIS:CG	1:A:151:SER:H	2.25	0.55
1:A:319:LEU:HD21	1:A:327:MET:HB3	1.90	0.53
1:A:345:ASN:O	1:A:349:VAL:HG23	2.08	0.53
1:A:175:LEU:HB2	1:A:381:GLU:HG2	1.93	0.51
1:A:465:ASP:HB2	1:A:547:MET:CE	2.42	0.50
1:A:536:LYS:HE3	1:A:540:LYS:HE3	1.95	0.49
1:A:239:THR:HG22	1:A:338:SER:HB3	1.93	0.49
1:A:465:ASP:HB2	1:A:547:MET:HE2	1.95	0.49
1:A:290:LEU:HD11	1:A:327:MET:HG3	1.95	0.48
1:A:9:HIS:HB3	1:A:433:THR:HG22	1.95	0.48
1:A:408:LYS:HE2	1:A:421:THR:HB	1.95	0.48
1:A:13:ILE:HG12	1:A:429:VAL:HG22	1.97	0.46
1:A:254:ALA:HB2	1:A:302:LEU:HD13	1.97	0.46
1:A:335:VAL:HG21	1:A:356:LEU:HD22	1.98	0.46
1:A:293:VAL:HG22	1:A:301:VAL:HG22	1.97	0.45
1:A:336:ILE:HG12	1:A:364:LEU:HD12	1.99	0.45
1:A:377:ALA:HB1	1:A:381:GLU:HB2	1.98	0.44
1:A:240:PHE:CZ	2:B:901:OMC:H5"	2.52	0.44
1:A:23:LYS:HB2	1:A:56:PRO:HD3	2.00	0.44
1:A:200:LEU:HD11	1:A:395:LEU:HB2	2.00	0.44
2:B:903:OMG:H8	2:B:903:OMG:H5"	1.83	0.43
1:A:228:ILE:HG12	1:A:235:VAL:HG11	1.99	0.43
1:A:293:VAL:HA	1:A:296:LEU:HD12	2.00	0.43
1:A:480:LEU:HD12	1:A:486:VAL:HB	2.01	0.43
1:A:4:GLY:HA3	1:A:434:ARG:HH22	1.84	0.42
1:A:25:ILE:HG21	1:A:56:PRO:HG3	2.02	0.42
1:A:72:VAL:HG21	1:A:143:LEU:HD22	2.00	0.42
1:A:397:TRP:CG	1:A:398:HIS:N	2.88	0.42
1:A:244:ILE:HD11	1:A:271:PHE:HB3	2.02	0.41
1:A:311:GLN:HG2	1:A:311:GLN:H	1.55	0.41
1:A:319:LEU:HD11	1:A:327:MET:HE2	2.02	0.41
1:A:238:THR:HA	1:A:305:ALA:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44:PRO:HB2	1:A:45:GLU:H	1.68	0.41
1:A:194:GLY:HA2	1:A:198:VAL:HG23	2.04	0.40
1:A:33:GLU:HG2	1:A:69:LYS:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	551/562 (98%)	517 (94%)	26 (5%)	8 (2%)	13	22

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PRO
1	A	47	GLY
1	A	94	LYS
1	A	322	GLU
1	A	65	ARG
1	A	25	ILE
1	A	232	PRO
1	A	531	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	461/470 (98%)	433 (94%)	28 (6%)	23	42

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	32	ASP
1	A	35	PHE
1	A	59	ASP
1	A	65	ARG
1	A	133	ASP
1	A	144	ASP
1	A	151	SER
1	A	172	ASP
1	A	219	GLU
1	A	230	ARG
1	A	266	ARG
1	A	273	ARG
1	A	293	VAL
1	A	311	GLN
1	A	313	MET
1	A	337	LEU
1	A	370	LYS
1	A	384	LEU
1	A	395	LEU
1	A	407	PHE
1	A	438	GLU
1	A	463	LEU
1	A	500	ARG
1	A	513	LEU
1	A	514	LYS
1	A	523	LEU
1	A	549	LEU
1	A	553	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	64	HIS
1	A	136	GLN
1	A	155	ASN
1	A	248	GLN
1	A	379	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	406	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	B	4/5 (80%)	3 (75%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	B	902	OMU
2	B	903	OMG
2	B	905	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	OMC	B	901	2	18,23,23	1.54	3 (16%)	26,34,34	1.92	7 (26%)
2	OMU	B	902	2	12,22,23	1.64	2 (16%)	19,31,34	3.08	3 (15%)
2	OMG	B	903	2	17,26,27	1.50	3 (17%)	21,38,41	2.31	3 (14%)
2	OMG	B	904	2	17,26,27	1.54	3 (17%)	21,38,41	2.30	4 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OMC	B	901	2	-	0/8/28/28	0/2/2/2
2	OMU	B	902	2	-	0/5/27/28	0/2/2/2
2	OMG	B	903	2	-	0/5/27/28	0/3/3/3
2	OMG	B	904	2	-	0/5/27/28	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	901	OMC	P-OP3	-4.11	1.39	1.54
2	B	903	OMG	C8-N7	-2.40	1.30	1.34
2	B	904	OMG	C8-N7	-2.31	1.30	1.34
2	B	902	OMU	C4-N3	2.75	1.38	1.33
2	B	901	OMC	C6-N1	2.82	1.39	1.35
2	B	901	OMC	O2'-CM2	3.02	1.53	1.42
2	B	903	OMG	C2-N1	3.02	1.40	1.35
2	B	904	OMG	C2-N1	3.22	1.41	1.35
2	B	903	OMG	C6-N1	4.11	1.40	1.33
2	B	902	OMU	O2'-CM2	4.14	1.57	1.42
2	B	904	OMG	C6-N1	4.37	1.41	1.33

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	904	OMG	C5-C6-N1	-7.61	113.18	123.59
2	B	903	OMG	C5-C6-N1	-7.57	113.23	123.59
2	B	901	OMC	CM2-O2'-C2'	-6.25	96.93	114.59
2	B	902	OMU	C5-C4-N3	-3.40	114.39	123.12
2	B	904	OMG	N3-C2-N1	-2.71	123.32	127.44
2	B	901	OMC	OP3-P-OP1	-2.68	101.95	110.58
2	B	901	OMC	C6-N1-C2	-2.62	117.03	121.28
2	B	903	OMG	N3-C2-N1	-2.60	123.48	127.44
2	B	904	OMG	CM2-O2'-C2'	-2.51	107.50	114.59
2	B	902	OMU	C2'-C1'-N1	2.08	119.50	113.53
2	B	901	OMC	OP2-P-OP1	2.46	118.49	110.58
2	B	901	OMC	OP3-P-O5'	2.75	114.48	106.56
2	B	901	OMC	C2'-C1'-N1	3.24	122.86	113.53
2	B	901	OMC	C2-N3-C4	3.41	120.42	115.61
2	B	903	OMG	C6-N1-C2	5.64	123.77	115.94
2	B	904	OMG	C6-N1-C2	5.71	123.87	115.94
2	B	902	OMU	C4-N3-C2	12.42	126.44	114.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	901	OMC	1	0
2	B	903	OMG	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	701	-	5,5,5	0.44	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	701	-	-	0/4/4/4	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	553/562 (98%)	0.59	63 (11%) 7 6	37, 65, 100, 116	0
2	B	1/5 (20%)	4.41	1 (100%) 0 0	114, 114, 114, 114	0
All	All	554/567 (97%)	0.60	64 (11%) 6 6	37, 65, 101, 116	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	GLY	6.0
1	A	47	GLY	5.4
1	A	325	ALA	5.4
1	A	48	MET	4.9
1	A	327	MET	4.8
1	A	3	GLN	4.7
2	B	905	U	4.4
1	A	239	THR	4.3
1	A	237	VAL	4.3
1	A	515	ASN	3.9
1	A	322	GLU	3.6
1	A	238	THR	3.6
1	A	524	GLU	3.4
1	A	295	ASP	3.4
1	A	305	ALA	3.2
1	A	532	TYR	3.2
1	A	2	SER	3.1
1	A	336	ILE	3.1
1	A	326	LYS	3.1
1	A	456	GLY	3.1
1	A	523	LEU	3.0
1	A	302	LEU	2.9
1	A	303	ILE	2.9
1	A	474	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	202	ILE	2.9
1	A	313	MET	2.8
1	A	525	ARG	2.8
1	A	121	PRO	2.8
1	A	37	LEU	2.8
1	A	519	GLU	2.7
1	A	236	PHE	2.7
1	A	118	GLY	2.7
1	A	511	GLU	2.6
1	A	344	GLY	2.6
1	A	169	HIS	2.6
1	A	168	VAL	2.5
1	A	171	GLY	2.5
1	A	531	TYR	2.5
1	A	536	LYS	2.5
1	A	50	GLY	2.5
1	A	201	LEU	2.4
1	A	304	LEU	2.4
1	A	268	MET	2.4
1	A	323	GLY	2.4
1	A	312	PRO	2.3
1	A	497	ALA	2.3
1	A	518	ARG	2.3
1	A	482	GLY	2.3
1	A	291	GLU	2.2
1	A	28	PHE	2.2
1	A	334	THR	2.2
1	A	520	LYS	2.2
1	A	39	GLY	2.1
1	A	500	ARG	2.1
1	A	6	PRO	2.1
1	A	123	ALA	2.1
1	A	434	ARG	2.1
1	A	463	LEU	2.1
1	A	41	LEU	2.1
1	A	367	PRO	2.1
1	A	54	LEU	2.1
1	A	117	PHE	2.0
1	A	324	HIS	2.0
1	A	55	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	OMC	B	901	22/22	0.88	0.21	-	102,104,105,105	0
2	OMU	B	902	21/22	0.88	0.18	-	97,102,105,105	0
2	OMG	B	903	24/25	0.89	0.17	-	102,104,106,106	0
2	OMG	B	904	24/25	0.88	0.20	-	104,109,113,113	0

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	GOL	A	701	6/6	0.89	0.25	0.51	77,78,78,79	0
3	ZN	A	600	1/1	0.94	0.05	-3.69	94,94,94,94	0

6.5 Other polymers ⓘ

There are no such residues in this entry.