wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 05:06 PM BST

PDB ID	:	1TDP
Title	:	NMR solution structure of the carnobacteriocin B2 immunity protein
Authors	:	Sprules, T.; Kawulka, K.E.; Vederas, J.C.
Deposited on	:	2004-05-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 93%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	111	13% 44% 26% 17%

2 Ensemble composition and analysis (i)

This entry contains 15 models. Model 10 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:1-A:58, A:70-A:103 (92)	0.43	10

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 2, 5, 7, 9, 10, 11, 13, 15
2	3, 8
3	4, 6
Single-model clusters	12; 14

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1811 atoms, of which 921 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called carnobacteriocin B2 immunity protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	111	1811	571	921	143	172	4	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: carnobacteriocin B2 immunity protein

Chain A: 13% 44% 26% 17%

M1 D2 I3 K4 S5 Q6 T7 L8 Y9 L10 N11 L12 S13 E14 A15 Y16 K17 D18 P19 E20 V21 K22 A23 N24 E25 F26 L27 S28 K29 L30 V31 V32 Q33 C34 A35 G36 K37 L38 T39 A40 S41 E44 Y47 I48 E49 V50 I51 S52 L53 L54 S55 R56 G57 I58 S59 S60 Y61 Y62

L63 S64 H65 K66 R67 I68 I69 P70 S71 S72 M73 L74 T75 I76 Y77 T78 Q79 I80 Q81 K82 D83 I84 K85 N86 I89 D90 T91 E92 K93 L94 R95 K96 Y97 E98 I99 A100 K101 G102 L103 M104 S105 V106 P107 Y108 I109 Y110 F111

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 10. Colouring as in section 4.1 above.

- Molecule 1: carnobacteriocin B2 immunity protein

Chain A: 16% 41% 25% • 17%

M1 D2 I3 K4 S5 Q6 T7 L8 Y9 L10 N11 L12 S13 E14 A15 Y16 K17 D18 P19 E20 V21 K22 A23 N24 E25 F26 L27 S28 K29 L30 V31 V32 Q33 C34 A35 G36 K37 L38 T39 A40 S41 N42 S43 E44 Y47 I48 E49 V50 I51 S52 L53 L54 S55 R56 G57 I58 S59 S60 Y61

Y62 L63 S64 H65 K66 R67 I68 I69 P70 S71 S72 M73 L74 T75 I76 Y77 T78 Q79 I80 Q81 K82 D83 I84 K85 N86 I89 D90 T91 E92 K93 L94 R95 K96 Y97 E98 I99 A100 K101 G102 L103 M104 S105 V106 P107 Y108 I109 Y110 F111

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	1.1
CNS	refinement	1.1

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6211
Number of chemical shift lists	1
Total number of shifts	1330
Number of shifts mapped to atoms	1330
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	93%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	722	753	753	130±11
All	All	10830	11295	11295	1945

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 88.

5 of 754 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:51:ILE:HG22	1:A:94:LEU:HD21	1.07	1.25	6	8
1:A:95:ARG:O	1:A:99:ILE:HG13	1.06	1.47	9	8
1:A:12:LEU:HD22	1:A:31:VAL:HG13	1.01	1.33	2	5
1:A:53:LEU:O	1:A:57:GLY:HA3	1.00	1.56	7	8
1:A:28:SER:O	1:A:32:VAL:HG23	0.99	1.56	11	15

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	91/111 (82%)	77±2 (85±2%)	11±2 (12±2%)	2±1 (3±1%)	11 47
All	All	1365/1665 (82%)	1162 (85%)	167 (12%)	36 (3%)	11 47

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	72	SER	14
1	A	58	ILE	4
1	A	56	ARG	4
1	A	74	LEU	3
1	A	40	ALA	2

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	83/102 (81%)	45±4 (55±4%)	38±4 (45±4%)	0 2
All	All	1245/1530 (81%)	679 (55%)	566 (45%)	0 2

5 of 69 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	3	ILE	15
1	A	6	GLN	15
1	A	47	TYR	15
1	A	4	LYS	15
1	A	16	TYR	15

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 93% for the well-defined parts and 90% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 6211

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1330
Number of shifts mapped to atoms	1330
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	106	-0.41 ± 0.10	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	101	0.25 ± 0.07	None needed (< 0.5 ppm)
$^{13}C'$	98	-0.50 ± 0.09	None needed (< 0.5 ppm)
^{15}N	102	0.03 ± 0.15	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 93%, i.e. 1063 atoms were assigned a chemical shift out of a possible 1137. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	444/456 (97%)	179/182 (98%)	178/184 (97%)	87/90 (97%)
Sidechain	570/632 (90%)	354/369 (96%)	206/237 (87%)	10/26 (38%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	49/49 (100%)	25/25 (100%)	24/24 (100%)	0/0 (—%)
Overall	1063/1137 (93%)	558/576 (97%)	408/445 (92%)	97/116 (84%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: