wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 05:07 PM BST

PDB ID	:	1TNN
Title	:	Tertiary structure of an immunoglobulin-like domain from the giant muscle protein titin: a new member of the I set
Authors	:	Pfuhl, M.; Pastore, A.
Deposited on	:	1995-01-17

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	100	15% 58% 17% • 9%

2 Ensemble composition and analysis (i)

This entry contains 16 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. Model 4 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:1-A:91 (91)	0.66	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16
2	1, 10

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1411 atoms, of which 698 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called TITIN MODULE M5.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	91	1411	440	698	122	149	2	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	MET	-	EXPRESSION TAG	UNP Q8WZ42
A	-7	HIS	-	EXPRESSION TAG	UNP Q8WZ42
A	-6	HIS	-	EXPRESSION TAG	UNP Q8WZ42
A	-5	HIS	-	EXPRESSION TAG	UNP Q8WZ42
A	-4	HIS	-	EXPRESSION TAG	UNP Q8WZ42
A	-3	HIS	-	EXPRESSION TAG	UNP Q8WZ42
A	-2	HIS	-	EXPRESSION TAG	UNP Q8WZ42
A	-1	SER	-	EXPRESSION TAG	UNP Q8WZ42
A	0	SER	-	EXPRESSION TAG	UNP Q8WZ42

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: TITIN MODULE M5

Chain A: 15% 58% 17% • 9%

MET HIS HIS HIS HIS HIS HIS HIS SER SER R1 I2 L3 T4 K5 F6 R7 S8 N9 T10 V11 Y12 E13 G14 E15 S16 A17 R18 F19 D22 T23 D24 G25 E26 P27 V28 P29 T30 V31 T32 W33 L34 R35 K36 G37 Q38 V39 L40 S41 T42 S43 A44 R45 H46 Q47 V48 T49 T50 T51 K52

Y53 K54 S55 T56 F57 E58 I59 S60 S61 V62 Q63 A64 S65 D66 E67 G68 N69 Y70 S71 V72 V73 V74 E75 N76 S77 E78 G79 K80 Q81 E82 A83 E84 F85 T86 L87 T88 I89 Q90 K91

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 4. Colouring as in section 4.1 above.

- Molecule 1: TITIN MODULE M5

Chain A: 27% 43% 19% • 9%

MET HIS HIS HIS HIS HIS HIS HIS SER SER R1 I2 L3 R7 S8 N9 E13 G14 E15 S16 A17 R18 F19 D22 T23 D24 G25 E26 P27 V28 P29 T30 V31 T32 W33 L34 R35 Q38 V39 L40 S41 A44 R45 H46 Q47 V48 T49 T50 T51 K52 Y53 K54 S55 T56 F57 E58

I59 S60 S61 V62 Q63 A64 S65 D66 E67 G68 N69 Y70 S71 V72 V73 V74 E75 N76 S77 E78 G79 K80 Q81 E82 A83 E84 F85 T86 L87 T88 I89 Q90 K91

5 Refinement protocol and experimental data overview (i)

Of the ? calculated structures, 16 were deposited, based on the following criterion: *?*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DIANA	refinement	
X-PLOR	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.08±0.01	0±0/724 (0.0±0.0%)	1.24±0.01	0±0/978 (0.0±0.0%)
All	All	1.08	0/11584 (0.0%)	1.24	3/15648 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	4.9±0.2
All	All	0	79

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	43	SER	N-CA-CB	-5.80	101.80	110.50	15	3

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	18	ARG	Sidechain	16
1	A	45	ARG	Sidechain	16
1	A	7	ARG	Sidechain	16
1	A	1	ARG	Sidechain	16
1	A	35	ARG	Sidechain	15

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	713	699	701	91±8
All	All	11408	11183	11216	1450

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 64.

5 of 599 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:59:ILE:HD12	1:A:87:LEU:HD23	1.14	1.16	2	6
1:A:59:ILE:HD13	1:A:87:LEU:HD23	1.09	1.10	9	3
1:A:17:ALA:HB1	1:A:87:LEU:HD21	1.09	1.17	4	10
1:A:17:ALA:HB3	1:A:87:LEU:HD22	1.00	1.28	9	2
1:A:59:ILE:HG22	1:A:62:VAL:HG12	0.99	1.32	10	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	89/100 (89%)	53±3 (60±4%)	23±4 (26±4%)	13±2 (14±2%)	1 5
All	All	1424/1600 (89%)	853 (60%)	370 (26%)	201 (14%)	1 5

5 of 41 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	69	ASN	14
1	A	28	VAL	14
1	A	16	SER	13

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	67	GLU	12
1	A	2	ILE	12

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	82/91 (90%)	60±3 (73±4%)	22±3 (27±4%)	2 23
All	All	1312/1456 (90%)	963 (73%)	349 (27%)	2 23

5 of 58 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	88	THR	16
1	A	30	THR	15
1	A	86	THR	13
1	A	80	LYS	12
1	A	35	ARG	12

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided