



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:23 PM GMT

PDB ID : 3TQS
Title : Structure of the dimethyladenosine transferase (ksgA) from *Coxiella burnetii*
Authors : Rudolph, M.; Cheung, J.; Franklin, M.C.; Cassidy, M.; Gary, E.; Burshteyn, F.; Love, J.
Deposited on : 2011-09-09
Resolution : 1.98 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9100 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribosomal RNA small subunit methyltransferase A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	Se	0	0	0
			1999	1286	345	362	3	3			
1	B	247	Total	C	N	O	S	Se	0	0	0
			1999	1286	345	362	3	3			
1	C	247	Total	C	N	O	S	Se	0	0	0
			1999	1286	345	362	3	3			
1	D	247	Total	C	N	O	S	Se	0	0	0
			1999	1286	345	362	3	3			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	279	Total	O	0	0
			279	279		
2	B	272	Total	O	0	0
			272	272		
2	C	283	Total	O	0	0
			283	283		
2	D	270	Total	O	0	0
			270	270		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Ribosomal RNA small subunit methyltransferase A

Chain A: 




- Molecule 1: Ribosomal RNA small subunit methyltransferase A

Chain B: 





- Molecule 1: Ribosomal RNA small subunit methyltransferase A

Chain C: 



- Molecule 1: Ribosomal RNA small subunit methyltransferase A

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.01Å 57.90Å 112.94Å 90.00° 90.01° 104.34°	Depositor
Resolution (Å)	17.75 – 1.98 17.75 – 1.98	Depositor EDS
% Data completeness (in resolution range)	92.9 (17.75-1.98) 93.2 (17.75-1.98)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	14.31 (at 1.97Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.183 , 0.246 0.184 , 0.245	Depositor DCC
R_{free} test set	3066 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	12.2	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.4	EDS
Estimated twinning fraction	0.478 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 60556 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9100	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 98.54 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 2.0715e-11. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2043	0.52	0/2772
1	B	0.35	0/2043	0.52	0/2772
1	C	0.35	0/2043	0.52	0/2772
1	D	0.35	0/2043	0.52	0/2772
All	All	0.35	0/8172	0.52	0/11088

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1999	0	2017	22	0
1	B	1999	0	2017	19	0
1	C	1999	0	2017	17	0
1	D	1999	0	2017	22	0
2	A	279	0	0	3	0
2	B	272	0	0	1	0
2	C	283	0	0	1	0
2	D	270	0	0	2	0
All	All	9100	0	8068	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:155:VAL:HG12	1:D:156:MSE:HE3	1.36	1.03
1:C:155:VAL:HG12	1:C:156:MSE:HE3	1.36	1.03
1:D:156:MSE:HE2	1:D:156:MSE:HA	1.42	1.00
1:A:156:MSE:HA	1:A:156:MSE:HE2	1.42	0.99
1:A:155:VAL:HG12	1:A:156:MSE:HE3	1.41	0.99
1:D:156:MSE:SE	2:D:263:HOH:O	2.32	0.98
1:A:156:MSE:SE	2:A:366:HOH:O	2.32	0.96
1:C:156:MSE:SE	2:C:369:HOH:O	2.34	0.96
1:C:156:MSE:HA	1:C:156:MSE:HE2	1.43	0.95
1:C:234:LEU:HD13	1:C:253:ILE:HD13	1.53	0.90
1:A:234:LEU:HD13	1:A:253:ILE:HD13	1.55	0.88
1:A:52:THR:HG23	1:A:53:GLU:OE2	1.77	0.84
1:D:234:LEU:HD13	1:D:253:ILE:HD13	1.63	0.81
1:B:234:LEU:HD13	1:B:253:ILE:HD13	1.62	0.80
1:D:156:MSE:HA	1:D:156:MSE:CE	2.16	0.75
1:A:156:MSE:HA	1:A:156:MSE:CE	2.18	0.73
1:A:148:HIS:HE1	1:D:64:ARG:HH12	1.34	0.72
1:C:156:MSE:HA	1:C:156:MSE:CE	2.21	0.71
1:B:64:ARG:HH12	1:C:148:HIS:HE1	1.37	0.70
1:C:155:VAL:HG12	1:C:156:MSE:CE	2.20	0.70
1:D:234:LEU:HD13	1:D:253:ILE:CD1	2.28	0.61
1:B:141:ILE:HD11	1:B:153:LEU:HD11	1.83	0.59
1:B:234:LEU:HD13	1:B:253:ILE:CD1	2.32	0.57
1:D:155:VAL:HG12	1:D:156:MSE:CE	2.23	0.57
1:A:43:ARG:HG2	1:A:43:ARG:O	2.05	0.57
1:C:234:LEU:HD13	1:C:253:ILE:CD1	2.33	0.56
1:C:43:ARG:HG2	1:C:43:ARG:O	2.06	0.56
1:D:43:ARG:HG2	1:D:43:ARG:O	2.07	0.53
1:A:148:HIS:CE1	1:D:64:ARG:HH12	2.22	0.53
1:B:64:ARG:HH12	1:C:148:HIS:CE1	2.23	0.53
1:B:159:TYR:O	1:B:190:ARG:HD3	2.10	0.51
1:B:43:ARG:HG2	1:B:43:ARG:O	2.11	0.50
1:A:155:VAL:HG12	1:A:156:MSE:CE	2.27	0.49
1:D:47:THR:HG23	1:D:57:LEU:HD21	1.93	0.49
1:C:156:MSE:CA	1:C:156:MSE:HE2	2.31	0.48
1:A:234:LEU:HD13	1:A:253:ILE:CD1	2.37	0.48
1:B:47:THR:HG23	1:B:57:LEU:HD21	1.95	0.48
1:A:248:GLU:HG2	2:A:293:HOH:O	2.14	0.47
1:A:249:ASP:O	1:A:253:ILE:HG12	2.14	0.47
1:B:194:THR:HG22	1:B:195:PRO:HA	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:13:HIS:HD2	1:D:176:PRO:O	1.98	0.47
1:A:194:THR:HG23	1:A:195:PRO:HA	1.97	0.47
1:B:139:ARG:NE	2:B:970:HOH:O	2.48	0.47
1:B:194:THR:CG2	1:B:195:PRO:HA	2.46	0.46
1:C:194:THR:HG23	1:C:195:PRO:HA	1.98	0.46
1:A:231:TRP:N	1:A:232:PRO:CD	2.79	0.46
1:C:249:ASP:O	1:C:253:ILE:HG12	2.16	0.45
1:D:139:ARG:NE	2:D:958:HOH:O	2.49	0.45
1:D:107:LEU:HD13	1:D:115:LEU:HD23	1.99	0.45
1:D:194:THR:CG2	1:D:195:PRO:HA	2.45	0.45
1:B:16:HIS:HD2	1:B:175:THR:OG1	2.00	0.45
1:B:123:ILE:HD11	1:B:189:PRO:HG3	1.99	0.44
1:B:16:HIS:CD2	1:B:175:THR:OG1	2.70	0.44
1:D:132:MSE:HE1	1:D:169:VAL:HG21	1.98	0.44
1:C:231:TRP:N	1:C:232:PRO:CD	2.80	0.44
1:C:113:THR:N	1:C:114:PRO:HD2	2.32	0.44
1:B:132:MSE:HE1	1:B:169:VAL:HG21	1.99	0.44
1:D:159:TYR:O	1:D:190:ARG:HD3	2.18	0.44
1:B:13:HIS:HD2	1:B:176:PRO:O	2.02	0.43
1:A:113:THR:N	1:A:114:PRO:HD2	2.34	0.43
1:A:159:TYR:O	1:A:190:ARG:HD3	2.19	0.42
1:D:128:ASP:CB	1:D:188:ILE:HD13	2.49	0.42
1:D:107:LEU:CD1	1:D:115:LEU:HD23	2.49	0.42
1:A:139:ARG:NH2	2:A:820:HOH:O	2.52	0.42
1:C:47:THR:HG23	1:C:57:LEU:HD21	2.02	0.42
1:A:156:MSE:CA	1:A:156:MSE:CE	2.94	0.41
1:A:13:HIS:HD2	1:A:176:PRO:O	2.03	0.41
1:C:159:TYR:O	1:C:190:ARG:HD3	2.20	0.41
1:D:231:TRP:N	1:D:232:PRO:CD	2.83	0.41
1:D:249:ASP:O	1:D:253:ILE:HG12	2.20	0.41
1:B:231:TRP:N	1:B:232:PRO:CD	2.84	0.41
1:D:156:MSE:CA	1:D:156:MSE:CE	2.93	0.41
1:B:144:GLU:O	1:B:147:SER:HB3	2.21	0.41
1:A:107:LEU:HA	1:A:108:PRO:HD3	1.96	0.41
1:A:194:THR:CG2	1:A:195:PRO:HA	2.51	0.40
1:B:107:LEU:HD13	1:B:115:LEU:HD23	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	245/255 (96%)	245 (100%)	0	0	100	100
1	B	245/255 (96%)	245 (100%)	0	0	100	100
1	C	245/255 (96%)	245 (100%)	0	0	100	100
1	D	245/255 (96%)	245 (100%)	0	0	100	100
All	All	980/1020 (96%)	980 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	227/230 (99%)	221 (97%)	6 (3%)	54	51
1	B	227/230 (99%)	222 (98%)	5 (2%)	60	59
1	C	227/230 (99%)	222 (98%)	5 (2%)	60	59
1	D	227/230 (99%)	222 (98%)	5 (2%)	60	59
All	All	908/920 (99%)	887 (98%)	21 (2%)	58	57

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	111	ILE
1	A	136	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	156	MSE
1	A	222	LEU
1	A	225	LEU
1	A	227	ASN
1	B	136	GLU
1	B	147	SER
1	B	192	ASN
1	B	222	LEU
1	B	225	LEU
1	C	136	GLU
1	C	156	MSE
1	C	222	LEU
1	C	225	LEU
1	C	227	ASN
1	D	111	ILE
1	D	136	GLU
1	D	147	SER
1	D	156	MSE
1	D	225	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (23) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	13	HIS
1	A	16	HIS
1	A	89	GLN
1	A	148	HIS
1	A	227	ASN
1	A	258	ASN
1	B	13	HIS
1	B	16	HIS
1	B	29	HIS
1	B	89	GLN
1	B	148	HIS
1	B	239	GLN
1	B	258	ASN
1	C	16	HIS
1	C	148	HIS
1	C	227	ASN
1	C	239	GLN
1	D	13	HIS
1	D	16	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	29	HIS
1	D	89	GLN
1	D	148	HIS
1	D	239	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	244/255 (95%)	-0.74	0 100 100	6, 12, 25, 40	0
1	B	244/255 (95%)	-0.70	0 100 100	6, 12, 24, 38	0
1	C	244/255 (95%)	-0.73	0 100 100	7, 12, 24, 37	0
1	D	244/255 (95%)	-0.67	0 100 100	7, 12, 25, 37	0
All	All	976/1020 (95%)	-0.71	0 100 100	6, 12, 25, 40	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.