Full wwPDB X-ray Structure Validation Report (i)

Feb 1, 2016 – 08:33 PM GMT

PDB ID	:	4TR7
Title	:	Crystal structure of DNA polymerase sliding clamp from Mycobaterium tuberculosis
Authors	:	Olieric, V.; Burnouf, D.; Ennifar, E.; Wolff, P.
Deposited on	:	2014-06-14
Resolution	:	2.29 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.29 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	402	5% 88% 5% 7%
1	B	402	2% 87% • 8%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 10662 atoms, of which 5161 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase III subunit beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	375	5266	1694	2605	449	512	6	0	2	0
1	B	370	5179	1678	2556	440	499	6	0	2	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	103	103	103	0	0
2	B	114	114	114	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA polymerase III subunit beta

Chain A:





- Molecule 1: DNA polymerase III subunit beta

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α, β, γ	40.49Å 83.65Å 84.03Å 63.58° 86.84° 85.20°	Depositor
Resolution (Å)	18.76 – 2.29 36.38 – 2.29	Depositor EDS
% Data completeness (in resolution range)	96.2 (18.76-2.29) 81.8 (36.38-2.29)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	2.90 (at 2.29Å)	Xtriage
Refinement program	BUSTER 2.10.0	Depositor
R, R_{free}	0.202 , 0.238 0.204 , 0.240	Depositor DCC
R_{free} test set	2152 reflections (5.33%)	DCC
Wilson B-factor (Å^2)	39.3	Xtriage
Anisotropy	0.888	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 53.1	EDS
Estimated twinning fraction	0.009 for -h,-l,-k	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 42569 reflections	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	10662	wwPDB-VP
Average B, all atoms (Å^2)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.73% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.48	0/2707	0.68	1/3696 (0.0%)
1	B	0.50	1/2672 (0.0%)	0.67	0/3657
All	All	0.49	1/5379 (0.0%)	0.68	1/7353 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	186	ARG	CG-CD	-5.28	1.38	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	A	375	PRO	N-CA-C	6.38	128.68	112.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	37	ARG	Mainchain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2661	2605	2602	8	0
1	B	2623	2556	2574	12	0
2	A	103	0	0	1	0
2	B	114	0	0	1	0
All	All	5501	5161	5176	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:61:ASP:H	1:B:64:VAL:HG12	1.56	0.70
1:B:221:GLY:HA3	1:B:222:GLY:O	1.94	0.67
1:B:219:GLY:O	1:B:221:GLY:N	2.28	0.67
1:A:64:VAL:HG22	1:A:251:THR:OG1	2.03	0.59
1:A:378:GLY:HA2	2:A:506:HOH:O	2.04	0.56
1:B:64:VAL:CG2	2:B:596:HOH:O	2.55	0.54
1:A:64:VAL:HG11	1:A:131:PRO:HD2	1.88	0.54
1:B:94:LEU:HD11	1:B:110:LEU:HD21	1.91	0.51
1:B:221:GLY:H	1:B:222:GLY:HA2	1.78	0.48
1:A:328:ALA:HB2	1:A:377:ALA:HB2	1.95	0.47
1:A:238:ASP:HA	1:A:239:GLY:HA2	1.67	0.46
1:B:26:ALA:HB1	1:B:57:ILE:HD12	1.97	0.46
1:B:273:VAL:HG23	1:B:328:ALA:HB3	1.96	0.46
1:B:64:VAL:HG22	1:B:251:THR:OG1	2.17	0.43
1:A:94:LEU:HD11	1:A:110:LEU:HD21	2.01	0.43
1:B:273:VAL:CG2	1:B:328:ALA:HB3	2.50	0.42
1:A:307:VAL:HB	1:A:325:VAL:HG21	2.00	0.42
1:A:117:PHE:CE2	1:B:319:ALA:HB2	2.55	0.41
1:B:221:GLY:N	1:B:222:GLY:HA2	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/402 (90%)	343 (94%)	17 (5%)	3 (1%)	24	27
1	B	364/402 (90%)	351 (96%)	10 (3%)	3 (1%)	24	27
All	All	727/804 (90%)	694 (96%)	27 (4%)	6 (1%)	24	27

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	375	PRO
1	B	220	ILE
1	A	237	LYS
1	A	61	ASP
1	B	61	ASP
1	B	223	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/316 (83%)	258 (98%)	4 (2%)	72	85
1	B	256/316 (81%)	254 (99%)	2 (1%)	86	94
All	All	518/632 (82%)	512 (99%)	6 (1%)	78	89

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	186	ARG
1	A	194	TRP
1	A	258	GLU
1	A	379	LEU
1	B	194	TRP
1	B	326	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	375/402 (93%)	0.19	19 (5%) 32 41	34, 53, 87, 109	0
1	B	370/402 (92%)	0.03	8 (2%) 65 73	34, 49, 76, 100	0
All	All	745/804 (92%)	0.11	27 (3%) 46 55	34, 51, 81, 109	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	221	GLY	8.2
1	A	370	SER	4.8
1	B	36	ALA	4.8
1	A	371	GLY	4.7
1	B	230	GLY	4.3
1	A	199	PRO	4.2
1	A	139	LEU	3.6
1	A	361	GLY	3.6
1	A	200	ASP	3.4
1	A	53	ASN	3.3
1	A	161	THR	2.9
1	A	126	ASP	2.6
1	A	43	LEU	2.6
1	A	197	SER	2.5
1	A	238	ASP	2.5
1	B	221	GLY	2.5
1	B	62	TYR	2.4
1	A	62	TYR	2.4
1	A	201	ILE	2.3
1	B	381	GLY	2.3
1	A	52	ASP	2.2
1	B	198	SER	2.1
1	A	258	GLU	2.1
1	A	125	GLU	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	142	ALA	2.0
1	B	319	ALA	2.0
1	A	123	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.