Full wwPDB X-ray Structure Validation Report (i)

Feb 1, 2016 – 05:47 AM GMT

PDB ID	:	2TSS
Title	:	TOXIC SHOCK SYNDROME TOXIN-1 FROM STAPHYLOCOCCUS AUREUS: ORTHORHOMBICC222(1) CRYSTAL FORM
Authors	:	Prasad, G.S.; Radhakrishnan, R.; Mitchell, D.T.; Earhart, C.A.; Dinges, M.M.; Cook, W.J.; Schlivert, P.M.; Ohlendorf, D.H.
Deposited on	:	1996-12-04
Resolution	:	2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1269 (2.04-2.04)
Ramachandran outliers	100387	1258 (2.04-2.04)
Sidechain outliers	100360	1258 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	194	91% 9% •
1	B	194	90% 10%
1	C	194	91% 9% •

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 5119 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TOXIC SHOCK SYNDROME TOXIN-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	194	1580	1005	258	315	2	0	5	0
1	B	194	1566	996	257	311	2	0	1	0
1	C	194	1568	997	257	312	2	0	2	0

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	142	142	142	0	0
2	B	141	141	141	0	0
2	C	122	122	122	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: TOXIC SHOCK SYNDROME TOXIN-1

Chain A:



- Molecule 1: TOXIC SHOCK SYNDROME TOXIN-1

Chain B:



- Molecule 1: TOXIC SHOCK SYNDROME TOXIN-1

Chain C:



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α, β, γ	108.70Å 177.50Å 97.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	5.00 – 2.05	Depositor
% Data completeness (in resolution range)	88.4 (5.00-2.05)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.154 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5119	wwPDB-VP
Average B, all atoms (Å^2)	24.0	wwPDB-VP

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.46	0/1637	0.71	1/2214 (0.0%)
1	B	0.50	0/1607	0.71	1/2173 (0.0%)
1	C	0.57	2/1613 (0.1%)	0.76	3/2181 (0.1%)
All	All	0.51	2/4857 (0.0%)	0.72	5/6568 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	29[A]	SER	CA-CB	6.83	1.63	1.52
1	C	29[B]	SER	CA-CB	6.83	1.63	1.52

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	29[A]	SER	N-CA-CB	6.68	120.52	110.50
1	C	29[B]	SER	N-CA-CB	6.68	120.52	110.50
1	A	65	ASN	N-CA-C	-5.93	94.98	111.00
1	B	65	ASN	N-CA-C	-5.10	97.22	111.00
1	C	65	ASN	N-CA-C	-5.10	97.24	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1580	0	1558	12	0
1	B	1566	0	1548	7	0
1	C	1568	0	1549	6	0
2	A	142	0	0	1	0
2	B	141	0	0	0	0
2	C	122	0	0	0	0
All	All	5119	0	4655	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:155:LYS:HG2	1:C:165:GLN:HB3	1.69	0.73
1:A:103:LYS:HE3	1:A:105:LYS:HG3	1.79	0.63
1:B:105:LYS:HB2	1:B:192:GLU:HG2	1.83	0.60
1:A:103:LYS:HE2	1:A:103:LYS:O	2.02	0.59
1:C:75:THR:OG1	1:C:77:GLU:HG2	2.09	0.53
1:C:1:SER:O	1:C:147:SER:HB3	2.08	0.52
1:C:152:GLY:HA3	1:C:192:GLU:O	2.12	0.50
1:A:51:TYR:O	1:A:179:PRO:HG3	2.12	0.49
1:A:175[A]:ASN:ND2	2:A:324:HOH:O	2.43	0.48
1:A:48:PRO:HB2	1:A:175[B]:ASN:HD22	1.78	0.48
1:C:35:ILE:HD12	1:C:45:ILE:HD12	1.95	0.47
1:A:105:LYS:HB2	1:A:192:GLU:HG2	1.98	0.46
1:A:103:LYS:HE2	1:A:103:LYS:C	2.36	0.46
1:B:107:HIS:CE1	1:B:194:ASN:OXT	2.71	0.44
1:B:70:LYS:HE3	1:B:83:PHE:CD2	2.53	0.44
1:A:1:SER:O	1:A:147:SER:HB3	2.17	0.44
1:B:135:HIS:O	1:B:139:GLN:HG2	2.19	0.43
1:A:47:PHE:CE1	1:A:86:SER:HA	2.54	0.43
1:C:105:LYS:HB2	1:C:192:GLU:HG2	2.02	0.42
1:B:23:SER:HB3	1:B:35:ILE:CG2	2.49	0.42
1:A:67:LYS:HE2	1:A:131[A]:PHE:CE1	2.55	0.42
1:A:111:SER:HA	1:A:112:PRO:HD2	1.94	0.42
1:B:111:SER:HA	1:B:112:PRO:HD2	1.83	0.41
1:B:100:LEU:HA	1:B:101:PRO:HD3	1.88	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/194 (102%)	191 (97%)	6 (3%)	0	100	100
1	B	193/194 (100%)	189 (98%)	4 (2%)	0	100	100
1	C	194/194 (100%)	188 (97%)	6 (3%)	0	100	100
All	All	584/582 (100%)	568 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/180 (103%)	181 (98%)	4 (2%)	60	53
1	B	181/180 (101%)	177 (98%)	4 (2%)	60	53
1	C	182/180 (101%)	176 (97%)	6 (3%)	45	37
All	All	548/540 (102%)	534 (97%)	14 (3%)	57	47

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	79	THR
1	A	99	GLU
1	A	175[A]	ASN
1	A	175[B]	ASN
1	B	76	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	96	THR
1	B	110	ASP
1	B	114	LYS
1	C	4	ASP
1	C	46	ILE
1	C	103	LYS
1	C	114	LYS
1	C	155	LYS
1	C	187	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	22	ASN
1	B	73	GLN
1	B	107	HIS
1	C	73	GLN
1	C	107	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates (i)

EDS was not executed - this section will therefore be empty.

6.4 Ligands (i)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers (i)

EDS was not executed - this section will therefore be empty.