



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:56 PM GMT

PDB ID : 4U91  
Title : GephE in complex with Para-Phenyl crosslinked Glycine receptor beta subunit derived dimeric peptide  
Authors : Kasaragod, V.B.; Maric, H.M.; Schindelin, H.  
Deposited on : 2014-08-05  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	419	<div> <div>7%</div> <div>82%</div> <div>14%</div> <div>••</div> </div>
2	B	11	<div> <div>27%</div> <div>55%</div> <div>27%</div> <div>18%</div> </div>
2	E	11	<div> <div>27%</div> <div>55%</div> <div>27%</div> <div>18%</div> </div>

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 3689 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Gephyrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	415	Total	C	N	O	S	0	7	0
			3206	2023	557	605	21			

- Molecule 2 is a protein called Glycine receptor subunit beta.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	11	Total	C	N	O	S	0	0	0
			83	52	14	16	1			
2	E	11	Total	C	N	O	S	0	0	0
			83	52	14	16	1			

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is 1,1'-benzene-1,4-diylbis(1H-pyrrole-2,5-dione) (three-letter code: 3F7) (formula:  $C_{14}H_8N_2O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	E	1	Total	C	N	O	0	0
			20	14	2	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	283	Total	O	0	0
			283	283		
7	B	3	Total	O	0	0
			3	3		
7	E	2	Total	O	0	0
			2	2		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gephyrin



- Molecule 2: Glycine receptor subunit beta



- Molecule 2: Glycine receptor subunit beta



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	87.44Å 99.21Å 112.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.00 56.24 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-2.00) 100.0 (56.24-2.00)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.39 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, $R_{free}$	0.169 , 0.226 0.179 , 0.229	Depositor DCC
$R_{free}$ test set	1688 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.4	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 33406 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3689	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.93% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 3F7, CA, EDO, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	0/3281	0.89	6/4459 (0.1%)
2	B	0.74	1/84 (1.2%)	0.82	1/111 (0.9%)
2	E	0.81	0/84	0.89	1/111 (0.9%)
All	All	0.79	1/3449 (0.0%)	0.88	8/4681 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	405	PRO	N-CD	5.03	1.54	1.47

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	736	LEU	CA-CB-CG	8.03	133.76	115.30
2	E	404	LEU	C-N-CD	6.04	141.09	128.40
1	A	653	ARG	NE-CZ-NH2	-5.78	117.41	120.30
1	A	736	LEU	CB-CG-CD2	5.78	120.82	111.00
1	A	653	ARG	NE-CZ-NH1	5.75	123.17	120.30
1	A	504	MET	CG-SD-CE	-5.71	91.06	100.20
2	B	404	LEU	C-N-CD	5.66	140.28	128.40
1	A	736	LEU	CB-CG-CD1	-5.61	101.47	111.00

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3206	0	3264	61	0
2	B	83	0	80	5	0
2	E	83	0	80	7	0
3	A	4	0	3	0	0
4	A	4	0	6	0	0
5	A	1	0	0	0	0
6	E	20	0	6	4	0
7	A	283	0	0	11	1
7	B	3	0	0	0	0
7	E	2	0	0	0	0
All	All	3689	0	3439	75	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:735[B]:ARG:NH1	1:A:735[B]:ARG:HB2	1.59	1.16
1:A:735[B]:ARG:HB2	1:A:735[B]:ARG:CZ	1.79	1.10
1:A:653:ARG:NH1	7:A:901:HOH:O	1.91	1.02
1:A:436:SER:O	1:A:439:GLY:N	2.05	0.89
1:A:404:MET:HG2	7:A:969:HOH:O	1.75	0.86
1:A:575:SER:OG	1:A:580:ASP:OD2	1.96	0.82
1:A:735[B]:ARG:HD3	7:A:1002:HOH:O	1.82	0.78
1:A:735[B]:ARG:CB	1:A:735[B]:ARG:CZ	2.60	0.78
1:A:319:SER:HA	1:A:320:PRO:C	2.03	0.77
1:A:438:ASP:HB2	1:A:440:THR:OG1	1.86	0.75
2:B:406:ARG:NH2	2:B:406:ARG:HG2	2.04	0.72
1:A:382:ASP:OD2	1:A:406:GLY:N	2.16	0.71
2:B:406:ARG:HG2	2:B:406:ARG:HH21	1.54	0.70
1:A:320:PRO:HB3	1:A:576:MET:CE	2.21	0.69
1:A:584:GLN:CD	1:A:584:GLN:H	1.95	0.69
1:A:458:ARG:NH1	1:A:463:ASP:OD2	2.26	0.68
1:A:578:GLU:O	1:A:579:LYS:HB2	1.92	0.68
1:A:524:ASP:OD1	7:A:1087:HOH:O	2.13	0.66
1:A:605:LEU:HB3	1:A:606:PRO:HD3	1.78	0.65
1:A:735[B]:ARG:NH1	1:A:735[B]:ARG:CB	2.49	0.63
2:E:406:ARG:CG	2:E:407:ASP:H	2.12	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:406:ARG:CG	2:B:406:ARG:HH21	2.13	0.61
1:A:578:GLU:OE2	1:A:578:GLU:HA	2.01	0.60
1:A:735[B]:ARG:HB2	1:A:735[B]:ARG:HH11	1.63	0.58
1:A:512:ASN:HD22	1:A:527:ARG:HH22	1.52	0.58
1:A:438:ASP:OD1	1:A:438:ASP:N	2.28	0.57
1:A:578:GLU:O	1:A:579:LYS:CB	2.50	0.57
6:E:501:3F7:OAC	6:E:501:3F7:H4	2.05	0.57
1:A:436:SER:HG	1:A:440:THR:H	1.52	0.56
1:A:690:GLN:OE1	1:A:701:MET:HG2	2.07	0.55
6:E:501:3F7:OAB	6:E:501:3F7:H3	2.07	0.55
1:A:504:MET:CG	1:A:570:THR:HG22	2.37	0.55
1:A:506:THR:HA	1:A:547:VAL:O	2.08	0.54
6:E:501:3F7:H6	6:E:501:3F7:OAD	2.06	0.54
1:A:336[B]:MET:CE	1:A:595:HIS:CE1	2.90	0.54
1:A:364:ASN:ND2	1:A:466:ARG:H	2.05	0.54
6:E:501:3F7:H5	6:E:501:3F7:OAA	2.09	0.53
1:A:378:VAL:HG12	1:A:408:VAL:HG12	1.91	0.53
1:A:511:LEU:HD22	7:A:1039:HOH:O	2.08	0.53
1:A:735[B]:ARG:CZ	1:A:735[B]:ARG:H	2.23	0.52
1:A:319:SER:HA	1:A:321:PHE:N	2.24	0.51
1:A:719:TYR:CD2	2:B:404:LEU:HD22	2.44	0.51
2:E:407:ASP:O	2:E:408:CYS:HB2	2.08	0.51
1:A:433:ILE:HG22	1:A:434:ARG:HG2	1.92	0.51
1:A:336[B]:MET:HE2	1:A:595:HIS:CE1	2.46	0.49
2:E:406:ARG:HG3	2:E:407:ASP:H	1.76	0.48
1:A:555:LEU:HD22	1:A:589:ASP:HB3	1.95	0.48
1:A:736:LEU:HB2	7:A:1010:HOH:O	2.14	0.47
1:A:394:GLN:HA	7:A:914:HOH:O	2.15	0.47
2:B:406:ARG:HG3	2:B:407:ASP:N	2.30	0.47
1:A:320:PRO:HB3	1:A:576:MET:HE1	1.93	0.46
1:A:379:ARG:HD2	1:A:405:PRO:HA	1.98	0.46
1:A:434:ARG:HB3	7:A:1001:HOH:O	2.16	0.46
1:A:735[B]:ARG:NH2	1:A:735[B]:ARG:H	2.14	0.45
2:E:406:ARG:CG	2:E:407:ASP:N	2.79	0.45
1:A:351:MET:HG2	1:A:482[B]:ILE:CD1	2.47	0.44
1:A:561:GLY:C	1:A:568:ILE:HD11	2.38	0.44
1:A:436:SER:O	1:A:439:GLY:CA	2.67	0.43
1:A:523:ARG:NH2	7:A:909:HOH:O	2.50	0.43
1:A:452:ARG:O	1:A:455:GLN:HB2	2.18	0.43
1:A:459:PRO:HD2	1:A:462:HIS:HB2	2.00	0.43
1:A:700:LEU:HD23	1:A:700:LEU:HA	1.95	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:715:LYS:NZ	7:A:1104:HOH:O	2.52	0.42
2:E:406:ARG:HG3	2:E:407:ASP:N	2.34	0.42
2:E:400:ILE:HA	2:E:400:ILE:HD12	1.82	0.42
1:A:351:MET:HG2	1:A:482[B]:ILE:HD11	2.02	0.42
1:A:588:ILE:O	1:A:591:HIS:CE1	2.73	0.42
1:A:699:ARG:CG	1:A:700:LEU:N	2.83	0.42
1:A:412:THR:HG23	7:A:914:HOH:O	2.20	0.41
1:A:378:VAL:HG12	1:A:408:VAL:CG1	2.50	0.41
1:A:346:ASN:ND2	1:A:348:ARG:H	2.18	0.41
1:A:436:SER:C	1:A:439:GLY:H	2.15	0.41
1:A:724:LYS:HE2	2:E:408:CYS:C	2.41	0.41
1:A:320:PRO:CB	1:A:576:MET:CE	2.96	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1090:HOH:O	7:A:1090:HOH:O[2_655]	1.62	0.58

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	418/419 (100%)	412 (99%)	6 (1%)	0	100	100
2	B	9/11 (82%)	9 (100%)	0	0	100	100
2	E	9/11 (82%)	8 (89%)	1 (11%)	0	100	100
All	All	436/441 (99%)	429 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/356 (101%)	342 (95%)	17 (5%)	32	27
2	B	10/10 (100%)	8 (80%)	2 (20%)	1	0
2	E	10/10 (100%)	8 (80%)	2 (20%)	1	0
All	All	379/376 (101%)	358 (94%)	21 (6%)	27	21

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	392	GLU
1	A	394	GLN
1	A	434	ARG
1	A	438	ASP
1	A	440	THR
1	A	441	GLU
1	A	578	GLU
1	A	579	LYS
1	A	584	GLN
1	A	632[A]	VAL
1	A	632[B]	VAL
1	A	647	GLN
1	A	698	SER
1	A	699	ARG
1	A	700	LEU
1	A	702	SER
1	A	736	LEU
2	B	406	ARG
2	B	408	CYS
2	E	406	ARG
2	E	408	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	346	ASN
1	A	364	ASN
1	A	496	ASN
1	A	512	ASN
1	A	591	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ACT	A	801	-	1,3,3	2.74	1 (100%)	0,3,3	0.00	-
4	EDO	A	802	-	3,3,3	0.40	0	2,2,2	0.28	0
6	3F7	E	501	-	22,22,22	4.17	10 (45%)	32,32,32	1.24	5 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ACT	A	801	-	-	0/0/0/0	0/0/0/0
4	EDO	A	802	-	-	0/1/1/1	0/0/0/0
6	3F7	E	501	-	-	0/8/34/34	0/3/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	E	501	3F7	CAK-CAQ	-7.51	1.33	1.48
6	E	501	3F7	CAJ-CAP	-7.47	1.33	1.48
6	E	501	3F7	CAL-CAR	-7.44	1.33	1.48
6	E	501	3F7	CAI-CAO	-7.44	1.33	1.48
6	E	501	3F7	CAM-NAS	-7.03	1.33	1.44
6	E	501	3F7	CAN-NAT	-6.90	1.33	1.44
6	E	501	3F7	CAO-NAS	-4.05	1.32	1.40
6	E	501	3F7	CAR-NAT	-4.03	1.32	1.40
6	E	501	3F7	CAP-NAS	-3.80	1.33	1.40
6	E	501	3F7	CAQ-NAT	-3.65	1.33	1.40
3	A	801	ACT	CH3-C	2.74	1.52	1.48

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	E	501	3F7	OAC-CAQ-CAK	-2.31	124.03	128.40
6	E	501	3F7	OAB-CAP-CAJ	-2.15	124.33	128.40
6	E	501	3F7	CAL-CAR-NAT	2.00	110.44	106.14
6	E	501	3F7	CAJ-CAP-NAS	2.11	110.68	106.14
6	E	501	3F7	CAK-CAQ-NAT	2.13	110.71	106.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	501	3F7	4	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	415/419 (99%)	0.23	29 (6%) 19 21	30, 41, 84, 114	0
2	B	11/11 (100%)	1.30	3 (27%) 1 1	47, 59, 148, 167	11 (100%)
2	E	11/11 (100%)	1.24	3 (27%) 1 1	38, 47, 93, 95	11 (100%)
All	All	437/441 (99%)	0.28	35 (8%) 15 16	30, 42, 88, 167	22 (5%)

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	699	ARG	5.5
1	A	440	THR	5.4
1	A	439	GLY	5.0
1	A	579	LYS	5.0
2	E	408	CYS	4.6
1	A	682	HIS	4.5
1	A	736	LEU	4.5
2	B	408	CYS	4.5
2	E	406	ARG	4.4
2	B	406	ARG	4.2
2	B	407	ASP	3.8
1	A	438	ASP	3.7
2	E	407	ASP	3.6
1	A	453	PRO	3.4
1	A	451	ALA	3.2
1	A	578	GLU	3.0
1	A	381	ALA	3.0
1	A	419	CYS	2.9
1	A	735[A]	ARG	2.8
1	A	694	ASN	2.7
1	A	693	GLY	2.7
1	A	420	GLY	2.7
1	A	319	SER	2.7

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	701	MET	2.7
1	A	436	SER	2.6
1	A	405	PRO	2.5
1	A	388	PHE	2.5
1	A	615	ASP	2.5
1	A	698	SER	2.5
1	A	577	GLY	2.3
1	A	437	ASP	2.2
1	A	576	MET	2.2
1	A	717	GLU	2.1
1	A	718	GLN	2.0
1	A	421	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
6	3F7	E	501	20/20	0.69	0.39	1.83	71,102,146,149	20
5	CA	A	803	1/1	0.96	0.04	-2.95	56,56,56,56	0
4	EDO	A	802	4/4	0.85	0.36	-	57,58,60,60	0
3	ACT	A	801	4/4	0.80	0.15	-	67,68,70,71	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.