



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 30, 2017 – 10:30 PM EST

PDB ID : 5UAG  
Title : Escherichia coli RNA polymerase mutant - RpoB D516V  
Authors : Molodtsov, V.; Scharf, N.T.; Stefan, M.A.; Garcia, G.A.; Murakami, K.S.  
Deposited on : 2016-12-19  
Resolution : 3.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442




# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1476 (3.50-3.30)
Clashscore	102246	1611 (3.50-3.30)
Ramachandran outliers	100387	1571 (3.50-3.30)
Sidechain outliers	100360	1571 (3.50-3.30)
RSRZ outliers	91569	1485 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	<div><div></div><div><div></div><div>39%</div><div>30%</div><div>•</div><div>27%</div></div></div>
1	B	320	<div><div>8%</div><div><div></div><div>32%</div><div>32%</div><div>•</div><div>32%</div></div></div>
1	G	320	<div><div>3%</div><div><div></div><div>36%</div><div>30%</div><div>•</div><div>29%</div></div></div>
1	H	320	<div><div>7%</div><div><div></div><div>33%</div><div>32%</div><div>•</div><div>33%</div></div></div>
2	C	1342	<div><div>4%</div><div><div></div><div>57%</div><div>39%</div><div>•</div></div></div>
2	I	1342	<div><div>7%</div><div><div></div><div>58%</div><div>38%</div><div>•</div></div></div>

Continued on next page...



Continued from previous page...

Mol	Chain	Length	Quality of chain
3	D	1407	<div><div><div></div><div></div><div></div><div></div></div><div>2%47%31%•17%</div></div>
3	J	1407	<div><div><div></div><div></div><div></div><div></div></div><div>4%45%33%•18%</div></div>
4	E	90	<div><div><div></div><div></div><div></div><div></div></div><div>3%60%30%9%•</div></div>
4	K	90	<div><div><div></div><div></div><div></div><div></div></div><div>20%54%27%7%12%</div></div>
5	F	613	<div><div><div></div><div></div><div></div><div></div></div><div>4%45%28%•24%</div></div>
5	L	613	<div><div><div></div><div></div><div></div><div></div></div><div>4%45%28%•23%</div></div>



## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 55066 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	233	Total	C	N	O	S	0	0	0
			1812	1127	323	356	6			
1	B	217	Total	C	N	O	S	0	0	0
			1677	1047	295	329	6			
1	G	227	Total	C	N	O	S	0	0	0
			1755	1093	311	345	6			
1	H	216	Total	C	N	O	S	0	0	0
			1662	1038	292	326	6			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1340	Total	C	N	O	S	0	0	0
			10569	6632	1841	2053	43			
2	I	1340	Total	C	N	O	S	0	0	0
			10565	6630	1840	2052	43			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	516	VAL	ASP	engineered mutation	UNP P0A8V2
I	516	VAL	ASP	engineered mutation	UNP P0A8V2

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1167	Total	C	N	O	S	0	0	0
			9065	5700	1622	1697	46			
3	J	1155	Total	C	N	O	S	0	0	0
			9001	5659	1612	1684	46			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	89	Total	C	N	O	S	0	0	0
			691	421	129	140	1			
4	K	79	Total	C	N	O	S	0	0	0
			627	382	118	126	1			

- Molecule 5 is a protein called RNA polymerase sigma factor RpoD.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	468	Total	C	N	O	S	0	0	0
			3813	2389	678	723	23			
5	L	469	Total	C	N	O	S	0	0	0
			3821	2393	679	726	23			

- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	J	1	Total	Mg	0	0
			1	1		
6	I	1	Total	Mg	0	0
			1	1		
6	D	2	Total	Mg	0	0
			2	2		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	J	2	Total	Zn	0	0
			2	2		
7	D	2	Total	Zn	0	0
			2	2		




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: DNA-directed RNA polymerase subunit alpha



#### • Molecule 1: DNA-directed RNA polymerase subunit alpha





Chain G:

Amino Acid	Percentage
Met	3%
Gln	36%
Gly	36%
Ser	36%
Val	36%
Thr	36%
Pro	36%
Leu	36%
Ala	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%
Val	36%
Leu	36%
Glu	36%
Gly	36%
Arg	36%
Asp	36%
Asn	36%
His	36%
Ile	36%
Tyr	36%
Cys	36%
Trp	36%
Phe	36%
Lys	36%


[illegible]

Chain C:

4% 57% 39%

Label	Color
RET	Grey
VAL	Grey
V3	Yellow
S4	Yellow
V5	Yellow
T6	Green
E7	Green
K8	Green
R9	Green
R10	Green
I11	Green
R12	Green
K13	Green
D14	Green
F15	Green
G16	Green
K17	Green
R18	Green
P19	Green
Q20	Green
V26	Green
L27	Green
L28	Green
S29	Green
T30	Green
K37	Green
F38	Green
I39	Orange
E40	Orange
Q41	Green
G45	Green
Q46	Green
V47	Green
G48	Green
F53	Green
R54	Green
S55	Green
V56	Green
F57	Green
F58	Green
I59	Green
Q60	Green
S61	Green
V62	Green
S63	Green
P64	Green
N65	Green
S66	Green
E67	Green
L68	Green
Q69	Green
V70	Green
V71	Green
S72	Green
V73	Green
T74	Green
L75	Green
G76	Green
E77	Green
F80	Green
D81	Green
R82	Green
Q83	Green
E84	Green
C85	Green
G86	Green
V90	Green
T91	Green
Y92	Green
S93	Green
A94	Green
P95	Green
L96	Green
R97	Green
V98	Green
K99	Green
L100	Green
R101	Green
L102	Green
V103	Green
I104	Green
Y105	Green
E106	Green
R107	Orange
E108	Orange
A109	Orange
P110	Green
E111	Green
G112	Green
T113	Green
V114	Green
K115	Green
D116	Green
I117	Green
K118	Green
E119	Green
Q120	Green
E121	Green
I122	Green
P123	Green
L129	Green
M124	Green
G125	Green
E126	Green
I127	Green
P128	Green
G129	Green
D132	Green
F136	Green
V137	Green
I138	Green
N139	Green
G140	Green
T141	Green





• Molecule 2: DNA-directed RNA polymerase subunit beta














• Molecule 3: DNA-directed RNA polymerase subunit beta'








• Molecule 3: DNA-directed RNA polymerase subunit beta'












- Molecule 5: RNA polymerase sigma factor RpoD





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	184.83Å 205.04Å 307.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.99 – 3.40 29.99 – 3.40	Depositor EDS
% Data completeness (in resolution range)	89.0 (29.99-3.40) 89.0 (29.99-3.40)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 3.39Å)	Xtriage
Refinement program	PHENIX (1.10.1 _2155: ???)	Depositor
R, $R_{free}$	0.277 , 0.311 0.273 , 0.310	Depositor DCC
$R_{free}$ test set	1970 reflections (1.38%)	DCC
Wilson B-factor (Å <sup>2</sup> )	135.9	Xtriage
Anisotropy	0.254	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 85.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	55066	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	163.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.84% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.46	1/1834 (0.1%)	0.85	0/2485
1	B	0.45	0/1697	1.00	3/2300 (0.1%)
1	G	0.52	1/1777 (0.1%)	0.93	5/2408 (0.2%)
1	H	0.44	0/1681	0.98	3/2278 (0.1%)
2	C	0.44	0/10738	0.83	7/14488 (0.0%)
2	I	0.42	1/10734 (0.0%)	0.81	12/14483 (0.1%)
3	D	0.45	0/9205	0.84	12/12430 (0.1%)
3	J	0.45	0/9140	0.88	22/12341 (0.2%)
4	E	0.43	0/693	0.76	1/935 (0.1%)
4	K	0.43	0/629	0.81	2/847 (0.2%)
5	F	0.42	0/3864	0.83	5/5194 (0.1%)
5	L	0.43	0/3872	0.83	3/5205 (0.1%)
All	All	0.44	3/55864 (0.0%)	0.85	75/75394 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1
2	C	0	4
2	I	0	2
3	D	0	2
3	J	0	3
4	E	0	1
5	L	0	1
All	All	0	14

All (3) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	G	29	GLU	C-N	9.36	1.52	1.34
2	I	373	GLY	C-N	7.20	1.50	1.34
1	A	29	GLU	C-N	6.31	1.46	1.34

The worst 5 of 75 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	J	425	ARG	NE-CZ-NH2	14.52	127.56	120.30
3	J	425	ARG	NE-CZ-NH1	-13.50	113.55	120.30
3	J	343	LEU	CB-CG-CD1	-11.67	91.16	111.00
1	G	12	ARG	NE-CZ-NH1	-10.66	114.97	120.30
3	D	343	LEU	CB-CG-CD2	-10.37	93.37	111.00

There are no chirality outliers.

5 of 14 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	C	109	ALA	Peptide
2	C	236	LYS	Peptide
2	C	658	GLN	Sidechain
2	C	985	GLU	Mainchain
3	D	1184	ASP	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1812	0	1839	111	0
1	B	1677	0	1703	127	0
1	G	1755	0	1773	105	0
1	H	1662	0	1687	111	0
2	C	10569	0	10587	439	0
2	I	10565	0	10581	437	0
3	D	9065	0	9210	430	0
3	J	9001	0	9165	461	0
4	E	691	0	695	26	0
4	K	627	0	634	26	0
5	F	3813	0	3880	150	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	L	3821	0	3884	149	0
6	D	2	0	0	0	0
6	I	1	0	0	0	0
6	J	1	0	0	0	0
7	D	2	0	0	0	0
7	J	2	0	0	0	0
All	All	55066	0	55638	2292	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 2292 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:I:1271:GLY:HA2	3:J:343:LEU:HD11	1.25	1.18
2:C:1269:ARG:HG2	3:D:343:LEU:HD12	1.24	1.11
2:C:696:ASP:HB2	2:C:798:GLN:HG2	1.35	1.08
3:D:1167:LYS:HZ3	3:D:1170:LYS:HB2	1.18	1.06
3:J:54:ASP:OD2	3:J:60:ARG:NH1	1.89	1.03

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	231/320 (72%)	201 (87%)	25 (11%)	5 (2%)	8	46
1	B	213/320 (67%)	186 (87%)	24 (11%)	3 (1%)	14	55
1	G	225/320 (70%)	193 (86%)	26 (12%)	6 (3%)	6	41
1	H	212/320 (66%)	189 (89%)	19 (9%)	4 (2%)	10	49
2	C	1338/1342 (100%)	1229 (92%)	99 (7%)	10 (1%)	26	70

*Continued on next page...*



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	I	1338/1342 (100%)	1227 (92%)	100 (8%)	11 (1%)	24	67
3	D	1163/1407 (83%)	1067 (92%)	86 (7%)	10 (1%)	21	65
3	J	1151/1407 (82%)	1054 (92%)	81 (7%)	16 (1%)	14	55
4	E	87/90 (97%)	82 (94%)	5 (6%)	0	100	100
4	K	77/90 (86%)	72 (94%)	3 (4%)	2 (3%)	7	42
5	F	462/613 (75%)	421 (91%)	36 (8%)	5 (1%)	17	61
5	L	463/613 (76%)	422 (91%)	39 (8%)	2 (0%)	39	79
All	All	6960/8184 (85%)	6343 (91%)	543 (8%)	74 (1%)	17	61

5 of 74 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	136	GLU
1	A	232	VAL
2	C	170	VAL
2	C	484	LEU
2	C	1137	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/279 (72%)	192 (96%)	9 (4%)	34	73
1	B	186/279 (67%)	173 (93%)	13 (7%)	19	58
1	G	193/279 (69%)	182 (94%)	11 (6%)	25	65
1	H	183/279 (66%)	172 (94%)	11 (6%)	24	64
2	C	1155/1157 (100%)	1066 (92%)	89 (8%)	16	53
2	I	1154/1157 (100%)	1063 (92%)	91 (8%)	15	52
3	D	962/1168 (82%)	885 (92%)	77 (8%)	15	51
3	J	960/1168 (82%)	881 (92%)	79 (8%)	14	50
4	E	72/74 (97%)	64 (89%)	8 (11%)	8	33

Continued on next page...



*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	K	67/74 (90%)	63 (94%)	4 (6%)	24	64
5	F	417/540 (77%)	381 (91%)	36 (9%)	13	48
5	L	418/540 (77%)	382 (91%)	36 (9%)	13	48
All	All	5968/6994 (85%)	5504 (92%)	464 (8%)	16	52

5 of 464 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
5	F	429	THR
2	I	116	ASP
5	L	98	VAL
5	F	476	ARG
1	G	50	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 89 such sidechains are listed below:

Mol	Chain	Res	Type
5	F	147	GLN
1	H	128	HIS
5	L	301	ASN
5	F	227	GLN
5	F	455	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.



## 5.6 Ligand geometry

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	233/320 (72%)	-0.03	4 (1%) 73 67	105, 144, 198, 236	0
1	B	217/320 (67%)	0.49	24 (11%) 7 7	121, 192, 243, 265	0
1	G	227/320 (70%)	0.25	10 (4%) 38 34	162, 195, 234, 271	0
1	H	216/320 (67%)	0.60	23 (10%) 8 7	175, 215, 240, 264	0
2	C	1340/1342 (99%)	0.07	55 (4%) 41 36	78, 133, 242, 297	0
2	I	1340/1342 (99%)	0.33	94 (7%) 19 18	104, 171, 269, 334	0
3	D	1167/1407 (82%)	0.08	31 (2%) 58 53	83, 120, 204, 261	0
3	J	1155/1407 (82%)	0.24	60 (5%) 31 28	99, 146, 228, 277	0
4	E	89/90 (98%)	-0.03	3 (3%) 49 44	118, 156, 179, 198	0
4	K	79/90 (87%)	1.22	18 (22%) 1 1	211, 251, 298, 305	0
5	F	468/613 (76%)	0.17	22 (4%) 35 32	115, 167, 295, 326	0
5	L	469/613 (76%)	0.20	26 (5%) 29 26	130, 181, 301, 320	0
All	All	7000/8184 (85%)	0.21	370 (5%) 30 27	78, 156, 252, 334	0

The worst 5 of 370 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	I	982	GLY	7.9
2	C	319	LEU	7.0
2	I	1000	LEU	6.8
2	I	1005	GLU	6.7
2	I	999	GLU	6.6

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	ZN	J	2003	1/1	0.88	0.22	0.34	189,189,189,189	0
7	ZN	D	2004	1/1	0.73	0.29	0.27	189,189,189,189	0
7	ZN	D	2003	1/1	0.70	0.20	-0.41	189,189,189,189	0
7	ZN	J	2002	1/1	0.94	0.13	-0.93	310,310,310,310	0
6	MG	D	2001	1/1	0.71	0.27	-	189,189,189,189	0
6	MG	J	2001	1/1	0.85	0.29	-	189,189,189,189	0
6	MG	D	2002	1/1	0.78	0.29	-	189,189,189,189	0
6	MG	I	1401	1/1	0.85	0.30	-	189,189,189,189	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.