



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 7, 2017 – 04:56 PM EST

PDB ID : 5UGS  
Title : Crystal structure of M. tuberculosis InhA inhibited by PT501  
Authors : Eltschkner, S.; Pschibul, A.; Spagnuolo, L.A.; Yu, W.; Tonge, P.J.; Kisker, C.  
Deposited on : 2017-01-10  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20028442  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20028442


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	<div> <div>86%</div> <div>6% • 7%</div> </div>
1	B	289	<div> <div>86%</div> <div>6% • 7%</div> </div>
1	C	289	<div> <div>89%</div> <div>• • 7%</div> </div>
1	D	289	<div> <div>3%</div> <div>86%</div> <div>6% • 7%</div> </div>
1	E	289	<div> <div>88%</div> <div>5% 7%</div> </div>
1	G	289	<div> <div>2%</div> <div>86%</div> <div>6% • 7%</div> </div>

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 12674 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-[acyl-carrier-protein] reductase [NADH].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total 2004	C 1270	N 348	O 375	S 11	0	2	0
1	B	268	Total 2005	C 1270	N 348	O 377	S 10	0	2	0
1	E	268	Total 1999	C 1266	N 348	O 375	S 10	0	1	0
1	G	268	Total 2004	C 1269	N 351	O 374	S 10	0	1	0
1	C	268	Total 1999	C 1266	N 348	O 375	S 10	0	1	0
1	D	268	Total 1996	C 1264	N 348	O 374	S 10	0	0	0

There are 120 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P9WGR1
A	-18	GLY	-	expression tag	UNP P9WGR1
A	-17	SER	-	expression tag	UNP P9WGR1
A	-16	SER	-	expression tag	UNP P9WGR1
A	-15	HIS	-	expression tag	UNP P9WGR1
A	-14	HIS	-	expression tag	UNP P9WGR1
A	-13	HIS	-	expression tag	UNP P9WGR1
A	-12	HIS	-	expression tag	UNP P9WGR1
A	-11	HIS	-	expression tag	UNP P9WGR1
A	-10	HIS	-	expression tag	UNP P9WGR1
A	-9	SER	-	expression tag	UNP P9WGR1
A	-8	SER	-	expression tag	UNP P9WGR1
A	-7	GLY	-	expression tag	UNP P9WGR1
A	-6	LEU	-	expression tag	UNP P9WGR1
A	-5	VAL	-	expression tag	UNP P9WGR1
A	-4	PRO	-	expression tag	UNP P9WGR1
A	-3	ARG	-	expression tag	UNP P9WGR1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P9WGR1
A	-1	SER	-	expression tag	UNP P9WGR1
A	0	HIS	-	expression tag	UNP P9WGR1
B	-19	MET	-	initiating methionine	UNP P9WGR1
B	-18	GLY	-	expression tag	UNP P9WGR1
B	-17	SER	-	expression tag	UNP P9WGR1
B	-16	SER	-	expression tag	UNP P9WGR1
B	-15	HIS	-	expression tag	UNP P9WGR1
B	-14	HIS	-	expression tag	UNP P9WGR1
B	-13	HIS	-	expression tag	UNP P9WGR1
B	-12	HIS	-	expression tag	UNP P9WGR1
B	-11	HIS	-	expression tag	UNP P9WGR1
B	-10	HIS	-	expression tag	UNP P9WGR1
B	-9	SER	-	expression tag	UNP P9WGR1
B	-8	SER	-	expression tag	UNP P9WGR1
B	-7	GLY	-	expression tag	UNP P9WGR1
B	-6	LEU	-	expression tag	UNP P9WGR1
B	-5	VAL	-	expression tag	UNP P9WGR1
B	-4	PRO	-	expression tag	UNP P9WGR1
B	-3	ARG	-	expression tag	UNP P9WGR1
B	-2	GLY	-	expression tag	UNP P9WGR1
B	-1	SER	-	expression tag	UNP P9WGR1
B	0	HIS	-	expression tag	UNP P9WGR1
E	-19	MET	-	initiating methionine	UNP P9WGR1
E	-18	GLY	-	expression tag	UNP P9WGR1
E	-17	SER	-	expression tag	UNP P9WGR1
E	-16	SER	-	expression tag	UNP P9WGR1
E	-15	HIS	-	expression tag	UNP P9WGR1
E	-14	HIS	-	expression tag	UNP P9WGR1
E	-13	HIS	-	expression tag	UNP P9WGR1
E	-12	HIS	-	expression tag	UNP P9WGR1
E	-11	HIS	-	expression tag	UNP P9WGR1
E	-10	HIS	-	expression tag	UNP P9WGR1
E	-9	SER	-	expression tag	UNP P9WGR1
E	-8	SER	-	expression tag	UNP P9WGR1
E	-7	GLY	-	expression tag	UNP P9WGR1
E	-6	LEU	-	expression tag	UNP P9WGR1
E	-5	VAL	-	expression tag	UNP P9WGR1
E	-4	PRO	-	expression tag	UNP P9WGR1
E	-3	ARG	-	expression tag	UNP P9WGR1
E	-2	GLY	-	expression tag	UNP P9WGR1
E	-1	SER	-	expression tag	UNP P9WGR1

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
E	0	HIS	-	expression tag	UNP P9WGR1
G	-19	MET	-	initiating methionine	UNP P9WGR1
G	-18	GLY	-	expression tag	UNP P9WGR1
G	-17	SER	-	expression tag	UNP P9WGR1
G	-16	SER	-	expression tag	UNP P9WGR1
G	-15	HIS	-	expression tag	UNP P9WGR1
G	-14	HIS	-	expression tag	UNP P9WGR1
G	-13	HIS	-	expression tag	UNP P9WGR1
G	-12	HIS	-	expression tag	UNP P9WGR1
G	-11	HIS	-	expression tag	UNP P9WGR1
G	-10	HIS	-	expression tag	UNP P9WGR1
G	-9	SER	-	expression tag	UNP P9WGR1
G	-8	SER	-	expression tag	UNP P9WGR1
G	-7	GLY	-	expression tag	UNP P9WGR1
G	-6	LEU	-	expression tag	UNP P9WGR1
G	-5	VAL	-	expression tag	UNP P9WGR1
G	-4	PRO	-	expression tag	UNP P9WGR1
G	-3	ARG	-	expression tag	UNP P9WGR1
G	-2	GLY	-	expression tag	UNP P9WGR1
G	-1	SER	-	expression tag	UNP P9WGR1
G	0	HIS	-	expression tag	UNP P9WGR1
C	-19	MET	-	initiating methionine	UNP P9WGR1
C	-18	GLY	-	expression tag	UNP P9WGR1
C	-17	SER	-	expression tag	UNP P9WGR1
C	-16	SER	-	expression tag	UNP P9WGR1
C	-15	HIS	-	expression tag	UNP P9WGR1
C	-14	HIS	-	expression tag	UNP P9WGR1
C	-13	HIS	-	expression tag	UNP P9WGR1
C	-12	HIS	-	expression tag	UNP P9WGR1
C	-11	HIS	-	expression tag	UNP P9WGR1
C	-10	HIS	-	expression tag	UNP P9WGR1
C	-9	SER	-	expression tag	UNP P9WGR1
C	-8	SER	-	expression tag	UNP P9WGR1
C	-7	GLY	-	expression tag	UNP P9WGR1
C	-6	LEU	-	expression tag	UNP P9WGR1
C	-5	VAL	-	expression tag	UNP P9WGR1
C	-4	PRO	-	expression tag	UNP P9WGR1
C	-3	ARG	-	expression tag	UNP P9WGR1
C	-2	GLY	-	expression tag	UNP P9WGR1
C	-1	SER	-	expression tag	UNP P9WGR1
C	0	HIS	-	expression tag	UNP P9WGR1
D	-19	MET	-	initiating methionine	UNP P9WGR1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-18	GLY	-	expression tag	UNP P9WGR1
D	-17	SER	-	expression tag	UNP P9WGR1
D	-16	SER	-	expression tag	UNP P9WGR1
D	-15	HIS	-	expression tag	UNP P9WGR1
D	-14	HIS	-	expression tag	UNP P9WGR1
D	-13	HIS	-	expression tag	UNP P9WGR1
D	-12	HIS	-	expression tag	UNP P9WGR1
D	-11	HIS	-	expression tag	UNP P9WGR1
D	-10	HIS	-	expression tag	UNP P9WGR1
D	-9	SER	-	expression tag	UNP P9WGR1
D	-8	SER	-	expression tag	UNP P9WGR1
D	-7	GLY	-	expression tag	UNP P9WGR1
D	-6	LEU	-	expression tag	UNP P9WGR1
D	-5	VAL	-	expression tag	UNP P9WGR1
D	-4	PRO	-	expression tag	UNP P9WGR1
D	-3	ARG	-	expression tag	UNP P9WGR1
D	-2	GLY	-	expression tag	UNP P9WGR1
D	-1	SER	-	expression tag	UNP P9WGR1
D	0	HIS	-	expression tag	UNP P9WGR1

- Molecule 2 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			44	21	7	14		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	E	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	G	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	C	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
2	D	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 3 is 5-[(4-cyclopropyl-1,2,3-triazol-1-yl)methyl]-2-(2-methylphenoxy)phenol (three-letter code: XT5) (formula: C<sub>19</sub>H<sub>19</sub>N<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O		0	0
			24	19	3	2			
3	A	1	Total	C	N	O		0	0
			24	19	3	2			
3	B	1	Total	C	N	O		0	0
			24	19	3	2			
3	E	1	Total	C	N	O		0	0
			24	19	3	2			
3	G	1	Total	C	N	O		0	0
			24	19	3	2			
3	C	1	Total	C	N	O		0	0
			24	19	3	2			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O	0	0
			24	19	3	2		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	G	2	Total	Cl	0	0
			2	2		
4	B	1	Total	Cl	0	0
			1	1		
4	A	3	Total	Cl	0	0
			3	3		
4	D	1	Total	Cl	0	0
			1	1		
4	C	3	Total	Cl	0	0
			3	3		

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Na	0	0
			1	1		

- Molecule 6 is water.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	43	Total	O	0	0
			43	43		
6	B	41	Total	O	0	0
			41	41		
6	E	31	Total	O	0	0
			31	31		
6	G	39	Total	O	0	0
			39	39		
6	C	38	Total	O	0	0
			38	38		
6	D	32	Total	O	0	0
			32	32		





### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]





- Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]



- Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]




- Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]





- Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]





● Molecule 1: Enoyl-[acyl-carrier-protein] reductase [NADH]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	91.92Å 130.36Å 176.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.30 – 2.80 46.30 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.30-2.80) 99.4 (46.30-2.80)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.87 (at 2.81Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, $R_{free}$	0.206 , 0.244 0.208 , 0.242	Depositor DCC
$R_{free}$ test set	2576 reflections (5.16%)	DCC
Wilson B-factor (Å <sup>2</sup> )	42.5	Xtriage
Anisotropy	0.185	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 31.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	12674	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.12% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, XT5, NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.81	1/2048 (0.0%)	0.82	7/2779 (0.3%)
1	B	0.84	2/2049 (0.1%)	0.89	8/2781 (0.3%)
1	C	0.80	1/2040 (0.0%)	0.84	6/2769 (0.2%)
1	D	0.82	3/2034 (0.1%)	0.79	3/2761 (0.1%)
1	E	0.87	4/2040 (0.2%)	0.79	1/2769 (0.0%)
1	G	0.79	0/2045	0.80	4/2775 (0.1%)
All	All	0.82	11/12256 (0.1%)	0.82	29/16634 (0.2%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	209	GLU	CD-OE1	8.93	1.35	1.25
1	E	219	GLU	CD-OE2	8.15	1.34	1.25
1	D	209	GLU	CD-OE2	7.81	1.34	1.25
1	E	210	GLU	CD-OE2	7.70	1.34	1.25
1	A	68	GLU	CD-OE2	7.00	1.33	1.25

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	218	LEU	CA-CB-CG	10.98	140.55	115.30
1	C	195	ARG	NE-CZ-NH2	9.89	125.24	120.30
1	A	195	ARG	NE-CZ-NH1	-9.31	115.64	120.30
1	B	214	GLN	CB-CA-C	8.74	127.88	110.40
1	B	225	ARG	NE-CZ-NH2	8.44	124.52	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2004	0	2027	16	0
1	B	2005	0	2024	9	0
1	C	1999	0	2018	2	1
1	D	1996	0	2013	9	0
1	E	1999	0	2018	10	0
1	G	2004	0	2026	10	0
2	A	44	0	26	0	0
2	B	44	0	26	1	0
2	C	44	0	26	0	0
2	D	44	0	26	0	0
2	E	44	0	26	1	0
2	G	44	0	26	0	0
3	A	48	0	0	2	0
3	B	24	0	0	1	0
3	C	24	0	0	0	0
3	D	24	0	0	1	0
3	E	24	0	0	0	0
3	G	24	0	0	0	0
4	A	3	0	0	1	0
4	B	1	0	0	0	0
4	C	3	0	0	1	0
4	D	1	0	0	0	0
4	G	2	0	0	0	0
5	B	1	0	0	0	0
6	A	43	0	0	2	0
6	B	41	0	0	0	0
6	C	38	0	0	0	0
6	D	32	0	0	1	0
6	E	31	0	0	0	0
6	G	39	0	0	0	0
All	All	12674	0	12282	54	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 54 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:216:GLN:NE2	1:G:220:GLU:OE1	1.83	1.11
1:A:67:ASN:ND2	1:A:70:HIS:CE1	2.53	0.76
1:B:211:ALA:O	1:B:215:ILE:HD12	1.92	0.69
1:A:67:ASN:CG	1:A:70:HIS:ND1	2.46	0.68
1:G:215:ILE:H	1:G:215:ILE:HD12	1.59	0.68

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:153:ARG:NH1	1:C:153:ARG:NH1[2_655]	1.87	0.33

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/289 (93%)	253 (94%)	14 (5%)	1 (0%)	39	74
1	B	268/289 (93%)	256 (96%)	11 (4%)	1 (0%)	39	74
1	C	267/289 (92%)	254 (95%)	12 (4%)	1 (0%)	39	74
1	D	266/289 (92%)	252 (95%)	13 (5%)	1 (0%)	39	74
1	E	267/289 (92%)	255 (96%)	11 (4%)	1 (0%)	39	74
1	G	267/289 (92%)	252 (94%)	14 (5%)	1 (0%)	39	74
All	All	1603/1734 (92%)	1522 (95%)	75 (5%)	6 (0%)	39	74

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	208	GLY
1	E	208	GLY
1	D	208	GLY
1	A	211	ALA

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	G	208	GLY

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/222 (93%)	203 (98%)	3 (2%)	72	93
1	B	206/222 (93%)	201 (98%)	5 (2%)	57	87
1	C	205/222 (92%)	202 (98%)	3 (2%)	72	93
1	D	204/222 (92%)	200 (98%)	4 (2%)	63	90
1	E	205/222 (92%)	201 (98%)	4 (2%)	63	90
1	G	205/222 (92%)	199 (97%)	6 (3%)	50	83
All	All	1231/1332 (92%)	1206 (98%)	25 (2%)	65	90

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	209	GLU
1	G	170	SER
1	D	199	MET
1	G	45	ARG
1	G	195[A]	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	139	ASN
1	D	70	HIS
1	C	106	ASN
1	A	139	ASN
1	C	139	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 24 ligands modelled in this entry, 11 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAD	A	301	-	42,48,48	0.93	2 (4%)	46,73,73	1.17	3 (6%)
3	XT5	A	302	-	26,27,27	3.49	8 (30%)	36,38,38	2.48	7 (19%)
3	XT5	A	303	-	26,27,27	2.80	7 (26%)	36,38,38	1.96	9 (25%)
2	NAD	B	301	-	42,48,48	0.86	2 (4%)	46,73,73	1.23	4 (8%)
3	XT5	B	302	-	26,27,27	2.43	6 (23%)	36,38,38	2.65	8 (22%)
2	NAD	C	301	-	42,48,48	1.08	3 (7%)	46,73,73	0.94	3 (6%)
3	XT5	C	302	-	26,27,27	3.60	8 (30%)	36,38,38	2.48	9 (25%)
2	NAD	D	301	-	42,48,48	0.96	2 (4%)	46,73,73	1.24	6 (13%)
3	XT5	D	302	-	26,27,27	3.72	9 (34%)	36,38,38	2.27	12 (33%)
2	NAD	E	301	-	42,48,48	1.00	2 (4%)	46,73,73	1.04	3 (6%)
3	XT5	E	302	-	26,27,27	4.05	10 (38%)	36,38,38	2.41	4 (11%)
2	NAD	G	301	-	42,48,48	1.00	2 (4%)	46,73,73	1.27	3 (6%)
3	XT5	G	302	-	26,27,27	3.58	8 (30%)	36,38,38	3.24	12 (33%)



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	301	-	-	1/22/62/62	0/5/5/5
3	XT5	A	302	-	-	1/12/14/14	0/3/4/4
3	XT5	A	303	-	-	0/12/14/14	0/3/4/4
2	NAD	B	301	-	-	0/22/62/62	0/5/5/5
3	XT5	B	302	-	-	3/12/14/14	0/3/4/4
2	NAD	C	301	-	-	0/22/62/62	0/5/5/5
3	XT5	C	302	-	-	2/12/14/14	0/3/4/4
2	NAD	D	301	-	-	0/22/62/62	0/5/5/5
3	XT5	D	302	-	-	2/12/14/14	0/3/4/4
2	NAD	E	301	-	-	0/22/62/62	0/5/5/5
3	XT5	E	302	-	-	1/12/14/14	0/3/4/4
2	NAD	G	301	-	-	0/22/62/62	0/5/5/5
3	XT5	G	302	-	-	3/12/14/14	0/3/4/4

The worst 5 of 69 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	302	XT5	NAO-NAX	-13.99	1.07	1.34
3	A	302	XT5	NAN-NAO	-13.75	1.14	1.34
3	G	302	XT5	NAN-NAO	-13.70	1.14	1.34
3	D	302	XT5	NAN-NAO	-12.52	1.16	1.34
3	C	302	XT5	NAN-NAO	-11.90	1.17	1.34

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	302	XT5	CAK-CAW-CAT	-10.05	110.36	120.12
3	G	302	XT5	CAK-CAW-CAT	-9.31	111.08	120.12
3	G	302	XT5	CAL-CAW-CAK	-6.52	53.40	58.73
3	C	302	XT5	CAK-CAW-CAT	-6.17	114.12	120.12
3	A	302	XT5	CAL-CAW-CAT	-5.89	114.40	120.12

There are no chirality outliers.

5 of 13 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	302	XT5	CAL-CAW-CAT-CAJ

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	302	XT5	CAK-CAW-CAT-CAJ
3	B	302	XT5	CAK-CAW-CAT-CAJ
3	D	302	XT5	CAK-CAW-CAT-NAN
3	B	302	XT5	CAL-CAW-CAT-NAN

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	303	XT5	2	0
2	B	301	NAD	1	0
3	B	302	XT5	1	0
3	D	302	XT5	1	0
2	E	301	NAD	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	268/289 (92%)	-0.28	3 (1%) 82 74	26, 37, 48, 56	0
1	B	268/289 (92%)	-0.08	4 (1%) 76 68	28, 39, 54, 59	0
1	C	268/289 (92%)	-0.19	3 (1%) 82 74	26, 38, 45, 51	0
1	D	268/289 (92%)	0.06	10 (3%) 45 33	27, 41, 52, 60	0
1	E	268/289 (92%)	-0.17	4 (1%) 76 68	26, 38, 48, 55	0
1	G	268/289 (92%)	-0.14	5 (1%) 70 59	26, 38, 52, 59	0
All	All	1608/1734 (92%)	-0.13	29 (1%) 71 61	26, 38, 50, 60	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	2	THR	5.8
1	C	2	THR	5.3
1	G	211	ALA	4.0
1	D	45	ARG	3.9
1	E	2	THR	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	CL	A	306	1/1	0.86	0.29	1.00	47,47,47,47	0
3	XT5	A	303	24/24	0.94	0.21	0.95	36,40,43,44	0
3	XT5	B	302	24/24	0.94	0.16	-0.05	33,36,40,41	0
3	XT5	A	302	24/24	0.97	0.14	-0.37	29,33,37,38	0
3	XT5	G	302	24/24	0.96	0.15	-0.39	30,33,37,38	0
3	XT5	C	302	24/24	0.97	0.14	-0.45	31,33,35,36	0
2	NAD	A	301	44/44	0.97	0.13	-0.55	30,35,38,39	0
2	NAD	B	301	44/44	0.96	0.14	-0.74	28,33,36,38	0
2	NAD	C	301	44/44	0.97	0.13	-0.77	29,34,38,38	0
5	NA	B	304	1/1	0.91	0.14	-0.78	33,33,33,33	0
4	CL	G	304	1/1	0.93	0.12	-0.85	48,48,48,48	0
3	XT5	E	302	24/24	0.98	0.14	-0.86	29,33,35,36	0
2	NAD	G	301	44/44	0.96	0.13	-0.94	25,32,36,36	0
2	NAD	E	301	44/44	0.97	0.13	-0.99	29,34,36,38	0
2	NAD	D	301	44/44	0.96	0.13	-1.33	26,32,36,39	0
3	XT5	D	302	24/24	0.96	0.13	-1.49	32,36,38,40	0
4	CL	D	303	1/1	0.99	0.11	-1.81	41,41,41,41	0
4	CL	C	303	1/1	0.94	0.08	-2.96	39,39,39,39	0
4	CL	A	304	1/1	0.98	0.05	-4.97	42,42,42,42	0
4	CL	B	303	1/1	0.99	0.22	-	40,40,40,40	0
4	CL	A	305	1/1	0.97	0.18	-	43,43,43,43	0
4	CL	C	304	1/1	0.97	0.18	-	37,37,37,37	0
4	CL	G	303	1/1	0.91	0.24	-	38,38,38,38	0
4	CL	C	305	1/1	0.95	0.10	-	36,36,36,36	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.