wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 05:16 PM BST

PDB ID	:	1UJO
Title	:	Solution Structure of the CH domain from Mouse Trangelin
Authors	:	Tomizawa, T.; Koshiba, S.; Kigawa, T.; Yokoyama, S.; RIKEN Structural Genomics/Proteomics Initiative (RSGI)
Deposited on	:	2003-08-08

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 94%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	144	35% 42% • 21%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:9-A:29, A:35-A:123, A:132-A:135 (114)	0.31	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13, 15, 16, 17, 18, 19, 20
2	2, 7
Single-model clusters	14

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2203 atoms, of which 1096 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Transgelin.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	144	2203	699	1096	188	213	7	0

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	CLONING ARTIFACT	UNP P37804
A	2	SER	-	CLONING ARTIFACT	UNP P37804
A	3	SER	-	CLONING ARTIFACT	UNP P37804
A	4	GLY	-	CLONING ARTIFACT	UNP P37804
A	5	SER	-	CLONING ARTIFACT	UNP P37804
A	6	SER	-	CLONING ARTIFACT	UNP P37804
A	7	GLY	-	CLONING ARTIFACT	UNP P37804
A	139	SER	-	CLONING ARTIFACT	UNP P37804
A	140	GLY	-	CLONING ARTIFACT	UNP P37804
A	141	PRO	-	CLONING ARTIFACT	UNP P37804
A	142	SER	-	CLONING ARTIFACT	UNP P37804
A	143	SER	-	CLONING ARTIFACT	UNP P37804
A	144	GLY	-	CLONING ARTIFACT	UNP P37804

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Transgelin

Chain A: 35% 42% • 21%

G1 S2 S3 G4 S5 S6 G7 E8 E9 L10 E11 E12 R13 L14 V15 E16 W17 I18 V19 V20 Q21 C22 G23 P24 D25 R28 P29 D30 R31 G32 R33 L34 G35 F36 Q37 V38 W39 L40 V44 I45 L46 L49 V50 N51 S52 L53 Y54 S58 K59 P60 V61 K62 V63 S69 M70

V71 M75 F81 L82 A85 E86 D87 Y88 G89 V90 I91 K92 T93 D94 N95 F96 Q97 T98 V99 D100 L101 Y102 E103 G104 K105 D106 M107 A108 A109 V110 Q111 R112 M115 G118 S119 L120 K124 N125 D126 G127 N128 Y129 R130 G131 D132 W135 F136 M137 K138 S139 G140 P141 S142

S143 G144

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

- Molecule 1: Transgelin

Chain A: 41% 35% • 21%

G1 S2 S3 G4 S5 S6 G7 E8 E9 L10 E11 L14 V15 E16 W17 I18 V19 V20 Q21 C22 G23 P24 D25 R28 P29 D30 R31 G32 R33 L34 G35 F36 Q37 V38 W39 L40 K41 N42 G43 V44 L53 Y54 P55 E56 K59 P60 V61 K62 S69 M70 K73 Q74 M75

V78 F81 L82 V90 I91 K92 T93 D94 N95 F96 Q97 T98 V99 D100 L101 Y102 E103 G104 K105 A109 V110 Q111 R112 M115 K124 N125 D126 G127 N128 Y129 R130 G131 W135 F136 M137 K138 S139 G140 P141 S142 S143 G144

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *torsion angle dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy, target function*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	1.0.7
CYANA	refinement	1.0.7

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 10068
Number of chemical shift lists	1
Total number of shifts	1744
Number of shifts mapped to atoms	1744
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	94%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	895	901	901	48±6
All	All	17900	18020	18020	951

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

5 of 241 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:82:LEU:HD21	1:A:96:PHE:CZ	0.93	1.98	11	17
1:A:91:ILE:HD12	1:A:93:THR:OG1	0.92	1.64	2	10
1:A:82:LEU:HD21	1:A:96:PHE:CE1	0.85	2.06	11	8
1:A:122:VAL:HG13	1:A:133:PRO:CB	0.82	2.04	13	4
1:A:15:VAL:HG13	1:A:39:TRP:CZ3	0.80	2.11	9	12

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	114/144 (79%)	80±3 (70±3%)	29±3 (26±3%)	5±1 (4±1%)	6 31
All	All	2280/2880 (79%)	1599 (70%)	582 (26%)	99 (4%)	6 31

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	53	LEU	17
1	A	60	PRO	14
1	A	89	GLY	11
1	A	54	TYR	9
1	A	104	GLY	8

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	99/121 (82%)	81±3 (82±3%)	18±3 (18±3%)	6 41
All	All	1980/2420 (82%)	1629 (82%)	351 (18%)	6 41

5 of 47 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	10	LEU	20
1	A	28	ARG	20
1	A	112	ARG	19
1	A	69	SER	17
1	A	115	MET	17

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 94% for the well-defined parts and 91% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 10068

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1744
Number of shifts mapped to atoms	1744
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	4

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	139	-0.36 ± 0.14	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	123	0.27 ± 0.08	None needed (< 0.5 ppm)
$^{13}C'$	126	-0.28 ± 0.17	None needed (< 0.5 ppm)
^{15}N	126	0.21 ± 0.38	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 94%, i.e. 1319 atoms were assigned a chemical shift out of a possible 1399. 0 out of 26 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	544/554 (98%)	219/220 (100%)	220/228 (96%)	105/106 (99%)
Sidechain	683/749 (91%)	420/438 (96%)	249/283 (88%)	14/28 (50%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	92/96 (96%)	48/50 (96%)	41/43 (95%)	3/3 (100%)
Overall	1319/1399 (94%)	687/708 (97%)	510/554 (92%)	122/137 (89%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	132	ASP	HB2	0.52	4.07 – 1.37	-8.1
1	A	29	PRO	HD2	0.74	5.45 – 1.85	-8.1
1	A	118	GLY	HA3	1.04	5.80 – 2.00	-7.5
1	A	132	ASP	H	4.06	11.17 – 5.47	-7.5

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: