



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:42 PM GMT

PDB ID : 3UOW
Title : Crystal Structure of PF10_0123, a GMP Synthetase from Plasmodium Falciparum
Authors : Wernimont, A.K.; Dong, A.; Hills, T.; Amani, M.; Perieteanu, A.; Lin, Y.H.; Loppnau, P.; Arrowsmith, C.H.; Edwards, A.M.; Bountra, C.; Weigelt, J.; Hui, R.; Structural Genomics Consortium (SGC)
Deposited on : 2011-11-17
Resolution : 2.72 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2439 (2.74-2.70)
Clashscore	102246	2771 (2.74-2.70)
Ramachandran outliers	100387	2726 (2.74-2.70)
Sidechain outliers	100360	2727 (2.74-2.70)
RSRZ outliers	91569	2443 (2.74-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	556	<div> <div>8%</div> <div>84%</div> <div>8%</div> <div>7%</div> </div>
1	B	556	<div> <div>9%</div> <div>78%</div> <div>7%</div> <div>14%</div> </div>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 7474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called GMP synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	517	Total	C	N	O	S	0	0	0
			3951	2562	638	734	17			
1	B	477	Total	C	N	O	S	0	0	0
			3442	2222	566	639	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	SEE REMARK 999	UNP Q8IJR9
B	0	GLY	-	SEE REMARK 999	UNP Q8IJR9

- Molecule 2 is XANTHOSINE-5'-MONOPHOSPHATE (three-letter code: XMP) (formula: $C_{10}H_{14}N_4O_9P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			24	10	4	9	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			24	10	4	9	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	21	Total	O	0	0
			21	21		
4	B	11	Total	O	0	0
			11	11		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GMP synthetase



• Molecule 1: GMP synthetase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	43.16 Å 67.76 Å 99.83 Å 103.07° 99.36° 94.33°	Depositor
Resolution (Å)	30.00 – 2.72 29.92 – 2.72	Depositor EDS
% Data completeness (in resolution range)	97.7 (30.00-2.72) 90.8 (29.92-2.72)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.29 (at 2.72 Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.259 , 0.296 0.256 , 0.295	Depositor DCC
R_{free} test set	1449 reflections (5.39%)	DCC
Wilson B-factor (Å ²)	61.4	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 28330 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7474	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.18% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, XMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.45	0/4036	0.44	0/5488
1	B	0.44	0/3507	0.43	0/4782
All	All	0.44	0/7543	0.44	0/10270

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3951	0	3629	26	0
1	B	3442	0	2968	15	0
2	A	24	0	12	2	0
2	B	24	0	12	1	0
3	A	1	0	0	0	0
4	A	21	0	0	0	0
4	B	11	0	0	0	0
All	All	7474	0	6621	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119:LEU:HD12	1:A:183:LEU:HD22	1.73	0.70
1:A:149:MET:HG3	1:A:150:ASN:N	2.09	0.67
1:A:149:MET:HG3	1:A:150:ASN:H	1.60	0.67
1:A:336:ARG:HH12	2:A:556:XMP:H7	1.43	0.67
1:A:99:MET:HE1	1:A:202:ILE:HD13	1.79	0.64
1:B:256:TYR:HB2	1:B:361:THR:HG22	1.83	0.61
1:A:336:ARG:NH1	2:A:556:XMP:H7	1.99	0.61
1:A:256:TYR:HB2	1:A:361:THR:HG22	1.86	0.57
1:B:224:PHE:HA	1:B:228:ILE:HD12	1.88	0.56
1:A:90:TYR:HA	1:A:93:GLN:HE21	1.72	0.54
1:B:332:PRO:HB3	1:B:555:GLU:HB3	1.88	0.54
1:B:365:GLN:HB2	1:B:404:GLU:HG2	1.89	0.54
1:B:90:TYR:HA	1:B:93:GLN:HE21	1.73	0.53
1:A:269:VAL:HG21	1:A:414:VAL:HG13	1.91	0.52
1:B:291:LEU:HD22	1:B:439:ALA:HB2	1.93	0.51
1:B:269:VAL:HG21	1:B:414:VAL:HG13	1.93	0.51
1:B:552:ILE:HD12	2:B:556:XMP:H5'1	1.92	0.50
1:A:443:ILE:HD12	1:A:555:GLU:HG3	1.93	0.49
1:B:94:GLU:O	1:B:98:GLN:HG2	2.13	0.48
1:A:464:ASP:OD1	1:A:529:ARG:NH1	2.47	0.47
1:A:435:GLY:N	1:A:436:PRO:HD2	2.29	0.47
1:A:224:PHE:HA	1:A:228:ILE:HD12	1.96	0.46
1:A:507:THR:HG23	1:A:513:ALA:HB2	1.98	0.46
1:A:508:SER:HB2	1:A:512:THR:HG23	1.98	0.46
1:B:435:GLY:N	1:B:436:PRO:HD2	2.32	0.45
1:B:423:LEU:HB3	1:B:427:ILE:HD11	1.99	0.45
1:A:257:VAL:HG22	1:A:362:PHE:HB2	1.99	0.45
1:A:187:SER:HB3	1:A:190:CYS:HB3	2.00	0.44
1:A:119:LEU:CD1	1:A:183:LEU:HD22	2.46	0.44
1:A:339:ILE:HG21	1:A:436:PRO:HA	1.99	0.43
1:A:364:LEU:HD23	1:A:364:LEU:C	2.38	0.43
1:B:275:HIS:CE1	1:B:311:MET:HA	2.55	0.42
1:A:538:ASN:HD22	1:A:538:ASN:C	2.23	0.42
1:A:245:LEU:O	1:A:248:ILE:HG22	2.20	0.42
1:A:549:PRO:HG3	1:B:531:LEU:HB3	2.02	0.41
1:B:116:VAL:HG12	1:B:187:SER:HB2	2.01	0.41
1:B:181:PHE:CZ	1:B:197:ASN:HB2	2.56	0.41
1:A:424:PRO:HB2	1:A:426:GLU:HG2	2.03	0.41
1:A:110:GLU:HB2	1:A:170:HIS:NE2	2.35	0.40
1:A:112:GLY:HA2	1:A:167:TRP:CZ3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	505/556 (91%)	485 (96%)	19 (4%)	1 (0%)	52	80
1	B	455/556 (82%)	442 (97%)	12 (3%)	1 (0%)	52	80
All	All	960/1112 (86%)	927 (97%)	31 (3%)	2 (0%)	52	80

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	179	GLU
1	B	179	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	386/506 (76%)	376 (97%)	10 (3%)	54	82
1	B	300/506 (59%)	288 (96%)	12 (4%)	38	68
All	All	686/1012 (68%)	664 (97%)	22 (3%)	46	76

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	165	THR
1	A	171	ASN
1	A	183	LEU
1	A	219	LEU
1	A	333	GLU
1	A	369	TYR
1	A	430	ARG
1	A	529	ARG
1	A	538	ASN
1	B	165	THR
1	B	183	LEU
1	B	201	ASN
1	B	219	LEU
1	B	241	HIS
1	B	314	THR
1	B	369	TYR
1	B	412	ASP
1	B	430	ARG
1	B	472	ASN
1	B	529	ARG
1	B	538	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	472	ASN
1	A	476	GLN
1	A	538	ASN
1	B	28	ASN
1	B	93	GLN
1	B	472	ASN
1	B	476	GLN
1	B	538	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	XMP	A	556	-	18,26,26	1.23	2 (11%)	22,40,40	2.25	4 (18%)
2	XMP	B	556	-	18,26,26	1.28	3 (16%)	22,40,40	2.29	4 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XMP	A	556	-	-	0/6/26/26	0/3/3/3
2	XMP	B	556	-	-	0/6/26/26	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	556	XMP	O4'-C1'	2.12	1.43	1.41
2	A	556	XMP	C5-C4	2.70	1.46	1.40
2	B	556	XMP	C5-C4	2.86	1.46	1.40
2	B	556	XMP	C6-C5	3.34	1.47	1.41
2	A	556	XMP	C6-C5	3.41	1.48	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	556	XMP	C5-C6-N1	-4.09	117.99	123.59
2	A	556	XMP	C5-C6-N1	-4.03	118.07	123.59
2	B	556	XMP	C6-C5-C4	-3.37	116.86	120.90
2	A	556	XMP	C6-C5-C4	-3.26	117.00	120.90
2	A	556	XMP	C4-C5-N7	-2.52	107.16	109.48
2	B	556	XMP	C4-C5-N7	-2.44	107.23	109.48
2	A	556	XMP	C6-N1-C2	7.93	122.10	115.25
2	B	556	XMP	C6-N1-C2	8.07	122.22	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	556	XMP	2	0
2	B	556	XMP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	517/556 (92%)	0.48	43 (8%)	14 12	33, 53, 69, 74	1 (0%)
1	B	477/556 (85%)	0.69	52 (10%)	7 6	41, 61, 82, 101	1 (0%)
All	All	994/1112 (89%)	0.58	95 (9%)	10 8	33, 57, 76, 101	2 (0%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	87	GLY	5.6
1	B	204	GLY	5.1
1	A	88	ILE	4.7
1	A	204	GLY	4.5
1	A	205	VAL	4.3
1	A	53	VAL	4.3
1	A	89	CYS	4.2
1	A	200	TYR	4.0
1	B	128	THR	3.9
1	A	54	ILE	3.9
1	A	500	CYS	3.8
1	A	91	GLY	3.8
1	A	47	ASP	3.7
1	B	113	CYS	3.6
1	B	32	PHE	3.6
1	A	487	VAL	3.5
1	A	212	TYR	3.4
1	A	400	PHE	3.3
1	B	142	LEU	3.3
1	A	57	GLY	3.3
1	B	22	ILE	3.3
1	A	86	PHE	3.3
1	A	55	LEU	3.2
1	B	177	ILE	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	48	MET	3.1
1	A	479	ALA	3.1
1	B	114	THR	3.1
1	B	46	LYS	3.1
1	B	129	TYR	3.0
1	B	60	TYR	3.0
1	A	130	CYS	2.9
1	B	487	VAL	2.9
1	A	501	VAL	2.8
1	B	205	VAL	2.8
1	A	232	LYS	2.8
1	B	479	ALA	2.8
1	A	207	TYR	2.8
1	B	165	THR	2.7
1	B	161	SER	2.7
1	B	130	CYS	2.7
1	B	207	TYR	2.7
1	B	263	GLY	2.6
1	A	206	GLN	2.6
1	A	195	ILE	2.6
1	A	203	TYR	2.6
1	B	354	ILE	2.6
1	A	48	MET	2.5
1	A	185	SER	2.5
1	B	47	ASP	2.5
1	B	76	GLU	2.5
1	B	189	ASN	2.5
1	B	541	LEU	2.5
1	A	92	MET	2.5
1	B	88	ILE	2.5
1	A	85	ILE	2.5
1	B	265	ILE	2.4
1	A	56	SER	2.4
1	B	400	PHE	2.4
1	A	58	GLY	2.3
1	B	87	GLY	2.3
1	B	360	LYS	2.3
1	A	511	MET	2.3
1	A	42	LEU	2.3
1	B	77	TYR	2.3
1	B	20	HIS	2.3
1	A	140	MET	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	143	TYR	2.2
1	B	200	TYR	2.2
1	B	126	ASN	2.2
1	A	240	TYR	2.2
1	B	167	TRP	2.2
1	B	358	ILE	2.2
1	B	468	TYR	2.2
1	B	175	THR	2.2
1	A	209	PRO	2.1
1	B	75	PHE	2.1
1	A	496	TYR	2.1
1	A	502	LEU	2.1
1	B	267	SER	2.1
1	B	502	LEU	2.1
1	B	500	CYS	2.1
1	A	222	TYR	2.1
1	A	480	VAL	2.1
1	A	226	TYR	2.1
1	A	398	LEU	2.0
1	B	61	SER	2.0
1	B	144	SER	2.0
1	B	368	LEU	2.0
1	B	124	ILE	2.0
1	B	56	SER	2.0
1	B	183	LEU	2.0
1	B	212	TYR	2.0
1	B	268	THR	2.0
1	B	501	VAL	2.0
1	B	180	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	XMP	B	556	24/24	0.93	0.14	-1.85	45,48,50,51	0
2	XMP	A	556	24/24	0.95	0.13	-1.90	37,38,39,40	0
3	CA	A	557	1/1	0.69	0.19	-	124,124,124,124	0

6.5 Other polymers [i](#)

There are no such residues in this entry.