



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:01 PM GMT

PDB ID : 4UO4
Title : Structure of the A_Canine_Colorado_17864_06 H3 haemagglutinin
Authors : Vachieri, S.G.; Collins, P.J.; Haire, L.F.; Ogradowicz, R.W.; Martin, S.R.;
Walker, P.A.; Xiong, X.; Gamblin, S.J.; Skehel, J.J.
Deposited on : 2014-05-31
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	
2	B	175	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	BMA	A	633	X	-	-	-
5	MAN	A	637	X	-	-	-
6	NAG	A	641	-	-	-	X

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 4098 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H3 HAEMAGGLUTININ HA1 CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	319	Total	C	N	O	S	0	0	0
			2473	1543	437	477	16			

- Molecule 2 is a protein called H3 HAEMAGGLUTININ HA2 CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	175	Total	C	N	O	S	0	0	0
			1416	881	246	283	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	173	SER	-	EXPRESSION TAG	UNP E0UVR5
B	174	GLY	-	EXPRESSION TAG	UNP E0UVR5
B	175	ARG	-	EXPRESSION TAG	UNP E0UVR5
B	131	GLU	ASP	CONFLICT	UNP E0UVR5

- Molecule 3 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	2	Total	C	N	O	0	0
			28	16	2	10		
4	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 5 is a polymer of unknown type called SUGAR (4-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	4	Total	C	N	O	0	0
			50	28	2	20		

- Molecule 6 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	5	Total	C	N	O	0	0
			61	34	2	25		

- Molecule 7 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	S	0	0
			5	4	1		
7	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	16	Total	O	0	0
			16	16		
8	B	2	Total	O	0	0
			2	2		

4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, α , β , γ	96.41Å 96.41Å 347.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	173.56 – 2.60 47.74 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.3 (173.56-2.60) 99.3 (47.74-2.60)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.33 (at 2.61Å)	Xtriage
Refinement program	REFMAC 5.8.0069	Depositor
R, R_{free}	0.199 , 0.244 0.204 , 0.246	Depositor DCC
R_{free} test set	1529 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	59.0	Xtriage
Anisotropy	0.685	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 37.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	2 of 30349 reflections (0.007%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4098	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, BMA, NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/2524	0.75	1/3424 (0.0%)
2	B	0.58	0/1441	0.83	4/1939 (0.2%)
All	All	0.55	0/3965	0.78	5/5363 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
5	A	2	0

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	110	LEU	CA-CB-CG	7.21	131.87	115.30
2	B	132	ASP	CB-CG-OD1	5.81	123.53	118.30
2	B	124	ARG	NE-CZ-NH2	-5.20	117.70	120.30
2	B	153	ARG	NE-CZ-NH1	5.17	122.88	120.30
1	A	109	ARG	NE-CZ-NH1	5.01	122.81	120.30

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	633	BMA	C1
5	A	637	MAN	C1

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2473	0	2421	23	0
2	B	1416	0	1324	21	0
3	A	14	0	13	0	0
4	A	56	0	50	0	0
5	A	50	0	43	0	0
6	A	61	0	52	0	0
7	A	5	0	0	1	0
7	B	5	0	0	0	0
8	A	16	0	0	0	0
8	B	2	0	0	0	0
All	All	4098	0	3903	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:56:ILE:HD12	2:B:56:ILE:O	1.79	0.82
2:B:51:LYS:HE2	2:B:107:THR:OG1	1.83	0.78
1:A:17:HIS:HB3	2:B:115:MET:HE1	1.69	0.73
1:A:191:GLN:HE22	1:A:250:ASN:HD21	1.42	0.68
1:A:321:ARG:NH1	7:A:1327:SO4:O2	2.26	0.68
2:B:107:THR:HA	2:B:110:LEU:HD13	1.76	0.68
1:A:155:THR:HG21	1:A:194:LEU:HD12	1.79	0.64
2:B:25:ARG:HE	2:B:34:GLN:HE21	1.48	0.60
1:A:132:GLN:HE21	1:A:152:ASN:HD21	1.50	0.59
1:A:131:THR:HG22	1:A:156:LYS:O	2.04	0.58
1:A:17:HIS:HB3	2:B:115:MET:CE	2.35	0.56
1:A:295:GLN:HE22	1:A:298:ASN:H	1.54	0.56
2:B:158:ASP:HB3	2:B:161:ILE:HD12	1.88	0.55
1:A:15:LEU:HD22	2:B:119:PHE:HA	1.89	0.53
2:B:19:ASP:HB3	2:B:36:ALA:HB2	1.89	0.53
2:B:110:LEU:HD22	2:B:111:THR:HG23	1.91	0.52
1:A:276:ILE:O	1:A:276:ILE:HG22	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:174:GLY:O	2:B:175:ARG:HB2	2.10	0.51
1:A:131:THR:HG23	1:A:155:THR:OG1	2.12	0.49
2:B:174:GLY:O	2:B:175:ARG:CB	2.60	0.49
1:A:133:ASN:HD22	1:A:255:ARG:HH22	1.60	0.49
2:B:128:GLU:O	2:B:170:ARG:NH1	2.46	0.48
1:A:27:LYS:NZ	2:B:97:GLU:OE2	2.46	0.48
2:B:98:LEU:HG	2:B:102:LEU:HD22	1.94	0.48
2:B:171:PHE:C	2:B:173:SER:H	2.18	0.46
1:A:187:SER:HB2	1:A:189:GLN:OE1	2.16	0.46
2:B:51:LYS:HE2	2:B:107:THR:HG1	1.79	0.45
1:A:54:LYS:HB3	1:A:55:SER:HA	1.97	0.45
1:A:182:ILE:HD11	1:A:215:PRO:HD3	2.00	0.44
2:B:52:LEU:HA	2:B:52:LEU:HD12	1.74	0.44
1:A:249:GLY:C	1:A:250:ASN:HD22	2.22	0.43
2:B:53:ASN:HD22	2:B:53:ASN:N	2.19	0.41
1:A:80:GLN:O	1:A:81:TYR:HB2	2.21	0.41
1:A:164:LEU:O	1:A:246:ASN:HA	2.20	0.41
1:A:127:TRP:CE3	1:A:166:VAL:HG21	2.56	0.40
1:A:17:HIS:CD2	2:B:115:MET:CE	3.04	0.40
1:A:17:HIS:CB	2:B:115:MET:HE1	2.45	0.40
1:A:180:TRP:CE2	1:A:204:VAL:HG21	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	317/328 (97%)	305 (96%)	12 (4%)	0	100	100
2	B	173/175 (99%)	162 (94%)	9 (5%)	2 (1%)	16	33
All	All	490/503 (97%)	467 (95%)	21 (4%)	2 (0%)	39	65

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	59	THR
2	B	172	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/289 (97%)	260 (93%)	20 (7%)	18	36
2	B	146/148 (99%)	134 (92%)	12 (8%)	14	27
All	All	426/437 (98%)	394 (92%)	32 (8%)	17	33

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	HIS
1	A	24	THR
1	A	25	LEU
1	A	40	THR
1	A	41	GLU
1	A	42	LEU
1	A	47	SER
1	A	53	ASN
1	A	70	MET
1	A	96	ASN
1	A	143	SER
1	A	151	LEU
1	A	189	GLN
1	A	193	LYS
1	A	194	LEU
1	A	201	ARG
1	A	260	LEU
1	A	264	LYS
1	A	295	GLN
1	A	325	GLU
2	B	53	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	58	ARG
2	B	72	GLU
2	B	73	VAL
2	B	86	ASP
2	B	102	LEU
2	B	110	LEU
2	B	120	GLU
2	B	126	LEU
2	B	127	ARG
2	B	150	GLU
2	B	167	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	53	ASN
1	A	96	ASN
1	A	132	GLN
1	A	133	ASN
1	A	250	ASN
1	A	295	GLN
1	A	312	ASN
2	B	34	GLN
2	B	42	GLN
2	B	47	GLN
2	B	53	ASN
2	B	78	GLN
2	B	125	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

13 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	A	611	1,4	14,14,15	0.74	0	15,19,21	1.05	1 (6%)
4	NAG	A	612	4	14,14,15	0.49	0	15,19,21	1.25	1 (6%)
4	NAG	A	621	1,4	14,14,15	0.73	0	15,19,21	1.90	3 (20%)
4	NAG	A	622	4	14,14,15	0.57	0	15,19,21	1.15	1 (6%)
5	NAG	A	631	1,5	14,14,15	0.66	0	15,19,21	1.33	4 (26%)
5	NAG	A	632	5	14,14,15	0.72	0	15,19,21	1.50	1 (6%)
5	BMA	A	633	5	11,11,12	0.49	0	14,15,17	2.49	6 (42%)
5	MAN	A	637	5	11,11,12	0.81	0	14,15,17	2.17	2 (14%)
6	NAG	A	641	1,6	14,14,15	0.93	1 (7%)	15,19,21	1.68	2 (13%)
6	NAG	A	642	6	14,14,15	0.65	0	15,19,21	1.57	2 (13%)
6	BMA	A	643	6	11,11,12	0.58	0	14,15,17	1.11	1 (7%)
6	MAN	A	644	6	11,11,12	0.78	0	14,15,17	2.07	4 (28%)
6	MAN	A	645	6	11,11,12	0.71	0	14,15,17	1.09	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	611	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	612	4	-	0/6/23/26	0/1/1/1
4	NAG	A	621	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	622	4	-	0/6/23/26	0/1/1/1
5	NAG	A	631	1,5	-	0/6/23/26	0/1/1/1
5	NAG	A	632	5	-	0/6/23/26	0/1/1/1
5	BMA	A	633	5	1/1/4/5	0/2/19/22	0/1/1/1
5	MAN	A	637	5	1/1/4/5	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	641	1,6	-	0/6/23/26	0/1/1/1
6	NAG	A	642	6	-	0/6/23/26	0/1/1/1
6	BMA	A	643	6	-	0/2/19/22	0/1/1/1
6	MAN	A	644	6	-	0/2/19/22	0/1/1/1
6	MAN	A	645	6	-	0/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	641	NAG	C1-C2	2.11	1.55	1.52

All (29) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	641	NAG	O4-C4-C3	-4.55	100.09	110.34
5	A	632	NAG	C3-C4-C5	-3.37	104.33	110.20
6	A	644	MAN	C3-C4-C5	-3.15	104.71	110.20
5	A	633	BMA	C2-C3-C4	-3.04	105.87	111.04
4	A	621	NAG	O5-C5-C6	-2.91	101.06	107.35
6	A	645	MAN	O5-C1-C2	-2.75	106.39	110.86
6	A	641	NAG	O7-C7-C8	-2.69	117.13	122.06
4	A	612	NAG	C4-C3-C2	-2.42	107.47	111.23
5	A	633	BMA	O2-C2-C1	-2.12	104.96	109.21
6	A	643	BMA	C1-C2-C3	-2.11	107.04	109.54
5	A	633	BMA	C6-C5-C4	-2.09	107.85	113.02
5	A	631	NAG	O7-C7-C8	-2.03	118.34	122.06
5	A	631	NAG	O3-C3-C4	-2.02	105.78	110.34
5	A	631	NAG	C3-C2-N2	2.06	115.49	110.56
4	A	621	NAG	C4-C3-C2	2.19	114.63	111.23
5	A	633	BMA	C1-C2-C3	2.19	112.13	109.54
5	A	637	MAN	C1-O5-C5	2.50	115.42	112.25
6	A	644	MAN	C1-O5-C5	2.67	115.64	112.25
6	A	642	NAG	C2-N2-C7	2.75	126.57	123.04
4	A	611	NAG	C1-O5-C5	2.75	115.74	112.25
5	A	631	NAG	C8-C7-N2	2.96	121.78	116.11
4	A	622	NAG	O5-C5-C6	3.08	114.01	107.35
6	A	644	MAN	O3-C3-C4	3.19	117.51	110.34
6	A	644	MAN	O3-C3-C2	3.76	116.79	110.00
6	A	642	NAG	C3-C4-C5	4.01	117.18	110.20
5	A	633	BMA	C1-O5-C5	4.88	118.44	112.25
4	A	621	NAG	C1-O5-C5	5.18	118.83	112.25
5	A	633	BMA	O5-C1-C2	5.75	120.18	110.86

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	637	MAN	O5-C1-C2	7.34	122.76	110.86

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	A	633	BMA	C1
5	A	637	MAN	C1

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.6 Ligand geometry

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	SO4	A	1327	-	4,4,4	0.48	0	6,6,6	0.22	0
3	NAG	A	601	1	14,14,15	0.82	0	15,19,21	1.45	3 (20%)
7	SO4	B	1176	-	4,4,4	0.71	0	6,6,6	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	SO4	A	1327	-	-	0/0/0/0	0/0/0/0
3	NAG	A	601	1	-	0/6/23/26	0/1/1/1
7	SO4	B	1176	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	601	NAG	O4-C4-C5	2.10	114.80	109.24
3	A	601	NAG	O3-C3-C2	2.27	113.60	109.11
3	A	601	NAG	C4-C3-C2	3.57	116.78	111.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	1327	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	319/328 (97%)	-0.37	1 (0%) 94 93	48, 70, 100, 124	0
2	B	175/175 (100%)	-0.13	0 100 100	48, 73, 100, 119	0
All	All	494/503 (98%)	-0.29	1 (0%) 95 95	48, 71, 100, 124	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	143	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	NAG	A	641	14/15	0.92	0.18	3.05	83,96,101,109	0
6	MAN	A	645	11/12	0.86	0.16	0.80	107,131,151,155	0
5	NAG	A	632	14/15	0.97	0.10	-1.44	80,92,105,107	0
4	NAG	A	621	14/15	0.89	0.42	-	100,121,131,139	0
6	NAG	A	642	14/15	0.91	0.33	-	103,127,136,142	0
4	NAG	A	612	14/15	0.89	0.30	-	132,140,148,149	0
6	BMA	A	643	11/12	0.87	0.32	-	124,134,138,140	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	NAG	A	622	14/15	0.90	0.50	-	121,135,147,147	0
4	NAG	A	611	14/15	0.93	0.15	-	94,105,117,133	0
5	BMA	A	633	11/12	0.84	0.16	-	112,124,132,132	0
5	MAN	A	637	11/12	0.78	0.22	-	129,138,146,148	0
6	MAN	A	644	11/12	0.84	0.32	-	119,133,137,142	0
5	NAG	A	631	14/15	0.97	0.11	-	84,93,97,98	0

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
7	SO4	B	1176	5/5	0.97	0.17	-0.34	75,78,81,91	0
7	SO4	A	1327	5/5	0.92	0.11	-	121,123,135,137	0
3	NAG	A	601	14/15	0.80	0.29	-	116,127,135,136	0

6.5 Other polymers [i](#)

There are no such residues in this entry.