



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 09:04 PM GMT

PDB ID : 4UPR
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with N,N''-{[(2S)-3-aminopropane-1,2-diyl]bis(oxymethanediylbenzene-3,1-diyl)}dithiophene-2-carboximidamide
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-06-17
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2910 (1.96-1.92)
Clashscore	102246	3095 (1.96-1.92)
Ramachandran outliers	100387	3062 (1.96-1.92)
Sidechain outliers	100360	3062 (1.96-1.92)
RSRZ outliers	91569	2915 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	<div> <div>12%</div> <div>83%</div> <div>7% • 9%</div> </div>
1	B	443	<div> <div>10%</div> <div>83%</div> <div>7% • 9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	7LN	B	800	-	-	-	X
4	ACT	B	860	-	-	-	X
5	GOL	A	880	-	-	-	X
5	GOL	B	880	-	-	-	X

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 7014 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	405	Total	As	C	N	O	S	0	0	0
			3223	1	2049	568	589	16			
1	B	403	Total	As	C	N	O	S	0	0	0
			3212	1	2042	566	587	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	SEE REMARK 999	UNP P29473
B	100	ARG	CYS	SEE REMARK 999	UNP P29473

- Molecule 2 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			17	9	5	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 3 is N,N''-{[(2S)-3-AMINOPROPANE-1,2-DIYL]BIS(OXYMETHANEDIYLBENZENE-3,1-DIYL)}DITHIOPHENE-2-CARBOXIMIDAMIDE (three-letter code: 7LN) (formula: C₂₇H₂₉N₅O₂S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			36	27	5	2	2		
3	B	1	Total	C	N	O	S	0	0
			36	27	5	2	2		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
6	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Zn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	200	Total	O	0	0
			200	200		
8	B	158	Total	O	0	0
			158	158		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.16Å 106.73Å 157.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.35 – 1.93 39.32 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.4 (39.35-1.93) 99.4 (39.32-1.93)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.171 , 0.207 0.170 , 0.206	Depositor DCC
R_{free} test set	3710 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.9	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 74162 reflections	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7014	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, H4B, CAS, 7LN, ACT, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.55	0/3303	0.66	2/4497 (0.0%)
1	B	0.53	0/3291	0.65	2/4480 (0.0%)
All	All	0.54	0/6594	0.66	4/8977 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	240	ARG	NE-CZ-NH2	-7.60	116.50	120.30
1	B	240	ARG	NE-CZ-NH1	6.39	123.49	120.30
1	A	240	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	A	294	LEU	CA-CB-CG	5.23	127.32	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3223	0	3127	25	0
1	B	3212	0	3117	20	1
2	A	17	0	15	1	0
2	B	17	0	15	1	0
3	A	36	0	28	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	36	0	28	0	0
4	A	8	0	6	0	0
4	B	8	0	6	0	0
5	A	6	0	8	0	0
5	B	6	0	8	0	0
6	A	43	0	30	1	0
6	B	43	0	30	1	0
7	A	1	0	0	0	0
8	A	200	0	0	2	1
8	B	158	0	0	1	0
All	All	7014	0	6418	44	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:384:CAS:AS	1:B:384:CAS:SG	2.49	1.30
1:A:384:CAS:SG	1:A:384:CAS:AS	2.53	1.26
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.59	0.84
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.73	0.70
1:A:236:ARG:NH1	1:A:236:ARG:HB2	2.07	0.70
1:A:240:ARG:HD3	1:A:298:PRO:CB	2.26	0.65
1:A:384:CAS:CB	1:A:384:CAS:AS	3.05	0.64
1:A:479:PRO:HA	3:A:800:7LN:H23	1.78	0.64
1:A:236:ARG:HH11	1:A:236:ARG:HB2	1.64	0.60
1:B:236:ARG:HD2	1:B:242:ASP:OD1	2.03	0.57
1:A:236:ARG:CB	1:A:236:ARG:HH11	2.17	0.57
1:A:378:LEU:HB2	8:A:2146:HOH:O	2.08	0.54
1:B:130:ARG:NH2	1:B:156:GLU:OE1	2.40	0.54
1:B:384:CAS:AS	1:B:384:CAS:CB	3.17	0.53
1:A:449:TRP:HA	2:A:600:H4B:N1	2.25	0.52
1:B:378:LEU:HB2	8:B:2126:HOH:O	2.09	0.51
1:A:384:CAS:SG	1:A:384:CAS:CE2	2.98	0.51
1:B:240:ARG:HD2	1:B:241:GLY:O	2.10	0.51
6:B:1483:HEM:HHC	6:B:1483:HEM:HBB2	1.92	0.50
1:A:384:CAS:AS	1:A:384:CAS:HB2	2.72	0.50
1:B:71:PRO:HG2	1:B:84:ASP:HB3	1.93	0.50
1:A:240:ARG:HD2	1:A:241:GLY:O	2.12	0.49
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.94	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:99:ARG:HG2	1:B:100:ARG:HD2	1.95	0.48
1:A:340:ASN:HD22	1:A:340:ASN:H	1.62	0.48
1:A:109:ARG:HB2	1:A:109:ARG:NH1	2.29	0.47
1:A:149:GLU:O	1:A:153:GLN:HG3	2.15	0.46
1:B:384:CAS:CE2	1:B:384:CAS:SG	3.04	0.46
1:A:392:THR:HB	1:B:423:HIS:HB2	1.98	0.45
1:B:281:TRP:HB2	1:B:304:LEU:HD21	1.98	0.45
8:A:2001:HOH:O	1:B:109:ARG:HG2	2.18	0.44
1:A:281:TRP:HB2	1:A:304:LEU:HD21	1.98	0.44
1:A:109:ARG:HB2	1:A:109:ARG:HH11	1.83	0.44
1:B:231:THR:O	1:B:353:ALA:HA	2.17	0.44
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.58	0.43
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.59	0.42
6:A:1483:HEM:HBB2	6:A:1483:HEM:HHC	2.01	0.42
1:A:423:HIS:HB2	1:B:392:THR:HB	2.01	0.42
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.49	0.42
1:A:83:TYR:HE1	1:B:109:ARG:HH21	1.69	0.41
1:A:240:ARG:HD3	1:A:298:PRO:CG	2.51	0.41
1:B:449:TRP:HA	2:B:600:H4B:N1	2.36	0.40
1:B:455:SER:HB3	1:B:458:LEU:HD22	2.01	0.40
1:A:363:GLU:OE1	3:A:800:7LN:N07	2.55	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:240:ARG:NH2	8:A:2067:HOH:O[2_555]	2.06	0.14

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/443 (90%)	389 (97%)	10 (2%)	1 (0%)	46	35
1	B	398/443 (90%)	384 (96%)	14 (4%)	0	100	100
All	All	798/886 (90%)	773 (97%)	24 (3%)	1 (0%)	56	47

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/375 (92%)	324 (94%)	19 (6%)	27	11
1	B	342/375 (91%)	329 (96%)	13 (4%)	40	24
All	All	685/750 (91%)	653 (95%)	32 (5%)	32	16

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	72	ARG
1	A	91	GLN
1	A	92	GLN
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	109	ARG
1	A	125	LEU
1	A	146	GLN
1	A	154	GLU
1	A	293	LEU
1	A	294	LEU
1	A	311	LEU
1	A	328	LEU
1	A	340	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	378	LEU
1	A	438	LYS
1	A	458	LEU
1	A	468	ASN
1	B	69	LYS
1	B	81	ILE
1	B	100	ARG
1	B	105	LEU
1	B	109	ARG
1	B	141	LYS
1	B	225	ASN
1	B	240	ARG
1	B	330	LEU
1	B	340	ASN
1	B	378	LEU
1	B	379	GLU
1	B	458	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	A	166	HIS
1	A	191	GLN
1	A	340	ASN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	384	1	5,8,9	0.98	0	2,9,11	1.37	0
1	CAS	B	384	1	5,8,9	1.25	0	2,9,11	1.64	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	0/0/0/0
1	CAS	B	384	1	-	0/0/7/9	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	384	CAS	O-C-CA	-2.30	119.50	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	384	CAS	4	0
1	B	384	CAS	3	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	HEM	A	1483	1	30,50,50	2.01	8 (26%)	24,82,82	2.35	9 (37%)
2	H4B	A	600	-	13,18,18	0.75	0	11,26,26	2.51	5 (45%)
3	7LN	A	800	-	38,39,39	1.27	7 (18%)	36,51,51	1.58	4 (11%)
4	ACT	A	860	-	1,3,3	1.19	0	0,3,3	0.00	-
4	ACT	A	861	-	1,3,3	1.32	0	0,3,3	0.00	-
5	GOL	A	880	-	5,5,5	0.33	0	5,5,5	0.46	0
6	HEM	B	1483	1	30,50,50	2.12	6 (20%)	24,82,82	2.36	9 (37%)
2	H4B	B	600	-	13,18,18	0.76	0	11,26,26	2.36	5 (45%)
3	7LN	B	800	-	38,39,39	1.28	6 (15%)	36,51,51	1.64	3 (8%)
4	ACT	B	860	-	1,3,3	1.57	0	0,3,3	0.00	-
4	ACT	B	861	-	1,3,3	1.25	0	0,3,3	0.00	-
5	GOL	B	880	-	5,5,5	0.29	0	5,5,5	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	HEM	A	1483	1	-	0/10/54/54	0/0/8/8
2	H4B	A	600	-	-	0/8/17/17	0/2/2/2
3	7LN	A	800	-	-	1/21/29/29	0/4/4/4
4	ACT	A	860	-	-	0/0/0/0	0/0/0/0
4	ACT	A	861	-	-	0/0/0/0	0/0/0/0
5	GOL	A	880	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	HEM	B	1483	1	-	0/10/54/54	0/0/8/8
2	H4B	B	600	-	-	0/8/17/17	0/2/2/2
3	7LN	B	800	-	-	1/21/29/29	0/4/4/4
4	ACT	B	860	-	-	0/0/0/0	0/0/0/0
4	ACT	B	861	-	-	0/0/0/0	0/0/0/0
5	GOL	B	880	-	-	0/4/4/4	0/0/0/0

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1483	HEM	C2D-C3D	-6.60	1.34	1.54
6	B	1483	HEM	C2D-C3D	-6.43	1.35	1.54
6	B	1483	HEM	C2C-C1C	-5.96	1.41	1.52
6	A	1483	HEM	C2C-C1C	-5.40	1.42	1.52
6	B	1483	HEM	C3B-C4B	-3.66	1.48	1.51
3	B	800	7LN	C11-N07	-3.21	1.37	1.42
3	A	800	7LN	C31-N27	-2.82	1.36	1.41
3	B	800	7LN	C31-N27	-2.62	1.36	1.41
6	B	1483	HEM	C2B-C1B	-2.53	1.43	1.51
3	A	800	7LN	C11-N07	-2.45	1.38	1.42
3	A	800	7LN	C26-N27	-2.43	1.34	1.39
3	B	800	7LN	C26-N27	-2.41	1.34	1.39
6	A	1483	HEM	C2B-C1B	-2.27	1.44	1.51
6	A	1483	HEM	C3D-C4D	-2.19	1.48	1.51
6	A	1483	HEM	CHC-C1C	2.03	1.41	1.36
3	A	800	7LN	C05-S01	2.18	1.75	1.72
3	B	800	7LN	C06-N07	2.33	1.33	1.30
6	A	1483	HEM	FE-NB	2.34	2.09	1.97
6	B	1483	HEM	FE-NB	2.36	2.10	1.97
6	A	1483	HEM	C1C-NC	2.38	1.38	1.36
3	A	800	7LN	C06-N07	2.56	1.34	1.30
6	A	1483	HEM	FE-NC	2.81	2.06	1.95
3	B	800	7LN	C05-C06	2.92	1.49	1.45
3	A	800	7LN	C25-C26	2.93	1.49	1.45
3	A	800	7LN	C05-C06	2.99	1.49	1.45
6	B	1483	HEM	FE-NC	3.22	2.08	1.95
3	B	800	7LN	C25-C26	3.29	1.50	1.45

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	800	7LN	C23-C22-S21	-6.12	106.45	113.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	800	7LN	C03-C02-S01	-5.77	106.84	113.23
3	B	800	7LN	C23-C22-S21	-5.70	106.92	113.23
3	A	800	7LN	C03-C02-S01	-4.72	108.00	113.23
6	B	1483	HEM	CBA-CAA-C2A	-3.97	105.41	112.53
2	B	600	H4B	N3-C2-N1	-3.87	119.19	125.53
6	B	1483	HEM	C1D-CHD-C4C	-3.56	119.88	125.82
6	A	1483	HEM	CBA-CAA-C2A	-3.41	106.42	112.53
6	A	1483	HEM	C1D-CHD-C4C	-3.31	120.29	125.82
2	A	600	H4B	N3-C2-N1	-3.15	120.37	125.53
6	B	1483	HEM	C3C-CAC-CBC	-2.41	120.76	124.46
3	A	800	7LN	C21-C20-C19	-2.13	106.95	110.72
3	B	800	7LN	C21-C20-C19	-2.11	106.99	110.72
2	A	600	H4B	C4A-C8A-N8	2.13	120.93	118.43
2	A	600	H4B	C2-N1-C8A	2.23	119.55	114.54
6	B	1483	HEM	CMD-C2D-C3D	2.37	124.82	114.35
6	B	1483	HEM	C2D-C3D-C4D	2.38	105.53	101.50
3	A	800	7LN	C11-N07-C06	2.41	127.30	121.46
6	A	1483	HEM	C2D-C3D-C4D	2.51	105.75	101.50
6	A	1483	HEM	CAA-CBA-CGA	2.55	117.42	112.75
6	A	1483	HEM	CMD-C2D-C3D	2.85	126.96	114.35
2	B	600	H4B	N2-C2-N3	2.94	122.07	117.20
2	B	600	H4B	C2-N1-C8A	3.17	121.68	114.54
2	B	600	H4B	C4-C4A-C8A	3.23	117.49	114.56
2	B	600	H4B	C4-N3-C2	3.32	120.55	115.94
6	B	1483	HEM	CMB-C2B-C3B	3.34	124.86	116.53
6	A	1483	HEM	CMB-C2B-C3B	3.95	126.40	116.53
6	B	1483	HEM	CAD-C3D-C2D	4.04	124.83	113.22
6	A	1483	HEM	CMC-C2C-C3C	4.18	126.96	116.53
6	B	1483	HEM	CMC-C2C-C3C	4.32	127.31	116.53
6	A	1483	HEM	CAD-C3D-C2D	4.43	125.95	113.22
6	A	1483	HEM	CAD-C3D-C4D	4.46	128.20	112.47
2	A	600	H4B	C4-N3-C2	4.58	122.30	115.94
2	A	600	H4B	C4-C4A-C8A	4.75	118.86	114.56
6	B	1483	HEM	CAD-C3D-C4D	4.87	129.64	112.47

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	800	7LN	N06-C06-N07-C11
3	A	800	7LN	N06-C06-N07-C11

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1483	HEM	1	0
2	A	600	H4B	1	0
3	A	800	7LN	2	0
6	B	1483	HEM	1	0
2	B	600	H4B	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/443 (91%)	0.63	52 (12%) 5 7	27, 38, 67, 115	0
1	B	402/443 (90%)	0.58	44 (10%) 7 11	26, 42, 72, 107	0
All	All	806/886 (90%)	0.60	96 (11%) 6 9	26, 40, 70, 115	0

All (96) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	6.8
1	B	259	GLN	6.3
1	A	160	ALA	6.2
1	A	109	ARG	6.2
1	A	259	GLN	6.0
1	A	121	PRO	5.7
1	A	91	GLN	5.5
1	A	122	ALA	5.5
1	A	108	PRO	5.5
1	B	143	SER	5.2
1	A	123	GLU	5.2
1	B	142	ARG	5.1
1	A	97	THR	5.1
1	B	121	PRO	4.8
1	B	261	GLY	4.8
1	B	98	PRO	4.7
1	A	69	LYS	4.6
1	B	146	GLN	4.6
1	A	92	GLN	4.5
1	B	99	ARG	4.5
1	A	67	GLY	4.3
1	A	98	PRO	4.2
1	A	338	VAL	4.0
1	B	123	GLU	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	239	GLY	4.0
1	A	126	LEU	3.9
1	A	159	VAL	3.8
1	B	122	ALA	3.7
1	A	99	ARG	3.6
1	A	260	ASP	3.6
1	B	93	ASP	3.6
1	A	124	GLN	3.6
1	B	94	GLY	3.6
1	B	223	ARG	3.5
1	A	238	PRO	3.3
1	B	260	ASP	3.3
1	B	109	ARG	3.3
1	A	156	GLU	3.3
1	A	153	GLN	3.2
1	B	338	VAL	3.2
1	B	141	LYS	3.2
1	B	144	GLY	3.2
1	B	258	GLN	3.1
1	B	145	SER	3.1
1	A	93	ASP	3.1
1	A	337	ALA	3.1
1	B	91	GLN	3.0
1	A	127	SER	3.0
1	B	414	LEU	2.9
1	B	96	CYS	2.9
1	A	230	ILE	2.9
1	A	101	CYS	2.9
1	A	261	GLY	2.9
1	B	97	THR	2.8
1	A	355	PHE	2.8
1	B	92	GLN	2.8
1	A	94	GLY	2.8
1	A	414	LEU	2.8
1	A	157	ALA	2.7
1	A	146	GLN	2.7
1	A	107	LEU	2.7
1	B	70	PHE	2.6
1	B	108	PRO	2.6
1	A	163	GLY	2.6
1	A	257	ARG	2.6
1	B	449	TRP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	95	PRO	2.5
1	B	69	LYS	2.5
1	A	155	VAL	2.5
1	A	161	SER	2.5
1	A	142	ARG	2.5
1	B	124	GLN	2.4
1	B	323	GLU	2.4
1	A	299	ASP	2.4
1	A	96	CYS	2.3
1	A	258	GLN	2.3
1	A	448	ALA	2.3
1	A	125	LEU	2.2
1	B	140	ILE	2.2
1	B	230	ILE	2.2
1	B	282	THR	2.2
1	B	147	ALA	2.2
1	B	448	ALA	2.2
1	A	339	SER	2.2
1	A	232	VAL	2.2
1	B	251	VAL	2.2
1	A	223	ARG	2.1
1	A	231	THR	2.1
1	A	449	TRP	2.1
1	B	337	ALA	2.1
1	B	339	SER	2.1
1	B	390	ARG	2.1
1	B	232	VAL	2.0
1	B	454	ILE	2.0
1	A	102	LEU	2.0
1	B	263	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CAS	B	384	9/10	0.96	0.09	-	49,52,65,68	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
1	CAS	A	384	9/10	0.97	0.09	-	36,37,57,61	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	ACT	B	860	4/4	0.95	0.16	6.40	47,50,50,51	0
5	GOL	A	880	6/6	0.89	0.29	4.13	54,60,61,61	0
5	GOL	B	880	6/6	0.90	0.25	2.87	50,58,60,60	0
3	7LN	B	800	36/36	0.91	0.27	2.61	29,67,146,148	0
3	7LN	A	800	36/36	0.93	0.26	1.67	26,73,135,139	0
4	ACT	B	861	4/4	0.97	0.21	1.57	40,40,42,42	0
4	ACT	A	861	4/4	0.97	0.17	1.12	33,34,34,36	0
6	HEM	A	1483	43/43	0.97	0.19	1.01	27,29,40,43	0
2	H4B	A	600	17/17	0.97	0.18	0.86	29,31,35,35	0
6	HEM	B	1483	43/43	0.98	0.15	0.57	28,31,36,41	0
2	H4B	B	600	17/17	0.98	0.16	0.26	28,30,34,35	0
4	ACT	A	860	4/4	0.98	0.08	-1.48	44,46,47,47	0
7	ZN	A	1484	1/1	1.00	0.06	-2.49	47,47,47,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.