



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:46 PM GMT

PDB ID : 3UYK
Title : Spinosyn Rhamnosyltransferase SpnG complexed with spinosyn aglycone
Authors : Isiorho, E.A.; Liu, H-W.; Keatinge-Clay, A.T.
Deposited on : 2011-12-06
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3190 (1.70-1.70)
Clashscore	102246	3585 (1.70-1.70)
Ramachandran outliers	100387	3527 (1.70-1.70)
Sidechain outliers	100360	3527 (1.70-1.70)
RSRZ outliers	91569	3200 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	 82% 13% • •
1	B	387	 80% 15% • 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OCX	B	387	-	-	-	X
4	BGC	B	388	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6079 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NDP-rhamnosyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	373	2780	1763	491	511	15	0	0	0
1	B	369	2747	1744	482	506	15	0	0	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	EXPRESSION TAG	UNP Q9ALM8
A	360	ALA	SER	SEE REMARK 999	UNP Q9ALM8
B	0	HIS	-	EXPRESSION TAG	UNP Q9ALM8
B	360	ALA	SER	SEE REMARK 999	UNP Q9ALM8

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is (2R,3AS,5AR,5BS,9S,13S,14R,16AS,16BR)-9-ETHYL-2,13-DIHYDROXY-14-METHYL-2,3,3A,5A,5B,6,9,10,11,12,13,14,16A,16B-TETRADECAHYDRO-1H-AS-INDACENO[3,2-D]OXACYCLODODECINE-7,15-DIONE (three-letter code: OCX) (formula: C₂₄H₃₄O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			29	24	5		

- Molecule 4 is SUGAR (BETA-D-GLUCOSE) (three-letter code: BGC) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	259	Total 259	O 259	0	0
5	B	251	Total 251	O 251	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NDP-rhamnosyltransferase



- Molecule 1: NDP-rhamnosyltransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	54.02Å 57.08Å 68.10Å 81.98° 74.10° 86.07°	Depositor
Resolution (Å)	29.83 – 1.70 29.83 – 1.70	Depositor EDS
% Data completeness (in resolution range)	96.9 (29.83-1.70) 93.4 (29.83-1.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.23 (at 1.71Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.189 , 0.234 0.188 , 0.233	Depositor DCC
R_{free} test set	4101 reflections (5.26%)	DCC
Wilson B-factor (Å ²)	18.4	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.3	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Outliers	0 of 82125 reflections	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6079	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BGC, 0CX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.19	5/2850 (0.2%)	1.11	9/3911 (0.2%)
1	B	1.15	5/2816 (0.2%)	1.17	13/3865 (0.3%)
All	All	1.17	10/5666 (0.2%)	1.14	22/7776 (0.3%)

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	320	TYR	CE1-CZ	8.91	1.50	1.38
1	A	142	TRP	CD2-CE2	8.71	1.51	1.41
1	A	112	TRP	CD2-CE2	5.68	1.48	1.41
1	B	142	TRP	CD2-CE2	5.66	1.48	1.41
1	B	112	TRP	CD2-CE2	5.65	1.48	1.41
1	A	210	TRP	CD2-CE2	5.58	1.48	1.41
1	A	340	SER	CA-CB	5.57	1.61	1.52
1	B	76	GLY	C-O	5.44	1.32	1.23
1	B	210	TRP	CD2-CE2	5.04	1.47	1.41
1	B	22	TRP	CD2-CE2	5.00	1.47	1.41

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	121	VAL	CG1-CB-CG2	-9.14	96.28	110.90
1	B	372	MET	CG-SD-CE	-9.14	85.58	100.20
1	A	364	LYS	CD-CE-NZ	-8.93	91.17	111.70
1	A	120	ASP	CB-CG-OD1	6.92	124.53	118.30
1	B	261	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	B	94	VAL	CG1-CB-CG2	-6.37	100.70	110.90
1	A	309	LEU	CB-CG-CD1	6.29	121.69	111.00
1	B	214	ARG	NE-CZ-NH1	6.22	123.41	120.30
1	A	314	TYR	CB-CG-CD1	6.20	124.72	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	141	ARG	NE-CZ-NH2	-5.99	117.30	120.30
1	B	265	THR	CA-CB-CG2	-5.89	104.16	112.40
1	A	120	ASP	CB-CG-OD2	-5.85	113.03	118.30
1	B	322	ARG	NE-CZ-NH1	-5.74	117.43	120.30
1	A	318	PHE	CB-CG-CD2	-5.72	116.79	120.80
1	B	261	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	B	214	ARG	NE-CZ-NH2	-5.59	117.51	120.30
1	B	247	LEU	CB-CG-CD1	-5.57	101.54	111.00
1	B	141	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	B	367	ASP	CB-CG-OD1	5.33	123.10	118.30
1	A	316	ASP	CB-CG-OD2	5.25	123.03	118.30
1	A	160	ASP	CB-CG-OD1	5.25	123.03	118.30
1	B	368	GLU	OE1-CD-OE2	-5.00	117.29	123.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2780	0	2774	35	0
1	B	2747	0	2742	39	0
2	A	1	0	0	0	0
3	B	29	0	34	5	0
4	B	12	0	12	0	0
5	A	259	0	0	7	0
5	B	251	0	0	4	0
All	All	6079	0	5562	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:175:GLU:HG2	1:A:385:THR:HG23	1.15	1.08
1:A:175:GLU:HG2	1:A:385:THR:CG2	1.89	1.01
1:A:113:ARG:HD2	5:A:599:HOH:O	1.61	1.01
1:B:196:VAL:HB	1:B:378:LEU:HD11	1.43	0.99
1:B:202:ASN:ND2	1:B:280:ASN:OD1	1.95	0.99
1:B:196:VAL:HB	1:B:378:LEU:CD1	2.00	0.91
1:A:202:ASN:OD1	1:A:280:ASN:OD1	1.88	0.91
1:A:107:ARG:HD3	5:A:485:HOH:O	1.77	0.84
1:B:196:VAL:CG2	1:B:378:LEU:HD12	2.10	0.82
1:A:175:GLU:CG	1:A:385:THR:HG23	2.07	0.80
1:B:90:THR:HG23	3:B:387:OCX:H34	1.68	0.76
1:A:53:ILE:O	1:A:54:ARG:HB2	1.86	0.75
1:B:94:VAL:CG2	3:B:387:OCX:H36	2.18	0.73
1:A:9:PRO:HB3	1:A:36:PRO:HG2	1.71	0.72
1:B:258:PRO:HA	1:B:275:GLU:OE2	1.91	0.70
1:B:256:VAL:HB	1:B:257:PRO:HD2	1.73	0.70
1:B:258:PRO:N	1:B:275:GLU:OE2	2.25	0.70
1:B:258:PRO:CA	1:B:275:GLU:OE2	2.42	0.68
1:A:259:GLU:N	1:A:259:GLU:OE1	2.28	0.66
1:B:257:PRO:C	1:B:275:GLU:OE2	2.34	0.66
1:B:294:GLY:O	1:B:298:ALA:HB3	1.97	0.65
1:B:380:ARG:HB3	5:B:414:HOH:O	1.96	0.64
1:A:79:ASP:HB2	5:A:653:HOH:O	1.98	0.62
1:B:342:HIS:O	1:B:346:THR:HG23	2.02	0.60
1:B:256:VAL:HB	1:B:257:PRO:CD	2.33	0.58
1:B:140:HIS:HE1	5:B:524:HOH:O	1.87	0.57
1:B:294:GLY:HA3	5:B:598:HOH:O	2.05	0.56
1:A:283:LEU:HD11	1:A:289:VAL:HG22	1.89	0.55
1:B:94:VAL:HG23	3:B:387:OCX:H36	1.89	0.54
1:B:196:VAL:CB	1:B:378:LEU:CD1	2.80	0.54
1:A:374:HIS:CE1	1:B:28:GLY:HA3	2.43	0.53
1:A:0:HIS:CE1	1:A:30:GLU:HG3	2.44	0.52
1:B:37:PRO:HD3	1:B:53:ILE:O	2.09	0.52
1:A:154:ARG:NH2	5:A:550:HOH:O	2.41	0.52
1:A:0:HIS:HE1	1:A:30:GLU:OE1	1.91	0.52
1:A:7:PRO:HG2	1:A:39:LEU:CD1	2.40	0.52
1:B:196:VAL:HG23	1:B:378:LEU:HD12	1.92	0.52
1:B:257:PRO:HG2	1:B:260:HIS:ND1	2.25	0.51
1:B:196:VAL:HB	1:B:378:LEU:HD12	1.91	0.51
1:A:294:GLY:N	5:A:415:HOH:O	2.44	0.51
1:A:104:GLU:HG3	1:A:107:ARG:NH2	2.26	0.51
1:A:215:THR:HG21	5:A:666:HOH:O	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:ARG:NH2	1:A:2:ARG:HG2	2.27	0.50
1:B:196:VAL:CB	1:B:378:LEU:HD12	2.41	0.49
1:B:110:GLU:OE2	1:B:113:ARG:NH1	2.45	0.49
1:A:343:GLU:HG2	5:A:536:HOH:O	2.13	0.49
1:B:12:THR:HG22	5:B:570:HOH:O	2.12	0.49
1:A:36:PRO:HA	1:A:53:ILE:O	2.13	0.49
1:A:226:ARG:NE	1:A:259:GLU:OE2	2.43	0.48
1:B:196:VAL:HG21	1:B:378:LEU:HD12	1.91	0.48
1:B:243:ALA:HA	1:B:346:THR:HG22	1.95	0.48
1:A:227:MET:HE3	1:A:227:MET:HA	1.94	0.47
1:A:54:ARG:N	1:A:104:GLU:OE2	2.40	0.47
1:B:180:PRO:HA	1:B:196:VAL:O	2.15	0.47
1:B:240:ALA:HA	1:B:345:PHE:CZ	2.50	0.46
1:B:305:GLY:HA3	1:B:363:ILE:HD13	1.98	0.45
1:A:160:ASP:N	1:A:161:PRO:CD	2.79	0.45
1:B:94:VAL:HG23	3:B:387:OCX:C2	2.47	0.44
1:B:27:SER:OG	1:B:379:VAL:HG11	2.17	0.44
1:A:298:ALA:HB1	1:A:320:TYR:CZ	2.52	0.44
1:B:9:PRO:HB3	1:B:36:PRO:HG2	2.01	0.43
1:A:54:ARG:HD3	1:A:54:ARG:HA	1.60	0.43
1:A:267:LEU:HD12	1:A:267:LEU:HA	1.91	0.43
1:A:38:GLU:N	1:A:38:GLU:OE2	2.44	0.42
1:B:283:LEU:HD11	1:B:289:VAL:HG22	2.00	0.42
1:A:8:LEU:HD12	1:A:13:HIS:CE1	2.54	0.42
1:A:269:ASP:N	1:A:269:ASP:OD1	2.52	0.42
1:B:298:ALA:HB1	1:B:320:TYR:CZ	2.55	0.42
1:B:182:PRO:HG2	1:B:185:LEU:HD12	2.02	0.41
1:B:94:VAL:CG2	3:B:387:OCX:C2	2.93	0.41
1:A:283:LEU:HD11	1:A:289:VAL:CG2	2.51	0.41
1:A:37:PRO:HD3	1:A:54:ARG:HB2	2.02	0.40
1:A:113:ARG:HG2	1:A:113:ARG:HH11	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	369/387 (95%)	359 (97%)	9 (2%)	1 (0%)	46	26
1	B	365/387 (94%)	356 (98%)	8 (2%)	1 (0%)	46	26
All	All	734/774 (95%)	715 (97%)	17 (2%)	2 (0%)	46	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	295	SER
1	A	146	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/300 (97%)	276 (95%)	14 (5%)	31	12
1	B	287/300 (96%)	278 (97%)	9 (3%)	47	25
All	All	577/600 (96%)	554 (96%)	23 (4%)	38	16

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	PRO
1	A	12	THR
1	A	40	GLN
1	A	54	ARG
1	A	69	ARG
1	A	113	ARG
1	A	202	ASN
1	A	227	MET
1	A	268	PRO
1	A	269	ASP
1	A	278	PRO

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	308	GLN
1	A	314	TYR
1	A	345	PHE
1	B	99	LEU
1	B	134	ASP
1	B	184	SER
1	B	279	LEU
1	B	314	TYR
1	B	345	PHE
1	B	364	LYS
1	B	380	ARG
1	B	382	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	0	HIS
1	A	25	GLN
1	A	96	GLN
1	A	166	HIS
1	A	197	GLN
1	A	230	ASN
1	A	308	GLN
1	A	313	GLN
1	A	374	HIS
1	B	25	GLN
1	B	140	HIS
1	B	166	HIS
1	B	202	ASN
1	B	280	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	0CX	B	387	-	30,32,32	1.53	6 (20%)	23,47,47	2.10	8 (34%)
4	BGC	B	388	-	12,12,12	1.09	1 (8%)	17,17,17	1.61	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0CX	B	387	-	-	0/28/65/65	0/3/4/4
4	BGC	B	388	-	-	0/2/22/22	0/1/1/1

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	387	0CX	C16-C15	-2.93	1.48	1.52
3	B	387	0CX	C3-C14	-2.45	1.47	1.51
3	B	387	0CX	O1-C1	2.13	1.40	1.34
3	B	387	0CX	C12-C13	2.65	1.53	1.50
3	B	387	0CX	C10-C9	2.72	1.58	1.52
4	B	388	BGC	O5-C1	2.85	1.48	1.43
3	B	387	0CX	C13-C14	5.23	1.39	1.33

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	387	OCX	C3-C2-C1	-4.20	107.38	113.76
3	B	387	OCX	O5-C17-C16	-2.89	101.21	109.06
4	B	388	BGC	C4-C3-C2	-2.69	105.78	110.79
3	B	387	OCX	O1-C1-O3	-2.68	116.47	123.67
3	B	387	OCX	C21-O1-C1	-2.12	112.79	117.89
3	B	387	OCX	C8-C7-C11	2.02	104.78	100.48
4	B	388	BGC	C1-C2-C3	2.73	114.49	110.43
4	B	388	BGC	O5-C5-C4	3.00	115.30	109.68
4	B	388	BGC	O5-C1-C2	3.31	115.08	109.80
3	B	387	OCX	C12-C4-C5	3.67	118.41	113.16
3	B	387	OCX	O4-C15-C14	3.96	126.06	120.35
3	B	387	OCX	O1-C1-C2	4.05	119.38	111.54

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	387	OCX	5	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/387 (96%)	-0.11	3 (0%) 87 90	9, 18, 36, 61	0
1	B	369/387 (95%)	-0.12	2 (0%) 91 93	10, 19, 31, 55	0
All	All	742/774 (95%)	-0.12	5 (0%) 89 91	9, 19, 34, 61	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	294	GLY	3.9
1	A	10	TYR	3.3
1	A	266	ASP	3.0
1	A	54	ARG	2.6
1	B	380	ARG	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	BGC	B	388	12/12	0.89	0.15	5.06	16,24,41,57	0
3	OCX	B	387	29/29	0.79	0.21	4.71	31,36,39,43	0
2	MG	A	387	1/1	0.99	0.04	-1.82	11,11,11,11	0

6.5 Other polymers [i](#)

There are no such residues in this entry.