Full wwPDB X-ray Structure Validation Report (i)

Feb 1, 2016 – 09:08 PM GMT

PDB ID	:	4UZN
Title	:	The native structure of the family 46 carbohydrate-binding module (CBM46) of endo-beta-1,4-glucanase B (Cel5B) from Bacillus halodurans
Authors	:	Venditto, I.; Santos, H.; Ferreira, L.M.A.; Sakka, K.; Fontes, C.M.G.A.; Najmudin, S.
Deposited on	:	2014-09-05
Resolution	:	2.46 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.46 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4776 (2.50-2.42)
Clashscore	102246	1030 (2.48-2.44)
Ramachandran outliers	100387	1024 (2.48-2.44)
Sidechain outliers	100360	1024 (2.48-2.44)
RSRZ outliers	91569	4787 (2.50-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	3%, 75%, 5% •, 19%
1	B	130	18%, 75%, 6%, 19%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1669 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ENDO-BETA-1,4-GLUCANASE (CELULASE B).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	105	816	518	135	161	2	0	0	0
1	B	105	830	528	139	161	2	0	2	0

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	436	MET	-	EXPRESSION TAG	UNP Q9KF82
A	437	GLY	-	EXPRESSION TAG	UNP Q9KF82
A	438	SER	-	EXPRESSION TAG	UNP Q9KF82
A	439	SER	-	EXPRESSION TAG	UNP Q9KF82
A	440	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	441	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	442	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	443	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	444	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	445	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	446	SER	-	EXPRESSION TAG	UNP Q9KF82
A	447	SER	-	EXPRESSION TAG	UNP Q9KF82
A	448	GLY	-	EXPRESSION TAG	UNP Q9KF82
A	449	LEU	-	EXPRESSION TAG	UNP Q9KF82
A	450	VAL	-	EXPRESSION TAG	UNP Q9KF82
A	451	PRO	-	EXPRESSION TAG	UNP Q9KF82
A	452	ARG	-	EXPRESSION TAG	UNP Q9KF82
A	453	GLY	-	EXPRESSION TAG	UNP Q9KF82
A	454	SER	-	EXPRESSION TAG	UNP Q9KF82
A	455	HIS	-	EXPRESSION TAG	UNP Q9KF82
A	456	MET	-	EXPRESSION TAG	UNP Q9KF82
A	457	ALA	-	EXPRESSION TAG	UNP Q9KF82
A	458	SER	-	EXPRESSION TAG	UNP Q9KF82
B	436	MET	-	EXPRESSION TAG	UNP Q9KF82
B	437	GLY	-	EXPRESSION TAG	UNP Q9KF82

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	438	SER	-	EXPRESSION TAG	UNP Q9KF82
B	439	SER	-	EXPRESSION TAG	UNP Q9KF82
B	440	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	441	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	442	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	443	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	444	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	445	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	446	SER	-	EXPRESSION TAG	UNP Q9KF82
B	447	SER	-	EXPRESSION TAG	UNP Q9KF82
B	448	GLY	-	EXPRESSION TAG	UNP Q9KF82
B	449	LEU	-	EXPRESSION TAG	UNP Q9KF82
B	450	VAL	-	EXPRESSION TAG	UNP Q9KF82
B	451	PRO	-	EXPRESSION TAG	UNP Q9KF82
B	452	ARG	-	EXPRESSION TAG	UNP Q9KF82
B	453	GLY	-	EXPRESSION TAG	UNP Q9KF82
B	454	SER	-	EXPRESSION TAG	UNP Q9KF82
B	455	HIS	-	EXPRESSION TAG	UNP Q9KF82
B	456	MET	-	EXPRESSION TAG	UNP Q9KF82
B	457	ALA	-	EXPRESSION TAG	UNP Q9KF82
B	458	SER	-	EXPRESSION TAG	UNP Q9KF82

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	12	12	12	0	0
			Total	O		
2	B	11	11	11	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ENDO-BETA-1,4-GLUCANASE (CELULASE B)

Chain A:



- Molecule 1: ENDO-BETA-1,4-GLUCANASE (CELULASE B)

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α, β, γ	121.19Å 121.19Å 77.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	85.69 – 2.46 44.37 – 2.46	Depositor EDS
% Data completeness (in resolution range)	99.4 (85.69-2.46) 99.5 (44.37-2.46)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.42 (at 2.45Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.214 , 0.240 0.223 , 0.245	Depositor DCC
R_{free} test set	523 reflections (5.14%)	DCC
Wilson B-factor (Å^2)	56.7	Xtriage
Anisotropy	0.555	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.39 , 65.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	5 of 10697 reflections (0.047%)	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	1669	wwPDB-VP
Average B, all atoms (Å^2)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.98% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.64	0/838	0.70	0/1139
1	B	0.61	0/858	0.74	0/1164
All	All	0.63	0/1696	0.72	0/2303

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	816	0	755	3	1
1	B	830	0	781	4	0
2	A	12	0	0	0	0
2	B	11	0	0	0	0
All	All	1669	0	1536	7	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (7) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:463:VAL:HG12	1:B:479:SER:HB2	1.74	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:463:VAL:HG12	1:A:479:SER:HB2	1.76	0.67
1:B:468:GLN:HE21	1:B:561:THR:CG2	2.25	0.49
1:B:468:GLN:HE21	1:B:561:THR:HG22	1.78	0.49
1:A:478:ALA:O	1:A:520:ASN:HB3	2.15	0.46
1:B:478:ALA:O	1:B:520:ASN:HB3	2.16	0.44
1:A:547:GLU:HG2	1:A:548:THR:N	2.34	0.42

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:484:SER:OG	1:A:484:SER:OG[10_555]	1.84	0.36

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/130 (79%)	96 (93%)	7 (7%)	0	100	100
1	B	105/130 (81%)	97 (92%)	8 (8%)	0	100	100
All	All	208/260 (80%)	193 (93%)	15 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/107 (80%)	84 (98%)	2 (2%)	58	74
1	B	88/107 (82%)	85 (97%)	3 (3%)	44	61
All	All	174/214 (81%)	169 (97%)	5 (3%)	58	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	468	GLN
1	A	484	SER
1	B	484	SER
1	B	523[A]	LYS
1	B	523[B]	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	468	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å^2)	Q<0.9
1	A	105/130 (80%)	0.56	4 (3%) 44 48	50, 68, 99, 103	0
1	B	105/130 (80%)	1.09	23 (21%) 1 1	49, 67, 103, 114	0
All	All	210/260 (80%)	0.82	27 (12%) 5 4	49, 68, 101, 114	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	482	GLY	5.3
1	B	480	PHE	4.5
1	B	461	THR	4.4
1	B	530[A]	ARG	4.0
1	B	543	PHE	3.5
1	B	546	GLY	3.4
1	B	548	THR	3.3
1	B	518	ASP	3.2
1	B	547	GLU	3.2
1	B	544	TRP	3.1
1	B	545	SER	3.1
1	B	549	VAL	3.0
1	A	546	GLY	2.9
1	B	481	ASN	2.9
1	A	565	ALA	2.8
1	B	558	ASN	2.8
1	B	463	VAL	2.8
1	B	462	PRO	2.8
1	B	487	THR	2.7
1	B	471	VAL	2.7
1	B	559	GLN	2.6
1	B	483	ASN	2.6
1	B	557	GLY	2.5
1	B	542	HIS	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	464	LEU	2.4
1	A	478	ALA	2.2
1	B	478	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.