



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 05:50 AM GMT

PDB ID : 2V0O
Title : FCHO2 F-BAR DOMAIN
Authors : Henne, W.M.; McMahon, H.T.; Kent, H.M.; Evans, P.R.
Deposited on : 2007-05-15
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	<div> <div>7%</div> <div>95%</div> <div>• •</div> </div>
1	B	276	<div> <div>14%</div> <div>92%</div> <div>6% •</div> </div>
1	C	276	<div> <div>•</div> <div>91%</div> <div>7% •</div> </div>

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 6757 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FCH DOMAIN ONLY PROTEIN 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	0	0
			2196	1383	373	432	8			
1	B	271	Total	C	N	O	S	0	0	0
			2180	1374	371	427	8			
1	C	272	Total	C	N	O	S	0	0	0
			2189	1378	372	431	8			

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	63	Total 63	O 63	0	0
3	B	31	Total 31	O 31	0	0
3	C	86	Total 86	O 86	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: FCH DOMAIN ONLY PROTEIN 2



• Molecule 1: FCH DOMAIN ONLY PROTEIN 2



• Molecule 1: FCH DOMAIN ONLY PROTEIN 2



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	254.44Å 65.68Å 89.91Å 90.00° 110.32° 90.00°	Depositor
Resolution (Å)	119.52 – 2.30 39.56 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.3 (119.52-2.30) 97.5 (39.56-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.278 , 0.316 0.276 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	43.0	Xtriage
Anisotropy	0.590	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 27.8	EDS
Estimated twinning fraction	0.136 for -h-2*l,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 61798 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6757	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 25.19 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.3225e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.30	0/2230	0.40	0/2989
1	B	0.31	0/2213	0.40	0/2967
1	C	0.31	0/2222	0.42	0/2978
All	All	0.31	0/6665	0.41	0/8934

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2196	0	2194	4	0
1	B	2180	0	2182	7	0
1	C	2189	0	2188	11	0
2	A	4	0	3	0	0
2	B	4	0	3	0	0
2	C	4	0	3	0	0
3	A	63	0	0	0	0
3	B	31	0	0	0	0
3	C	86	0	0	0	0
All	All	6757	0	6573	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:98:ILE:HG22	1:C:102:GLN:HE21	1.61	0.65
1:B:61:SER:HB3	1:B:77:LYS:HD2	1.84	0.58
1:A:52:SER:HB3	1:B:45:ILE:HG12	1.92	0.51
1:C:53:MET:HE2	1:C:83:LEU:HD23	1.94	0.48
1:C:53:MET:HE3	1:C:80:THR:HG23	1.94	0.48
1:A:95:GLN:HA	1:A:98:ILE:HD12	1.96	0.47
1:C:47:GLU:O	1:C:51:ARG:HG2	2.15	0.47
1:B:3:MET:HG2	1:B:4:ALA:H	1.80	0.46
1:B:87:HIS:O	1:B:91:VAL:HG23	2.15	0.46
1:C:246:THR:HG23	1:C:249:SER:H	1.81	0.46
1:B:95:GLN:HA	1:B:98:ILE:HD12	1.98	0.45
1:C:263:GLU:H	1:C:263:GLU:CD	2.20	0.45
1:C:95:GLN:HA	1:C:98:ILE:HD12	1.99	0.45
1:C:87:HIS:O	1:C:91:VAL:HG23	2.17	0.45
1:B:160:GLN:HA	1:B:163:ILE:HD12	1.99	0.44
1:A:246:THR:HG23	1:A:249:SER:H	1.82	0.44
1:C:53:MET:CE	1:C:80:THR:HG23	2.49	0.42
1:C:36:ALA:O	1:C:40:ARG:HB2	2.20	0.40
1:C:101:VAL:HG22	1:C:216:ILE:HG21	2.03	0.40
1:A:45:ILE:HG12	1:B:52:SER:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	271/276 (98%)	269 (99%)	2 (1%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	269/276 (98%)	264 (98%)	5 (2%)	0	100	100
1	C	270/276 (98%)	266 (98%)	4 (2%)	0	100	100
All	All	810/828 (98%)	799 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/240 (99%)	232 (98%)	6 (2%)	55	73
1	B	236/240 (98%)	231 (98%)	5 (2%)	61	78
1	C	237/240 (99%)	234 (99%)	3 (1%)	76	87
All	All	711/720 (99%)	697 (98%)	14 (2%)	63	79

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	GLN
1	A	68	THR
1	A	150	GLN
1	A	179	LEU
1	A	252	GLN
1	A	258	LYS
1	B	9	ASN
1	B	26	LYS
1	B	179	LEU
1	B	258	LYS
1	B	263	GLU
1	C	68	THR
1	C	150	GLN
1	C	179	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	23	HIS
1	A	65	GLN
1	A	127	GLN
1	A	134	GLN
1	B	112	HIS
1	C	23	HIS
1	C	102	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	1275	-	1,3,3	1.13	0	0,3,3	0.00	-
2	ACT	B	1275	-	1,3,3	1.15	0	0,3,3	0.00	-
2	ACT	C	1275	-	1,3,3	1.00	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ACT	A	1275	-	-	0/0/0/0	0/0/0/0
2	ACT	B	1275	-	-	0/0/0/0	0/0/0/0
2	ACT	C	1275	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/276 (98%)	0.81	20 (7%) 18 25	37, 63, 99, 109	0
1	B	271/276 (98%)	1.07	39 (14%) 3 5	39, 77, 110, 114	0
1	C	272/276 (98%)	0.51	4 (1%) 76 81	33, 53, 84, 89	0
All	All	816/828 (98%)	0.80	63 (7%) 16 23	33, 62, 103, 114	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163	ILE	6.9
1	B	151	GLU	6.4
1	B	152	ARG	5.7
1	B	186	LEU	5.2
1	B	155	LYS	5.0
1	C	160	GLN	4.3
1	A	161	ARG	4.0
1	A	155	LYS	3.8
1	B	156	GLU	3.7
1	B	160	GLN	3.6
1	A	163	ILE	3.6
1	B	168	VAL	3.5
1	A	270	PHE	3.4
1	A	164	GLU	3.4
1	B	6	PHE	3.3
1	A	267	LEU	3.1
1	B	129	ILE	3.1
1	B	167	ALA	3.0
1	B	159	THR	3.0
1	B	220	LEU	2.9
1	A	182	GLU	2.9
1	A	186	LEU	2.8
1	B	193	GLN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	158	ALA	2.8
1	B	179	LEU	2.8
1	B	210	LEU	2.7
1	A	160	GLN	2.7
1	B	97	LEU	2.6
1	B	213	ILE	2.6
1	A	274	ASP	2.6
1	B	199	ALA	2.5
1	B	21	LEU	2.5
1	B	35	LEU	2.5
1	B	153	LEU	2.4
1	B	83	LEU	2.4
1	C	182	GLU	2.4
1	B	123	LEU	2.4
1	A	263	GLU	2.4
1	B	165	LYS	2.3
1	A	158	ALA	2.3
1	B	132	ILE	2.3
1	A	66	LEU	2.3
1	B	148	VAL	2.3
1	A	251	ILE	2.3
1	A	83	LEU	2.3
1	B	177	TYR	2.3
1	B	189	ALA	2.3
1	B	30	ILE	2.2
1	A	152	ARG	2.2
1	B	98	ILE	2.1
1	A	273	CYS	2.1
1	B	205	ILE	2.1
1	A	151	GLU	2.1
1	B	32	THR	2.1
1	A	172	LYS	2.1
1	C	163	ILE	2.1
1	B	164	GLU	2.0
1	B	229	LEU	2.0
1	A	177	TYR	2.0
1	B	181	VAL	2.0
1	B	196	THR	2.0
1	B	101	VAL	2.0
1	C	220	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ACT	C	1275	4/4	0.88	0.19	0.32	67,67,67,67	0
2	ACT	B	1275	4/4	0.96	0.20	0.12	65,65,65,65	0
2	ACT	A	1275	4/4	0.94	0.10	-2.78	77,77,77,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.